Otology for Audiology

Otology and Audiology

Speak One Language

“Two Specialties. One Patient. One Language.”

◆

Dr. Yahia Al Ahmari

Otology, Neurotology & Cochlear Implant Leader

Dr. Hany Abdou

Audio-Vestibular Medicine

بسم الله الرحمن الرحيم

وَقُل رَّبِّ زِدْنِي عِلْمًا

A Decision That Could Not Be Delayed

The clinic was already behind schedule.

A child sat quietly on the edge of the chair, feet not touching the floor.

The parents sat closer than necessary, holding a file that felt heavier than paper.

Inside the file:

- *An audiogram labeled “severe to profound”*
- *ABR results marked “consistent with auditory neuropathy”*
- *Imaging reported as “structurally unremarkable”*
- *A previous note that read:*
  “Consider cochlear implantation.”

No urgency in the wording.

No hesitation in the conclusion.

Just enough confidence to move forward.

The otologist reviewed the images.

The anatomy was acceptable.



The cochlea looked navigable.

Nothing surgical stood in the way.

The audiologist reviewed the data.

The responses were present—but unstable.

Repeatability was questionable.

Functional access was unclear.

Both were correct.

Both were experienced.

Both cared about the same child.

Yet the room felt uneasy.

The parents asked a simple question:

"So… what do we do now?"

Not:

- *What does the test show?*
- *What does the guideline say?*
- *What is the next step in the protocol?*

But:

"What do we do now?"

Silence followed.



Not because no one knew the data

but because data alone could not answer the question.

A cochlear implant was possible.

Waiting was defensible.

More testing could be justified.

Referral could be delayed—or expedited.

Every option had a rationale.

Every option carried risk.

And whatever was chosen

would live with the child long after this meeting ended.

The decision would not be remembered as:

- An audiogram
- An ABR waveform
- A radiology report

It would be remembered as:

- Speech—or silence
- Integration—or isolation
- Opportunity—or regret



This moment did not require more tests.

It required shared understanding.

Understanding of:

- *What the data truly meant*
- *What surgery could—and could not—guarantee*
- *What delay might cost*
- *What intervention might risk*

It required otology and audiology

to speak one language.

This book begins here.

Not with certainty.

Not with protocols.

But with a decision that could not be delayed—

and a patient who would live with its consequences.



“Before guidelines, before protocols,

there is a decision.”

“Why Otology and Audiology Must Speak One Language”

There has always been an invisible line in hearing care.

A line not drawn by science—but by training paths, silos, and assumptions.

On one side stands Otology—

the world of anatomy, surgery, pathology, and decisive intervention.

On the other stands Audiology—

the world of measurement, interpretation, functional outcomes, and lifelong follow-up.

Both serve the same patient.

Both aim for the same goal.

Yet too often, they speak different languages.

This divide has cost us clarity.

It has delayed decisions.

It has created misunderstandings—not out of ego, but out of perspective.

And the patient stands in the middle.

This Project Was Born From Reality — Not Theory

This is not a traditional textbook.

It is not written to impress.

It is written because clinical reality demands it.

Every cochlear implant discussion.

Every borderline candidacy.

Every conflicting report.

Every moment where one specialty asks:

"Why did they decide this way?"

This project exists to answer that question.

Otology & Audiology
One Mission.

— This Book

Not to teach audiologists surgery—

but to teach them how otologists think.

How surgical risk is weighed.

How anatomy changes decisions.

How timing, prognosis, and feasibility are judged.

How “possible” is different from “appropriate”.



Otology without audiology is blind to function.

Audiology without otology is blind to structure.

True excellence lives between the two.

Dr. Yahia Al Ahmari

Otology, Neurotology & Cochlear Implant Leader

Dr. Hany Abdou

Audio-Vestibular Medicine

The Philosophy Behind The Books

- ***Evidence before opinion***
- ***Clinical reasoning before protocols***
- ***Outcomes before numbers***
- ***Communication before conflict***
- ***The patient before the system***

Each chapter ends with a distilled clinical insight not found in guidelines,but learned at the intersection of experience, evidence, and responsibility.

Our Commitment

We commit to honesty—even when uncomfortable.

We commit to clarity—even when complex.

We commit to respect—across disciplines.

And above all, we commit to the patient who trusts both of us.

These books are written for:

- ***Audiologists who want to think beyond the booth***
- ***Otologists who want to hear beyond the audiogram***
- ***Residents, fellows, and consultants who refuse fragmented care***

“Two Specialties. One Patient. One Language.”

- *Where Otology Meets Audiology*
- *Clinical Thinking Beyond Boundaries*
- *From Anatomy to Function — Together*

How Otologic Surgeons Think, Decide, and Judge Risk

How Audiologists Interpret Data, Define Function, and Guide Decisions

The Shared Language Series is not designed to teach one specialty the tools of the other.

It is designed to teach how each specialty thinks.

Because better outcomes begin when experts understand not only the data—but the reasoning behind it.

Dr. Yahia Al Ahmari & Dr. Hany Abdou

A Final Word

The future of hearing care will not be led by one specialty.

It will be led by those who understand both.

This is our contribution to that future.

"Medicine does not fail when knowledge is missing.
It fails when knowledge is fragmented."
"When experts stop listening to each other,
patients pay the price

Final Clinical Truth

“The most dangerous clinician is not the one who lacks knowledge,
but the one who believes that a test result is a decision.”

“Audiograms do not treat patients.
Surgeries do not define outcomes.
Reports do not replace judgment.”

“Hearing care succeeds only when structure, function, data, and ethics
are interpreted by a clinician who understands their limits.”

“If Otology forgets function, it risks harm.
If Audiology forgets anatomy, it risks illusion.”



“The patient does not care who is right.
The patient lives with the outcome.”

“This is why Otology and Audiology must speak one language.”

“Skipping this step saves time—but costs accuracy.”
“Assuming this saves money—but costs trust.”

“Most clinical errors are not due to ignorance, but to comfort.”

“We rarely make the wrong decision—we make the easy one.”

“If a test makes everyone feel safe, question whether it made anyone wiser.”



“Reports don’t document reality—they create it.”

“A single careless sentence can undo a hundred correct tests.”
“Write as if this sentence will be read in court.”



“Miscommunication is the most expensive complication.”

“Different specialties don’t disagree on facts—they disagree on meaning.”

“If you don’t translate your data, someone else will misinterpret it.”

“A good clinician plans success. A great one anticipates failure.”

“Most disasters were once ‘acceptable risks’.”

“The earlier you imagine failure, the safer the patient becomes.”

“If you can’t explain it to a family, you don’t fully understand it.”

“Families don’t need certainty—they need honesty.”
“Trust is built by clarity, not confidence.”

“Clinical excellence is not about knowing more.
It is about deciding better—under pressure.”

“Clinical excellence is not about knowing more.
It is about deciding better—under pressure.”



Preface

Before You Read Another Chapter

This book was not written to add information.

The world does not suffer from lack of information.

It suffers from fragmented thinking.

In modern hearing care, knowledge is everywhere.

Guidelines are accessible.

Protocols are published.

Devices are advancing.

Yet patients still fall between decisions.

Not because clinicians are careless—

but because decisions are often made in isolation.

The Reality We Face in the Clinic

Every week, we encounter the same patterns:

- ***Reports that are technically correct, yet clinically misleading***
- ***Surgical plans that ignore functional hearing reality***

- Audiological results interpreted without anatomical context
- Referrals made too early, or far too late
- Families confused by confident but conflicting opinions

None of this comes from bad intentions.

It comes from specialties speaking past each other.

Why This Book Exists

This book exists because:

- ***An audiogram is not a decision***
- ***A surgical option is not an outcome***
- ***A report is not neutral***
- ***And silence between specialties is not harmless***

Hearing care does not fail at the level of effort.

It fails at the level of understanding.

What This Book Is — and Is Not

This book is not:

- A guideline replacement
- A protocol checklist
- A showcase of devices or techniques
- A defense of one specialty over another

This book is:

- A map of clinical thinking
- A translation between disciplines
- A guide to safer decisions
- A reminder of professional responsibility

One Voice. One Philosophy.

Although written from two disciplines,

this book speaks with one voice.

A voice that values:

- Judgment over numbers
- Context over convenience
- Communication over confidence
- And outcomes over opinions

Every chapter is written with the same mindset:

"If this decision were about someone we love, would it still be acceptable?"

At the end of every chapter, you will find the Framework

Not a summary.

Not a revision tool.

But a clinical closure:

- *Where mistakes are named*
- *Where risks are exposed*
- *Where ethics are addressed*
- *Where decisions are owned*

This framework exists because real learning happens after the chapter ends.

How to Read This Book

Do not rush.

This book is not meant to be consumed—it is meant to be engaged.

- ***Pause at the uncomfortable truths***
- ***Re-read the dangerous sentences***
- ***Argue with the decisions***
- ***Translate the language into your own practice***

If a chapter makes you uncomfortable,

it is doing its job.

A Shared Responsibility

Otology and Audiology do not compete.

They complete each other.

The patient does not benefit from who is right.

The patient lives with what was decided.

This book is our attempt to narrow that gap.

Not with louder voices—

but with a shared language.



A Final Note

If this book changes the way you:

- ***Write one report***
- ***Explain one decision to a family***
- ***Pause before one irreversible step***

Then it has succeeded.

Because in hearing care,
the smallest decision can echo for a lifetime.

Dr. Yahia Al Ahmari

Otology, Neurotology & Cochlear Implant Leader

Dr. Hany Abdou

Audio-Vestibular Medicine

THE SHARED LANGUAGE SERIES

Bridging Otology & Audiology in Modern Hearing Care

Otology for Audiology

How Otologic Surgeons Think, Decide, and Judge Risk

AND

Audiology for Otology

How Audiologists Interpret Data, Define Function, and Guide Decisions

Clinical Reference • Educational Framework • Professional Responsibility

Two Volumes. One Clinical Language.

Dr. Yahia Al Ahmari

Otology & Cochlear Implant Surgery

Dr. Hany Abdou

Audiology & Audio-Vestibular Medicine



How to Use This Book

A Guide for the Clinician, Not the Reader

This book is not meant to be read passively.

It is meant to be used.

It was written for moments of uncertainty,

for decisions made under pressure,

and for clinicians who understand that knowing is not the same as deciding.

Read With a Clinical Mindset

Each chapter is structured to mirror real clinical thinking:

- *A problem appears*
- *Data is gathered*
- *Interpretation follows*
- *A decision must be made*
- *Responsibility is owned*

Do not skip to conclusions.

The value of this book lies in how conclusions are reached.

Use the Chapters as Decision Tools

You can approach this book in multiple ways:

- *Sequential reading to build a shared philosophy*
- *Problem-based reading when facing a real patient*
- *Selective reading to understand how the other specialty thinks*

Each chapter stands alone—

but together, they form a unified clinical framework.

Pause at the Framework

The end of each chapter contains the Framework™.

This is not a summary.

It is the most important part of the chapter.

Use it to:

- ***Re-check your assumptions***
- ***Identify blind spots***
- ***Anticipate errors***
- ***Strengthen documentation***
- ***Prepare for discussion with colleagues or families***

If time is limited, read the chapter—

then read the Framework twice.

Engage With the Discomfort

Some sections are intentionally uncomfortable.

They expose:

- ***Common but unspoken mistakes***
- ***Medico-legal vulnerabilities***
- ***Miscommunication between specialties***
- ***Decisions made for convenience rather than clarity***

If a section challenges your routine practice,

pause.

That moment is where learning happens.

Translate, Don’t Imitate

This book does not offer templates to copy blindly.

Instead, it offers:

- ***Ways of thinking***
- ***Patterns of judgment***
- ***Questions to ask before acting***

Translate the principles into your own clinical environment,

your own regulations,

and your own scope of practice.

Use It as a Shared Reference

This book is designed to be used:

- ***In multidisciplinary meetings***
- ***During case discussions***
- ***In training and mentoring***
- ***As a reference for report writing and referral decisions***

When both specialties read from the same page,

disagreements become productive, not personal.

A Final Instruction

Do not ask:

“What does this book say I should do?”

Ask instead:

“How does this book change the way I think?”

If it sharpens your judgment,

clarifies your communication,

and protects your patient—

then it has done its job.

About the Authors

Dr. Yahia Al Ahmari

Otology & Cochlear Implant Leader

Dr. Yahia Al Ahmari is a leading otologic surgeon and a recognized figure in cochlear implant practice and advanced ear surgery.

His work is grounded in a deep understanding of anatomy, surgical feasibility, and long-term patient outcomes.

Beyond the operating room, he is known for his precise clinical judgment knowing not only what can be done, but what should be done, and when restraint is the safest decision.

Throughout his clinical career, Dr. Al Ahmari has been actively involved in multidisciplinary decision-making, particularly at the interface between otology and audiology.

His experience has highlighted a recurring challenge: technically sound data that fails to translate into shared understanding or optimal outcomes.

This book reflects his belief that successful otologic care depends not only on surgical skill, but on clear communication, realistic expectations, and a shared clinical language with audiology.

Dr. Hany Abdou

Audio-Vestibular Medicine

Dr. Hany Abdou is an audiology physician specializing in audio-vestibular medicine, with extensive experience in diagnostic audiology, clinical decision-making, and evidence-based report writing.

His work focuses on transforming test results into meaningful clinical conclusions—bridging the gap between measurements, function, and real-life patient impact. He is particularly known for emphasizing clinical reasoning, documentation integrity, and medico-legal awareness in audiological practice.

Dr. Abdou's professional interest lies in redefining how audiology is practiced, taught, and communicated—moving beyond isolated test interpretation toward integrated, patient-centered decision-making that supports surgical, rehabilitative, and long-term care pathways.

This book reflects his commitment to making audiology not just accurate, but clinically decisive, communicative, and accountable.

A Shared Perspective

Although trained in different disciplines, the authors share a common philosophy:

- ***That no single test defines a patient***
- ***That no single specialty owns the outcome***
- ***That responsibility lies not in data, but in decisions***

This collaboration was born not from theory, but from real clinical encounters—where better outcomes depended on better understanding.

Together, they write not as representatives of two specialties,
but as clinicians committed to one patient.



Acknowledgments

This work did not emerge from isolation.

It was shaped by conversations, disagreements, shared cases, and moments of uncertainty—

the kind that define real clinical practice.

We acknowledge, first and foremost, the patients and families whose trust carries the greatest responsibility.

Every difficult decision, every unresolved question, and every long discussion behind closed doors has left a mark on how these pages were written.

They are the silent contributors to every chapter.

We are deeply grateful to the clinicians—otologists, audiologists, residents, fellows, nurses, and technicians—who practice with honesty and humility.

Those who ask uncomfortable questions.

Those who challenge assumptions.

Those who choose clarity over convenience.

Their daily work in clinics, operating rooms, and multidisciplinary meetings exposed the gaps this book attempts to address.

We acknowledge the mentors who taught us that clinical excellence is not measured by confidence alone, but by restraint, reflection, and

accountability.

Their influence is present not in citations, but in judgment.

We also recognize colleagues who may not always agree with every position presented here.

Disagreement, when grounded in respect and shared purpose, strengthens medicine rather than weakens it.

This book welcomes that dialogue.

Finally, we acknowledge the families, colleagues, and teams who support clinical work beyond visible hours.

Those who understand that decisions made during the day often follow clinicians home at night.

This book is not a declaration of authority.

It is a reflection of responsibility—

shared, debated, and continuously evolving.

If these pages help even one clinician pause, reconsider, or communicate more clearly,

then everyone acknowledged here has contributed to that outcome.

Dr. Yahia Al Ahmari

Otology & Cochlear Implant Surgery

Dr. Hany Abdou

Audiology & Audio-Vestibular Medicine

Dedications

To the patients

who trusted us at moments when certainty was impossible,
and whose lives were shaped by decisions made in minutes—
yet carried for a lifetime.

To the families

who asked the hardest questions,
waited through the longest silences,
and reminded us that behind every test, report, and procedure
stands a human story.

To the clinicians

who choose to think before acting,
to question before concluding,
and to place responsibility above comfort.
Those who understand that
clinical excellence is not loud,
but precise.

To the mentors

who taught us that experience without reflection is dangerous,
and that restraint is often the highest form of expertise.

And to every otologist and audiologist

who believes that better outcomes begin
when we listen—to data, to colleagues, and to each other.
This work is dedicated to a shared language,
a shared responsibility,
and the patient who depends on both.

"To the patient who lives with our decisions long after we leave the clinic."

"Because the patient remembers outcomes, not specialties."

"Every decision echoes beyond the clinic."

"Clinical decisions end in pages — but live in people."

Dr. Yahia Al Ahmari

Dr. Hany Abdou

Disclaimer

This book is an educational and professional reference designed to support clinical thinking, communication, and decision-making at the interface of otology and audiology.

It is not intended to replace:

- *Individual clinical judgment*
- *Institutional policies or regulations*
- *Local laws, licensing requirements, or scope-of-practice rules*
- *Formal training, supervision, or credentialing*

Clinical decisions must always be made in the context of:

- *The individual patient*
- *The clinician's qualifications and privileges*
- *Available resources and institutional guidelines*

All clinical actions, interpretations, and decisions must comply with:

- *Regulations of the Saudi Ministry of Health (MOH)*

- Requirements of the Saudi Commission for Health Specialties (SCFHS)
- Institutional bylaws, credentialing processes, and approved clinical policies

References to medical devices, software platforms, manufacturers, or commercial products are provided solely for educational illustration.

Such references do not constitute endorsement, recommendation, sponsorship, or preferential promotion of any company, product, or service.

Scope of practice, credentialing, and clinical privileges vary between institutions and jurisdictions.

Any examples, templates, or policy models included are provided for reference only and must not be implemented without formal institutional approval.

The authors assume no responsibility for clinical decisions, patient outcomes, or professional actions resulting from the use or misuse of this content.

Ultimate responsibility for patient care rests with the licensed practitioner acting within approved scope and privileges.

Although every effort has been made to ensure accuracy, clarity, and alignment with current evidence-based and international guidelines at the time of publication, medicine is a constantly evolving field.

Recommendations, standards, and best practices may change.

The authors do not assume responsibility for:

- ***Misinterpretation or misuse of the information presented***
- ***Decisions made outside the clinician's scope of practice***
- ***Actions taken without appropriate clinical correlation, supervision, or consent***

Examples, case scenarios, and clinical discussions presented in this book are intended for educational purposes only.

They do not represent directives, mandates, or universally applicable solutions.

This book emphasizes shared responsibility, interdisciplinary communication, and ethical practice.

However, the ultimate responsibility for patient care rests solely with the treating clinician.

By using this book, the reader acknowledges the necessity of independent clinical judgment, professional accountability, and adherence to local regulations.

Table of Contents

Front Matter

- Otology & Audiology One Mission
- The Philosophy Behind The Books
- Our Commitment
- Why This Book Exists
- How to Read This Book
- Preface
- About the Authors
- Acknowledgments
- Dedications
- Disclaimer



Section 1

Otology & Audiology

Language, Scope & Ethics

Chapter 1: *Why Otology & Audiology Must Speak One Language ……*

Chapter 2: *Understanding Surgical Thinking Before Irreversible Decisions ……*

Chapter 3: *Otology Scope of Practice ……*

Chapter 4: *Audiology Scope of Practice ……*

Chapter 5: *Audio-Vestibular Medicine Scope of Practice ……*

Section 2

Ear Embryology, Anatomy &Physiology

Chapter 6: *Anatomy and Embryology of the Ear ……*

Chapter 7: *Auditory Physiology ……*

Chapter 8: Vestibular Physiology ……

Chapter 9: Pharmacology of Otologic Drugs ……

Section 3

Outer Ear, Tympanic Membrane & Middle Ear Pathologies and Management

Chapter 10: Disorders of the Auricle ……

Chapter 11: Aural Atresia ……

Chapter 12: Tympanic Membrane ……

Chapter 13: Trauma of the Middle Ear & Tympanic Membrane Perforations ……

Chapter 14: Retraction Pockets & Atelectasis ……

Chapter 15: Tympanometry Revisited, Beyond Type A / B / C ……

Chapter 16: Otitis Media with Effusion (Foundations) ……

Chapter 17: Acute Otitis Media, Why Audiology Should Be Careful ……

Chapter 18: *When to Test, When to Wait, Middle Ear Edition ……*

Chapter 19: *Chronic Suppurative Otitis Media (CSOM) ……*

Chapter 20: *Complications of Otitis Media ……*

Chapter 21: *Otitic Barotrauma ……*

Chapter 22: *Otosclerosis ……*

Section 4

Neuro-Otology ،Inner Ear &Implantable Hearing Devices

Chapter 23: *Tumors of the Middle Ear (Glomus Tumors) ……*

Chapter 24: *Diseases of the Inner Ear ……*

Chapter 25: *Meniere Disease ……*

Chapter 26: *The Facial Nerve and Bell's Palsy ……*

Chapter 27: *Hearing Loss ……*

Chapter 28: *Tinnitus ……*

Chapter 29: *Otalgia ……*

Chapter 30: Otorrhea ……

Chapter 31: Vertigo ……

Chapter 32: Ear Operations ……

Chapter 33: Rehabilitation of the Impaired Hearing ……

Chapter 34: Hearing Aid Fitting and Follow Up Protocol ……

Chapter 35: Middle Ear Implantable Hearing Devices ……

Section 4

Pediatric Otology

Chapter 36: Pediatric Otology ……

Chapter 37: What Otology **Expects from Audiology ……**

بسم الله الرحمن الرحيم

SECTION 1

Otology & Audiology

Language, Scope & Ethics

Chapter

1

Why Otology & Audiology Must Speak One Language

The problem was never lack of expertise.

The problem was lack of understanding.

Modern hearing care is filled with experts.

Highly trained otologists.

Highly skilled audiologists.

Advanced technology.

International guidelines.

Yet patients still experience delayed decisions, conflicting opinions, and fragmented care.

Not because clinicians are unqualified—

but because each specialty often thinks in isolation.

Otology and audiology do not fail independently.

They fail at the interface.

Two Ways of Seeing the Same Patient

Otology sees:

- ***Anatomy***
- ***Pathology***
- ***Surgical feasibility***

- *Risk, timing, and irreversibility*

Audiology sees:

- *Function*
- *Variability*
- *Reliability of data*
- *Long-term auditory access and outcomes*

Both perspectives are correct.

Neither is sufficient alone.

A normal cochlea does not guarantee usable hearing.

A measurable response does not guarantee functional benefit.

The patient lives in the space between structure and function.

Where the Gap Becomes Dangerous

The most critical decisions in hearing care occur when:

- *Data is incomplete*
- *Findings are borderline*
- *Outcomes are uncertain*
- *Time matters*

This is where:

- *Audiological reports can be technically accurate yet clinically misleading*
- *Surgical decisions can be anatomically sound yet functionally inappropriate*
- *Silence between specialties becomes interpreted as agreement*

The absence of a shared language does not delay decisions—

it distorts them.

Why Guidelines Are Not Enough

Guidelines define boundaries.

They do not make decisions.

They cannot account for:

- ***Individual variability***
- ***Contextual risk***
- ***Family dynamics***
- ***Real-world functional demands***

When clinicians rely on guidelines without shared interpretation, they replace judgment with comfort.

True clinical excellence begins where guideline interpretation requires clinical judgment.

A Shared Language Is Not Compromise

This book does not ask:

- ***Otologists to abandon surgical rigor***
- ***Audiologists to dilute diagnostic precision***

It asks both to:

- ***Understand how the other reasons***
- ***Recognize the limits of their own data***
- ***Anticipate how their conclusions will be interpreted***

A shared language does not blur roles.

It clarifies responsibility.

The Cost of Not Speaking One Language

When specialties fail to align:

- *Decisions are delayed*
- *Families receive mixed messages*
- *Documentation becomes vulnerable*
- *Accountability becomes unclear*

And when outcomes fall short,

the patient does not ask who decided—

only why it happened.

The Purpose of This Chapter

This chapter is not about agreement.

It is about alignment.

Alignment in:

- *Interpretation*
- *Communication*
- *Decision-making*
- *Documentation*

Because hearing care succeeds only when:

Structure and function are interpreted together,

by clinicians who understand both.

Clinical Summary

Otology and audiology approach the same patient through different lenses. When these lenses are not aligned, technically correct decisions may lead to clinically harmful outcomes.

Key Points

- *Expertise does not prevent fragmentation*

- ***Data without context misleads***
- ***Silence between specialties is never neutral***
- ***Patients live with outcomes, not rationales***

Red Flags

- ***Confident conclusions based on incomplete data***
- ***Decisions made without multidisciplinary discussion***

Pitfalls

- ***Decisions made outside defined scope***
- ***Lack of documented interdisciplinary communication***

Ethics

Ethical care demands transparency, humility, and shared responsibility across specialties.

Documentation

Document not only results, but reasoning and interdisciplinary discussions.

Avoid absolute language.

Describe limitations, context, and uncertainty explicitly.

Final Clinical Message

Shared language is not optional.

It is the foundation of safe hearing care.

Take-Home Messages

- ***Speak together before acting***
- ***Interpret before deciding***

Chapter



2

Understanding Surgical Thinking Before Irreversible Decisions

This chapter is not about how surgery is performed.

It is about how surgery is decided.

For many audiologists, otologic surgery appears as a moment of action.

An intervention.

A solution.

For the otologist, it is something else entirely.

It is a point of no return.

How Otologists Think Differently

Otologists are trained to think in terms of:

- Anatomy that cannot be negotiated
- Pathology that may progress
- Risk that must be justified
- Timing that cannot be reversed

Every surgical decision carries a permanent signature.

Once made, it cannot be recalled, repeated, or corrected without consequence.

This reality shapes how otologists weigh uncertainty.

Why This Matters to Audiologists

Audiological data often arrives before surgical discussion.

It frames expectations.

It influences urgency.

It guides consent.

When audiologists do not understand how otologists interpret:

- *Variability*
- *Borderline findings*
- *Functional uncertainty*

They may unintentionally push decisions toward action—or delay—without meaning to.

Understanding surgical thinking does not mean agreeing with it.

It means anticipating how your data will be used.

The Most Common Misunderstanding

Audiologists often ask:

"Is this patient see-through for surgery?"

Otologists ask something different:

"If this goes wrong, can I justify why I acted when I did?"

This difference is not philosophical.

It is legal, ethical, and permanent.

What This chapter Will Teach You

This chapter will not teach audiologists how to operate.

It will teach them how otologists:

- ***Define acceptable risk***
- ***Interpret uncertainty***
- ***Balance anatomy with prognosis***
- ***Decide when not to operate***

It will also show where otologists rely—often heavily—on audiological interpretation.

Reading This chapter Correctly

Read these chapters not as explanations, but as translations.

When you see:

- ***Hesitation → understand accountability***
- ***Requests for repetition → understand legal defense***
- ***Delayed action → understand irreversible risk***

This volume exists so audiologists can:

- ***Write reports that guide rather than mislead***
- ***Communicate findings in a way surgeons can safely act upon***
- ***Participate meaningfully in high-stakes decisions***

A Responsibility Shared

Otologic decisions are never owned by one person.

They are shaped by:

- ***Data***
- ***Interpretation***
- ***Communication***
- ***Documentation***

This volume is an invitation to step into that responsibility before the incision is ever made.

Because once surgery begins, thinking has already ended.

How Otologists Think When Decisions Cannot Be Undone

Surgery is not a skill.

It is a commitment.

To the outside observer, otologic surgery appears decisive.

An incision.

A plan.

An intervention.

To the otologist, it is something far heavier.

It is a decision that will be questioned—not today, but years later.

By families.

By colleagues.

By courts.

By the patient's outcome.

And once made, it cannot be undone.

The Weight of Irreversibility

Otologists are trained under a constant, silent question:

"Can I justify this decision if the outcome is poor?"

This question shapes everything:

- *Timing*

- ***Caution***
- ***Requests for confirmation***
- ***Apparent hesitation***

Irreversibility changes how uncertainty is tolerated.

Audiological uncertainty may invite repetition.

Surgical uncertainty invites restraint.

Anatomy Is Not Permission

A surgically accessible cochlea does not equal a surgically appropriate decision.

Otologists think in layers:

1. ***Can I technically do this?***
2. ***Should I do this now?***
3. ***What happens if I don't?***
4. ***What happens if I do—and it fails?***

Audiological data feeds directly into layers 2, 3, and 4.

This is why wording matters more than numbers.

How Risk Is Really Calculated

Risk is not calculated mathematically.

It is calculated defensively and ethically.

Otologists consider:

- ***Permanence of harm***
- ***Likelihood of benefit***
- ***Ability to explain the decision later***
- ***Documentation strength***

A decision that cannot be defended should not be performed.

Why Otologists Ask for "More Tests"

Requests for repetition are often misinterpreted as doubt or delay.

In reality, they are often:

- ***Attempts to narrow uncertainty***
- ***Efforts to strengthen justification***
- ***Protection against irreversible harm***

When audiologists understand this,

they stop hearing mistrust

and start hearing responsibility.

The Silence That Means "No"

Otologists rarely say "no" loudly.

They say it quietly.

- ***By delaying***
- ***By requesting more discussion***
- ***By reframing urgency***
- ***By asking different questions***

Silence is not indecision.

It is often a decision to protect.

Where Audiology Shapes Surgical Thinking

Audiological interpretation influences:

- ***Surgical timing***
- ***Family counseling***
- ***Consent framing***

- *Outcome expectations*

A single sentence in a report can:

- *Accelerate surgery*
- *Delay it responsibly*
- *Or expose the surgeon legally*

Audiology does not just inform surgery.

It steers it.

The Responsibility Before the Incision

The incision is the final act.

The decision is made long before.

Otologists do not fear operating.

They fear operating without justification.

Understanding this mindset allows audiologists to:

- *Communicate findings safely*
- *Frame uncertainty responsibly*
- *Participate fully in irreversible decisions*

Clinical Summary

Otologic decision-making is defined by irreversibility.

Every surgical choice must be defensible long after the moment of action.

Key Points

- *Irreversible decisions demand restraint*
- *Anatomy does not equal appropriateness*
- *Silence often reflects responsibility, not hesitation*
- *Audiological wording shapes surgical risk*

The Golden Rule

Never frame data in a way that pressures an irreversible decision you cannot defend later.

Common Mistakes

- ***Interpreting caution as lack of expertise***

Red Flags

- ***Reports implying certainty where variability exists***
- ***Pressure to "just proceed"***

Pitfalls

- ***Underestimating medico-legal weight of reports***
- ***Ignoring long-term outcome accountability***
- ***Absolute language in reports***
- ***Missing documentation of interdisciplinary discussion***
- ***Acting outside defined privileges***

Clinical Decision-Making

Pause irreversible decisions until interpretation is shared and documented.

The Uncomfortable Truth

Most surgical regret begins with a sentence that sounded confident.

The Sentence That Changes Everything

"Findings suggest potential benefit, but functional reliability and long-term outcome remain uncertain."

Clinical Case

A surgically successful procedure with disappointing functional outcome due to premature decision-making.

Ethics

The safest decision was not the fastest—but the most defensible.

Report Writing

Avoid language that implies inevitability or guarantees.

Final Clinical Message

Understanding surgical thinking is not optional for audiologists—it is protective.

3

Otology Scope of Practice

Where Authority Ends and Responsibility Begins

How Surgical Decisions Are Actually Made

"Audiology does not need to know how to operate.

It needs to know how surgeons decide."

This book exists because many of the most critical audiological decisions are shaped by how otologists think under irreversible risk.

If audiology misunderstands that thinking,

its data—no matter how accurate—can become dangerous.

This book teaches audiologists to:

- ***Anticipate surgical thinking***
- ***Frame data for irreversible decisions***
- ***Communicate uncertainty safely***
- ***Protect patients before the incision***
- ***Protect colleagues after outcomes***

It is written for audiologists who:

- ***Participate in CI pathways***
- ***Write reports that trigger action***

- *Sit in MDT rooms*
- *Influence consent—often silently*

Scope of practice is not about power.

It is about protection.

In modern hearing care, the most dangerous errors rarely come from lack of skill.

They come from blurred boundaries.

Boundaries between:

- ***What a clinician can do***
- ***What a clinician should do***
- ***And what a clinician is accountable for***

This chapter exists to define those boundaries clearly—

not to restrict practice,

but to protect patients, clinicians, and outcomes.

Scope of Practice Is Not Skill

Otologists are highly trained.

Their technical abilities often exceed what is required in daily practice.

But skill does not define scope.

Scope of practice is defined by:

- ***Training and credentialing***
- ***Institutional privileges***
- ***Legal and regulatory frameworks***
- ***Ethical responsibility***

Being able to perform an act does not grant authority to decide it in isolation.

This distinction is critical in otology, where decisions are often irreversible.

What Otology Is Fully Accountable For

Otology carries direct responsibility for:

- ***Confirming that a surgical indication is valid***
- ***Assessing anatomical feasibility and risk***
- ***Ensuring informed, documented consent***
- ***Timing of intervention***
- ***Ownership of surgical outcomes***

Once an otologic decision is executed,
responsibility does not disappear—it concentrates.

What Otology Is Not Responsible For

Otology should not be solely responsible for:

- ***Interpreting audiological data outside its methodological limits***
- ***Making decisions based on incomplete or unreliable functional assessment***
- ***Replacing audiological judgment with surgical opinion***
- ***Acting on reports that lack transparency, context, or limitations***

When this line is crossed, responsibility becomes unclear—
and patients pay the price.

The Gray Zones — Where Shared Responsibility Is Mandatory

Some clinical areas do not belong to one specialty.

These include:

- ***Cochlear implant candidacy***
- ***Borderline or asymmetric hearing loss***
- ***Pediatric and pre-lingual cases***
- ***Auditory neuropathy spectrum disorder***
- ***Complex medical or developmental profiles***

In these zones,

no single specialty owns the decision.

Decisions must be:

- ***Shared***
- ***Documented***
- ***Defensible from multiple perspectives***

Silence or unilateral action in gray zones is a scope violation.

When Otology Must Say "Stop"

Stopping is not failure.

It is professional maturity.

Otology must pause or refuse to proceed when:

- ***Audiological data is unreliable or poorly contextualized***
- ***Counseling has not addressed uncertainty or alternatives***
- ***Documentation does not support defensible consent***
- ***External pressure replaces clinical judgment***

Proceeding under these conditions exposes:

- ***The patient to harm***
- ***The clinician to legal risk***

- *The system to ethical failure*

Scope, Privileges, and Institutional Reality

Scope of practice is enforced not only by ethics,

but by privileges and policy.

Otologic authority exists only within:

- ***Granted privileges***
- ***Defined service scope***
- ***Institutional protocols***
- ***Multidisciplinary governance***

Acting outside these boundaries—even with good intentions—

is indefensible.

The Relationship with Audiology

Otology depends on audiology more than it often admits.

Otology should expect:

- ***Clear reporting of reliability and limitations***
- ***Functional interpretation—not numbers alone***
- ***Explicit statements of uncertainty***

Otology must not:

- ***Assume audiological conclusions are absolute***
- ***Pressure audiologists into certainty***
- ***Use reports as shields for predetermined decisions***

Respecting scope strengthens collaboration.

The Real Cost of Scope Violations

Scope violations rarely announce themselves.

They appear later as:

- *Poor functional outcomes*
- *Family dissatisfaction*
- *Conflicting documentation*
- *Legal scrutiny*

By the time they surface,
it is too late to redefine boundaries.

Clinical Summary

Otology scope of practice defines responsibility, not authority.
Clear boundaries protect patients, clinicians, and outcomes—
especially in irreversible decisions.

Key Points

- *Skill does not equal scope*
- *Gray zones require shared ownership*
- *Silence is not consent*
- *Scope violations often appear late—but cost early*

The Golden Rule

Never act beyond your scope.

Common Mistakes

- *Acting on incomplete reports*
- *Equating privileges with absolute authority*
- *Proceeding under pressure*

Red Flags

- *Unilateral decisions in gray zones*
- *Documentation lacking interdisciplinary input*
- *Urgency without shared justification*

Pitfalls

- *Inadequate consent framing*
- *Acting outside granted privileges*

Clinical Decision-Making

Pause when boundaries are unclear.

Clarify scope before committing to irreversible action.

The Uncomfortable Truth

Most scope violations begin with good intentions.

The Sentence That Changes Everything

"This decision falls within a shared scope and requires multidisciplinary agreement."

What I Would Explain to the Family

"This decision involves more than one specialty.
We will not proceed until everyone agrees it is right."

Clinical Case

A technically successful surgery later questioned due to missing shared documentation.

Thinking Behind the Scene

Stopping protected the patient—and the team.

Ethics

Ethical practice respects boundaries as much as expertise.

Documentation

Document scope, shared decisions, and limits explicitly.

Report Writing

Avoid language that transfers responsibility silently.

If scope is unclear → Do not proceed.

Dangerous Sentence Alert

“No contraindication from our side.”

without shared review and documented agreement

Evidence-Based & International Guidelines

Scope and governance are integral to safe clinical practice.

Teaching Pearl

Boundaries prevent more harm than boldness ever will.

Final Clinical Message

Clear scope protects everyone—especially the patient.

Take-Home Messages

- ***Define boundaries early***
- ***Share responsibility explicitly***
- ***Document defensively and ethically***

Chapter



4

Audiology Scope of Practice

Where Interpretation Becomes Responsibility

Audiology does not end at measurement.

It begins at interpretation.

In hearing care, few specialties influence decisions as quietly—and as powerfully—as audiology.

Audiologists rarely make incisions.

They rarely sign surgical consent.

Yet their words often initiate irreversible pathways.

This chapter exists to clarify where audiological responsibility begins,
where it must stop,
and how crossing that boundary—intentionally or not—can harm patients and colleagues alike.

Scope of Practice Is Not Testing

Audiology is often misunderstood as a technical service.

A collection of tests.

A set of measurements.

This view is dangerously incomplete.

Audiology scope of practice includes:

- ***Selection of appropriate tests***
- ***Assessment of test reliability***
- ***Interpretation of findings within context***
- ***Communication of limitations and uncertainty***
- ***Anticipation of how reports will be used***

Producing data is not neutral. Interpretation always carries responsibility.

What Audiology Is Fully Accountable For

Audiology carries direct responsibility for:

- ***Test validity and reliability***
- ***Methodological transparency***
- ***Functional interpretation of results***
- ***Explicit documentation of uncertainty***
- ***Framing conclusions that guide—not pressure—decisions***

Once a report is issued,

it becomes a clinical instrument—not a passive record.

What Audiology Is Not Responsible For

Audiology is not responsible for:

- ***Making surgical decisions***
- ***Determining anatomical feasibility***
- ***Predicting guaranteed outcomes***
- ***Replacing multidisciplinary discussion with report conclusions***

When audiological language implies inevitability,

scope is silently violated.

The Most Dangerous Assumption in Audiology

That numbers speak for themselves.

They do not.

Audiologists must constantly ask:

- ***How will this sentence be interpreted?***
- ***Who will act on this report?***
- ***What decision might this wording accelerate?***

Silence about limitations is not objectivity.

It is exposure.

Shared Zones — Where Audiology Must Not Act Alone

Audiology does not own decisions in:

- ***Cochlear implant candidacy***
- ***Pediatric uncertainty***
- ***Auditory neuropathy***
- ***Medically or developmentally complex cases***

In these areas,

audiology must:

- ***Inform***
- ***Contextualize***
- ***Clarify***

But never conclude alone.

When Audiology Must Say "I Don't Know"

Uncertainty is not weakness.

It is integrity.

Audiology must pause when:

- ***Reliability is questionable***
- ***Results conflict across tests***
- ***Functional performance does not match thresholds***
- ***Time pressure threatens interpretation***

Clear uncertainty protects:

- ***The patient***
- ***The surgeon***
- ***The audiologist***

Reports as Decision Triggers

Audiological reports are often treated as:

- ***Permission slips***
- ***Gatekeepers***
- ***Legal shields***

They are none of these by default.

Every report should answer:

- ***What is known***
- ***What is not known***
- ***What must not be assumed***

Failure to do so shifts responsibility unfairly—and dangerously.

Relationship with Otology

Audiology should expect:

- ***Surgical judgment, not blind acceptance***
- ***Questions, not compliance***

- *Requests for clarification*

Audiology must not:

- *Write to please*
- *Write to hurry decisions*
- *Write to avoid follow-up discussion*

Respecting scope strengthens credibility.

Clinical Summary

Audiology scope of practice centers on interpretation, context, and communication—not decision ownership.

Clear boundaries protect patients and preserve multidisciplinary integrity.

Key Points

- *Testing is not the endpoint*
- *Reports shape irreversible decisions*
- *Silence about uncertainty is harmful*
- *Audiology influences more than it decides*

The Golden Rule

Never write a conclusion that pressures a decision you are not responsible for making.

Common Mistakes

- *Overconfident language*
- *Conclusions without limitations*
- *Equating clarity with certainty*

Pitfalls

- *Assuming the reader understands limitations*

- ***Writing for speed rather than safety***
- ***Missing documentation***
- ***Acting outside defined scope***

What I Would Explain to the Family

"These tests help us understand hearing—but they don't decide alone.

The next step requires shared discussion."

Chapter

5

Audio-Vestibular Medicine Scope of Practice

Where Medical Authority, Diagnostic Integration, and Clinical Accountability Converge

Audio-Vestibular Medicine is a medical specialty.

In modern hearing and balance care, few areas generate more confusion—and more silent conflict—than the role of the Audio-Vestibular Medicine physician.

In many clinical settings, the distinction between:

- ***Audiologist***
- ***Audio-Vestibular Physician***

is blurred, undocumented, or entirely misunderstood.

This chapter exists to correct that confusion—
not to elevate one role over another,
but to define boundaries, protect patients, and assign responsibility clearly.

Why This Confusion Exists

The confusion surrounding Audio-Vestibular Medicine is not accidental.

It arises from:

- *Variability in training models across countries*
- *Overlapping clinical environments*
- *Inconsistent institutional definitions*
- *Absence of written scope frameworks*

When roles are undefined,
authority becomes negotiable,
and responsibility becomes fragmented.
Where scope is unclear, conflict replaces structure.

Who Is the Audio-Vestibular Medicine Physician?

An Audio-Vestibular Medicine physician is a medically trained doctor whose core responsibility is the diagnosis and medical management of hearing and balance disorders.

This role is defined not by test performance,
but by diagnostic ownership.

The Audio-Vestibular physician is trained to:

- *Integrate audiological and vestibular findings*
- *Construct differential diagnoses*
- *Identify medical, neurological, and systemic causes*
- *Decide when medical treatment, monitoring, referral, or escalation is required*
- *Assume medico-legal responsibility for diagnostic conclusions*

The defining feature of Audio-Vestibular Medicine is not testing—it is medical judgment.

What Audio-Vestibular Medicine Fully Owns

Audio-Vestibular Medicine carries direct authority and accountability for:

- ***Medical diagnosis of auditory and vestibular disorders***
- ***Etiological interpretation of audiological and vestibular data***
- ***Ordering and interpreting relevant investigations***
- ***Medical decision-making and treatment planning***
- ***Determining urgency and referral pathways***
- ***Issuing final diagnostic impressions***
- ***Medico-legal responsibility for diagnostic conclusions***

Once a diagnosis is declared,

the responsibility is medical, legal, and ethical.

What Audio-Vestibular Medicine Does NOT Own

Audio-Vestibular Medicine does not replace other specialties.

It does not own:

- ***Surgical decision-making***
- ***Acting outside multidisciplinary collaboration***
- ***Overriding ENT or Neurology scope***

Medical authority does not justify professional isolation.

Relationship with Audiology

The relationship between Audio-Vestibular Medicine and Audiology

is complementary, not hierarchical.

Neither role substitutes the other.

Each protects the other when boundaries are respected.

- ***Audio-Vestibular Physicians are responsible for:***
  - ***Medical integration***

  - *Diagnostic labeling*
  - *Clinical risk assessment*
- *Integrates findings medically*
- *Establishes diagnosis or differential*

Relationship with Otology / ENT

Audio-Vestibular Medicine is not surgical.

Its role in relation to Otology includes:

- ***Pre-surgical diagnostic clarification***
- ***Functional interpretation of auditory and vestibular status***
- ***Identification of medical contraindications***
- ***Post-operative diagnostic follow-up***

Audio-Vestibular physicians must:

- ***Avoid implying surgical authority***

Diagnosis informs surgery—
but does not authorize it.

Where Conflict Most Commonly Occurs

Conflict often arises in:

- ***CI candidacy discussions***
- ***Borderline or fluctuating hearing loss***
- ***ANSD***
- ***Vestibular syndromes***
- ***Report wording and diagnostic labels***

In most cases, the conflict is not clinical—it is linguistic.

Poorly defined language leads to misplaced authority.

Silence does not protect.

Documentation does.

The Patient's Perspective

Patients do not understand professional boundaries.

They understand outcomes.

They need:

- ***One medically accountable voice***
- ***Clear explanations***

Clinical Summary

Audio-Vestibular Medicine is a medical specialty defined by diagnostic ownership, etiological integration, and medico-legal accountability—not only by test performance.

Key Points

- ***Scope defines accountability, not status***
- ***Integration is the physician's core role***
- ***Clear boundaries protect collaboration***

The Golden Rule

Never issue a diagnostic conclusion unless you are prepared to own its medical and legal consequences.

Common Mistakes

- ***Acting as an advanced tester rather than a physician***
- ***Issuing diagnoses without full integration***
- ***Blurring boundaries with audiology or ENT***

Red Flags

- ***Diagnostic labels without etiological reasoning***
- ***Reports lacking medical context***
- ***Acting beyond institutional privileges***

Pitfalls

- ***Assuming shared responsibility without documentation***
- ***Using ambiguous language to avoid accountability***
- ***Acting outside defined medical scope***

The Uncomfortable Truth

Most conflicts arise when medical authority is implied rather than defined.

The Sentence That Changes Everything

"Based on integrated audiological and clinical findings, the medical diagnosis is…"

"This decision was reached through multidisciplinary discussion with shared accountability."

What I Would Explain to the Family

"These tests help us understand the problem.
My role is to put them together and explain what they mean medically."

Take-Home Messages

- ***Protect collaboration through clarity***

SECTION 2

Ear Embryology, Anatomy &Physiology

Chapter

6

Anatomy and Embryology of the Ear

Structure Before Function

Why This Chapter Matters

This chapter is not an anatomy review for surgeons.

It is an anatomy orientation for audiologists.

Otologists do not look

at the ear as:

- *Thresholds*
- *Curves*
- *Waveforms*

They look at it as:

- *Spaces*
- *Boundaries*
- *Blood supply*
- *Nerves*
- *Structures that cannot be negotiated*

Understanding ear anatomy and embryology helps audiologists appreciate:

- *Why some ears are operable*
- *Why others are not*

1. Embryology of the Ear — Why Development Matters

The ear develops from three embryological components, each with different vulnerabilities:

- ***External ear** → ectoderm*
- ***Middle ear** → endoderm*
- ***Inner ear** → otic placode (ectoderm)*

Embryology explains why "normal-looking" ears may function abnormally.



▲ *Figure 6–1. Development of external ear*

Structures Derived from Various Hillocks

1. Tragus
2. Crus of the helix
3. Helix
4. Antihelix
5. Scapha and the antitragus
6. Ear lobule


Invaginating neural tube
Otic placode
22 days
Endoderm
Pharynx
Root of rhombencephalon
Endolymphatic duct
Utricular portion of otic vesicle
Saccular portion
Mesenchymal condensation
1st pharyngeal cleft
Tubotympanic recess
Auditory ossicles embedded in loose mesenchyme
Wall of inner ear
Auditory tube
External auditory meatus
Meatal plug
Primitive tympanic cavity

▲ *Figure 6–2. Development of external ear*
Development of middle and inner ear


Cartilage
Stapes
Semicircular canals
Incus
Malleus
VIII N
Cochlea
Middle ear
Tympanic membrane
Eustachian tube
Bony part of ear canal
Lobule
Cartilaginous part of ear canal

▲ *Figure 6–3. Parts of ear*

2. External Ear Anatomy — More Than a Sound Funnel

The external ear includes:

- ***Auricle***
- ***External auditory canal***

Key surgical relevance:

- ***Canal stenosis or atresia affects access***
- ***External ear anomalies often coexist with middle ear defects***
- ***External anatomy influences device placement and outcomes***

For audiology:

- ***Conductive components must be interpreted anatomically***
- ***External ear findings should never be ignored as "minor"***


Helix
Posterior crus of anthelix
Triangular fossa
Anterior crus of anthelix
Cymba concha
Scapha
Antihelix
Tragus
Cavum concha
External auditory meatus
Antitragus
Lobule

▲ *Figure 6–4. Anatomy of the pinna.*
The pinna consists of a cartilaginous framework covered by skin.


1
10
2
5
3
9
4
6
7
8

▲ *Figure 6–5. Development of external ear*
Parts of pinna. 1. helix, 2. triangular fossa, 3. antihelix,
4. tragus, 5. cymba concha, 6. cavum concha, 7. antitragus,
8. lobule, 9. supratragal notch, 10. scaphoid fossa


Temporal branch of CN VII
Auricular branches of CN X (and CN VII and IX via Arnold's nerve)
Greater auricular nerve
Auriculo-temporal nerve
Mastoid branch of lesser occipital nerve
Posterior auricular branch of CN VII
Lesser auricular nerve
Greater auricular nerve
A
Mastoid branch of lesser occipital nerve
Lasser auricular nerve
Posterior auricular branch of CN VII
Greater auricular nerve
B

▲ *Figure 6–6. Nerve supply of pinna*


Superficial temporal artery
Posterior auricular artery
External carotid artery
Lateral
Posterior

▲ *Figure 6–7. Blood supply of pinna*



▲ *Figure 6–8. Coronal section of the ear canal.*
The skin of the cartilaginous and osseous canals is magnified.


Sebaceous gland
Capillary network
Epidermis
Ductal opening
(hair removed)
Duct of sebaceous
gland
Hair
Ductal
opening
Duct of modified
apocrine gland
Nerve
Modified apocrine gland

▲ *Figure 6–9*. ***Skin of the cartilaginous portion of the external auditory canal***


Superior
Anterior
Middle cranial fossa
Temporomandibular joint
superficial temporal
artery and vein
Auriculotemporal nerve
Parotid gland
preauricular lymph node
Medial
Middle ear
Outside world
Lateral
Mastoid
Posterior
Jugular bulb
Carotid
Facial nerve
Styloid process
Parotid gland
Digastric muscle
Inferior

▲ *Figure 6–10.* ***Relationships of the right external auditory canal.***


Sound waves
Ossicles
Cochlea
Auditory nerve
Tympanic membrane
Eustachian tube
External ear
Middle
Inner

▲ *Figure 6–11. Anatomy of the ear. The external ear collects sound pressure waves and funnels them toward the tympanic membrane.*
The middle ear ossicles transmit the sound waves to the inner ear (cochlea).
The middle ear acts to match the impedance difference between the air of the external environment to the fluid within the cochlea.
This permits maximal sound transmission.

3. Middle Ear Anatomy — The Mechanical Gatekeeper

The middle ear consists of:

- ***Tympanic membrane***
- ***Ossicular chain***
- ***Middle ear cavity***
- ***Eustachian tube***

Otologists focus on:

- *Mobility*
- *Continuity*
- *Ventilation*

Audiological relevance:

- *Air-bone gaps are **mechanical stories***
- *Stability matters more than magnitude*
- *Chronic pathology alters surgical risk*

A conductive loss is never "just conductive" to a surgeon.


M
I
Epitympanum
Aditus
Mastoid
ATI
ET
Protympanum
Chorda-
tympani
S
Retro-
tympanum
Mesotympanum
P
Po
ST
Su
F
Hypotympanum
Jugular bulb

▲ Figure 6–12. Parts of the middle ear cavity and the ventilation pathways of middle ear cleft (ATI: Anterior tympanic isthmus; I: Incus, M: Malleus; S: Stapes; ET: Eustachian tube; C: Chorda tympani; ST: Sinus tympani; P: Promontory; F: Funiculus; Su: Subicu lum; Po: Ponticulus; J: Jugular bulb)


Mastoid antrum
Stapes
Tympanal orifice
Bone
Isthmus (eustachian tube)
Tubal cartilage
Pharyngeal orifice
Mastoid air cells
Promontory
Round window
Hypotympanum
Mixed gland
Levator veli palatini muscle

▲ *Figure 6–13. Diagrammatic representation of middle ear cleft*


Notch of rivinus
Pars flaccida (Shrapnell's membrane)
Posterior malleal fold
Anterior malleal fold
Handle of malleus (manubrium)
Umbo
Tympanic ring
Annulus

▲ *Figure 6–14. Anatomy of the tympanic membrane (a left ear).*



▲ *Figure 6–15.Endoscopic photograph of the right tympanic membrane.*


Head of malleus
Short process of malleus
Neck of malleus
Lateral process
Anterior process
Manubrium of malleus (The long process)
Umbo
Lenticular process
Incudostapedial joint
Incudomallear joint
Short process of incus
Body of incus
Long process of incus
Stapes capitulum
Posterior crus
Anterior crus
Footplate

▲ *Figure 6–16. The middle ear ossicles.*


MCF
Superior
Aditus
Antrum
Posterior
Eustachian tube
Anterior
Carotid artery
Mastoid tip
Jugular bulb
Inferior
Processus cochleariformis
LSC
OW
Facial nerve
ET
Aditus
RW
Medial wall
Promontary
Anterior wall
Lateral wall
Posterior wall
Stylomastoid foramen

▲ *Figure 6–17. Relation of middle ear with surrounding important structures*


Malleus
Incus
Head
Neck
Anterior process
Lateral process
Handle
Body
Short process
Long process
Lenticular process
Anterior crus
Head
Neck
Footplate
Posterior crus
Stapes

▲ *Figure 6–18. Ossicular chain*

4. Inner Ear Anatomy — Where Irreversibility Begins

The inner ear includes:

- *Cochlea*
- *Vestibular system*
- *Endolymphatic structures*

Critical surgical considerations:

- *Cochlear turns and patency*
- *Modiolus and neural proximity*
- *Vestibular-cochlear relationship*

Audiological relevance:

- *Inner ear anomalies affect candidacy*
- *Thresholds do not reveal neural survival*
- *Anatomy defines limits of benefit*


Stapes
Incus
Oval window
Auditory nerve
Malleus
Basilar membrane
Perilymph
Tympanic membrane
Endolymph
Sound waves
Ear canal
Tectorial membrane
Round window
Eustachian tube
Middle ear
Inner ear

▲ *Figure 6–19. Structure of cochlea*


Scala vestibuli
Reissner's membrane
Inner hair cell
Scala media
Tectorial membrane
Stria vascularis
Outer hair cells
Deiter's cells
Basilar membrane
Cochlear nerve
Tunnel of Corti
Scala tympani

▲ *Figure 6–20. Structure of organ of Corti*


Superior semicircular canal
Incus
Malleus
Crus commune
Posterior semicircular canal
Geniculate ganglion
Chorda tympani nerve
Lateral semicircular canal
Stapes
Endolymphatic duct
Ponticulus
Sinus tympani
Stapedius muscle
Subiculum
Facial nerve
Round window niche

▲ *Figure 6–21.* **Relationship of the middle ear structures with the inner ear (a right ear).**


Superior
Anterior
Middle cranial fossa
Temporal lobe
Tensor tympani muscle
Eustachian tube
Medial
Lateral
Inner ear
Aditus to mastoid
External auditory meatus
Mastoid
Jugular bulb
Carotid artery
Posterior
Inferior

▲ *Figure 6–22. The relationships of the right middle ear.*


Superior semicircular canal
Depression for utricle
Vestibular crest
Depression for saccule
Scala vestibuli
Scala tympani
Osseous spiral lamina
Lateral semicircular canal
Cochlear recess
Cochlear canaliculus
Posterior semicircular canal
Fenestra cochleae
Opening of crus commune
Opening of aqueduct of vestibule

▲ *Figure 6–23. Diagrammatic representation of bony labyrinth*


Superior semicircular duct
Posterior semicircular duct
Utricle
Lateral semicircular duct
Ductus reuniens
Endolymphatic sac
Saccule
Cochlear duct

▲ *Figure 6–24. Membranous labyrinth shown floating inside the bony*


Basilar artery
Anterior inferior cerebellar artery
Vestibular artery
Common cochlear artery
Superior SSC
Anterior vestibular artery
Horizontal/ lateral SSC
Posterior vestibular artery
Posterior SSC
Cochlea
Vestibule

▲ *Figure 6–25. Blood supply of membranous labyrinth*


Cochlea
Foramen spinosum
Foramen ovale
Internal carotid
Cochlear nerve
Facial nerve
Greater petrosal nerve
Geniculate ganglion
Endolymphatic duct
Endolymphatic sac
Lateral semicircular canal
Superior semicircular canal
Posterior semicircular canal
Inferior vestibular nerve
Jugular foramen
Superior vestibular nerve

▲ *Figure 6–26. The location of the inner ears within the skull base.*


Vestibular membrane
Scala vestibuli
Tectorial membrane
Cochlear duct
Organ of Corti
Scala tympani
Zona pectinata
Cochlear nerve
A
Zona arcuata
Basilar membrane
Tectorial membrane
Reticular lamina
Outer hair cells
Organ of Corti
Inner hair cell
Supporting cells
B
Cochlear nerve
Inner tunnel
Nerve fibers
Pillar cell (rod of Corti)
Basilar membrane

▲ *Figure 6–27. Structure of organ of Corti*


Anterior vestibular artery
Common cochlear artery
Endolymphatic duct
Spiral modiolar artery
Vestibulo-cochlear artery
Posterior vestibular artery
Anterior inferior cerebellar artery
Main cochlear artery
Labyrinthine artery
Basilar artery

▲ *Figure 6–28. Blood supply of inner ear*

5. Facial Nerve and Vascular Relationships

Otologists are constantly aware of:

- ***Facial nerve course***
- ***Vascular variants***
- ***Risk of injury***

Audiological importance:

- ***Facial nerve anomalies may coexist with hearing loss***
- ***Syndromic patterns must be recognized early***
- ***Some "routine" surgeries carry non-routine risk***

6. Embryology Meets Clinical Reality

When audiologists understand embryology, they can:

- ***Anticipate associated anomalies***
- ***Recommend appropriate imaging***
- ***Frame reports with anatomical awareness***

This strengthens MDT discussions and protects patients.

Clinical Summary

Accurate interpretation of hearing and balance tests depends on knowing how each part of the ear is built and what it is designed to do.

The ear is not a single organ.

It is a highly organized system responsible for hearing and balance.

The ear performs two main functions:

- ***Hearing***
- ***Balance***

Anatomically, it is divided into three main parts:

1. *External Ear*
2. *Middle Ear*
3. *Inner Ear*

Each part has a distinct structure and a specific function.

I. External Ear

Components

The external ear consists of:

- *Auricle (Pinna)*
- *External Auditory Canal*
- *Tympanic Membrane*

A. Auricle (Pinna)

Structure

The auricle is a cartilaginous structure covered by skin.

It is attached to the skull by ligaments and small muscles.

Main Anatomical Features

- *Helix*
- *Antihelix*
- *Concha*
- *Tragus*
- *Lobule*

The lobule contains no cartilage.

Function

- *Collects sound waves*
- *Helps direct sound into the ear canal*

- *Assists in sound localization*

B. External Auditory Canal

Anatomy

- *Extends from the concha to the tympanic membrane*
- *Approximately 24 mm in length*
- *Not a straight tube*

It has two parts:

1. *Cartilaginous part, outer one third*
2. *Bony part, inner two thirds*

Important Features

- *The cartilaginous part contains hair follicles, sebaceous glands, and ceruminous glands*
- *The bony part has thin skin and no glands*

Functional Importance

- *Conducts sound toward the tympanic membrane*
- *Acts as a resonator that enhances speech frequencies*

C. Tympanic Membrane

Description

The tympanic membrane separates the external ear from the middle ear. It is oval in shape and positioned obliquely.

Layers

- *Outer epithelial layer*
- *Middle fibrous layer*
- *Inner mucosal layer*

Parts

- ***Pars tensa***
- ***Pars flaccida***

Key Landmark

- ***Umbo, the point of maximum concavity, attached to the handle of the malleus***

II. Middle Ear

Middle Ear Cleft

The middle ear cleft includes:

- ***Tympanic cavity***
- ***Eustachian tube***
- ***Mastoid air cells***

Tympanic Cavity

Description

- ***An air filled space within the temporal bone***
- ***Contains the ossicular chain***
- ***Lined by mucous membrane***

Contents

- ***Ossicles, malleus, incus, stapes***
- ***Muscles***
- ***Nerves***

Function

- ***Transmits sound vibrations from the tympanic membrane to the inner ear***

- *Amplifies sound through mechanical leverage*

III. Inner Ear

Location

The inner ear lies deep within the petrous part of the temporal bone.

Structure

It consists of:

- ***Bony labyrinth***
- ***Membranous labyrinth***

Main Components

- ***Cochlea, responsible for hearing***
- ***Vestibule, involved in balance***
- ***Semicircular canals, responsible for rotational balance***

Function

- ***Converts mechanical sound vibrations into nerve impulses***
- ***Detects head movement and position***

Key Points

- ***The ear is divided into external, middle, and inner parts***
- ***Each part has a specific role in hearing and balance***
- ***Structural damage in any part leads to predictable functional changes***

Golden Rule

Understanding ear anatomy is essential for correct audiologic interpretation.

Chapter

7

Auditory Physiology



From Sound to Neural Signal

Why This Chapter Matters

This chapter is not about test batteries.

It is about how hearing actually works—clinically.

Otologists rarely question:

- ***How to run a test***

They question:

- ***What does this result really represent physiologically?***
- ***Which part of the auditory system is functioning—and which is not?***
- ***Can this physiology support intervention?***

Audiology becomes clinically powerful

when physiology—not numbers—guides interpretation.

1. The Auditory System as a Functional Chain

Auditory physiology functions as a continuous chain:

1. ***Sound transmission***
2. ***Mechanical conversion***
3. ***Sensory transduction***

4. *Neural synchrony*
5. *Central processing*

Failure at any level produces different clinical meanings,

even if audiograms look similar.

Similar thresholds do not imply similar physiology.

2. Peripheral Auditory Physiology — More Than Thresholds

Outer Ear & Middle Ear

- *Sound conduction and impedance matching*
- *Mechanical efficiency affects stimulus quality*

Audiological relevance:

- *Poor conduction alters input, not neural integrity*
- *Stable mechanics predict better surgical predictability*

Cochlear Physiology

Key components:

- *Outer hair cells (amplification)*
- *Inner hair cells (transduction)*
- *Endocochlear potential*

Clinical relevance:

- *OHC loss ≠ IHC loss*
- *Presence of OAE does not guarantee usable hearing*
- *Absence of OAE does not define neural status*

Cochlear physiology determines quality, not just audibility.


Stapes footplate (mobile)
Reissner's membrane
Perilymph scala vestibuli
Tectorial membrane
Cochlear duct—scala media
Helicotrema
Cilia
Hair cells
Perilymph scala tympani
Sound waves
Round window membrane (mobile)
Organ of Corti
Basilar membrane

▲ *Figure 7–1. Mechanism of hearing showing tympanocochlear relationship*

3. Neural Physiology — Where Interpretation Becomes Dangerous

Auditory nerve function depends on:

- ***Neural survival***
- ***Synchrony***
- ***Temporal precision***

Clinical implications:

- ***ABR reflects synchrony, not hearing sensitivity alone***
- ***Absent or distorted waves may reflect neural timing failure***
- ***Neural disorders can masquerade as severe cochlear loss***

This is where audiology directly influences surgical thinking.

4. Central Auditory Pathways — Beyond the Cochlea

Central auditory structures:

- *Brainstem*
- *Midbrain*
- *Auditory cortex*

Clinical relevance:

- *Central integrity affects outcomes*
- *Peripheral correction does not fix central processing*
- *Developmental timing matters*

Otologists worry about ***downstream limitations*** *even when surgery is perfect.*

5. Clinical Assessment as Physiological Mapping

Clinical audiology is not testing.

It is ***physiological localization****.*

Each test answers a different question:

- *Pure-tone audiometry → detection*
- *Speech testing → functional integration*
- *OAE → cochlear amplification*
- *ABR → neural synchrony*
- *Immittance → mechanical integrity*

No single test defines the system.

Clinical assessment succeeds when tests are interpreted as physiology, not results.

6. Variability, Reliability, and Clinical Confidence

Physiology is dynamic.

So are patients.

Audiologists must always ask:

- ***Is this result repeatable?***
- ***Is it state-dependent?***
- ***Does it fit the rest of the physiological picture?***

Otologists interpret confidence not by certainty—but by consistency.

7. Why Physiology Matters to Surgical Outcomes

Surgical success depends on:

- ***Neural integrity***
- ***Functional pathways***
- ***Developmental timing***

Audiological physiology informs:

- ***Candidacy***
- ***Timing***
- ***Expectation setting***
- ***Risk-benefit analysis***

Good surgery cannot overcome poor physiology.


Primary auditory cortex (AI)
Non-primary auditory cortex
Medial geniculate nucleus
Cerebral cortex
Fifth order neuron
Dorsal division
Laminated, ventral division
Magnocellular division
Nonlaminated, ventral division
Thalamus
Fourth order neuron
Brachium of inferior colliculus
Commissure of inferior collicu
Pericentral region
Central region
Midbrain
Third order neuron
Cochlear nuclei
Dorsal nuclei
Second order neuron
Lateral lemniscus
Medulla
Trapezoid body
First order neuron
Superior olivary nuclei
Lateral
Medial

▲ *Figure 7–2. The auditory pathway*


Auditory cortex
Medial geniculate body
Inferior colliculus
Dorsal cochlear nucleus
Ventral cochlear nucleus
Cochlear nerve
Superior olivary complex
Cochlea

▲ *Figure 7–3.* ***Central auditory pathway***


Broca's area (Brodmann area 44)
Superior temporal gyrus
Wernicke's area (Brodmann area 22)

▲ *Figure 7–4.* ***Auditory cortex–Brodmann Area 41, 42***

Clinical Summary

Auditory physiology explains why patients hear the way they do.

Clinical audiology succeeds when physiology—not thresholds—guides interpretation.

Hearing is the process by which sound energy is collected, transmitted, amplified, and finally converted into electrical signals that can be interpreted by the brain.

Physiology of hearing can be divided into two main parts:

- ***Conductive part.***
- ***Sensory-neural part***

Each part performs a specific role, and failure at any level produces a predictable type of hearing loss.

A. Conductive Part

(Transmission of sound as mechanical vibrations)

The conductive part is responsible for delivering sound energy efficiently to the inner ear.

1. External Ear

Auricle (Pinna)

- ***Collects air-borne sound waves***
- ***Helps direct sound into the external auditory canal***
- ***Contributes to sound localization***

External Auditory Canal

- *Conducts sound waves toward the tympanic membrane*
- *Acts as a resonator*
- *Enhances speech frequencies*

Tympanic Membrane

- *Vibrates when sound waves reach it*
- *Converts air-borne sound waves into mechanical vibrations*

Clinical Insight

Any pathology affecting the external ear or tympanic membrane results in conductive hearing loss.

2. Middle Ear

The middle ear transmits and amplifies sound vibrations.

Functions of the Ossicular Chain

1. *Transmission of vibrations from the tympanic membrane to the oval window*
2. *Amplification of sound energy*

Mechanisms of Amplification

Areal Ratio

- *Large vibrating area of the tympanic membrane*
- *Small area of the oval window*
- *Ratio approximately 17 : 1*

Lever Action

- *Lever system formed by malleus and incus*
- *Lever ratio approximately 1.3 : 1*

Total Amplification

- *17 × 1.3 ≈ 22 times*
- *This compensates for sound transmission from air to cochlear fluid*
- *Known as the impedance matching mechanism*

Phase Difference

- *When the oval window moves inward, the round window moves outward*
- *Prevents excessive pressure buildup within the cochlea*


Malleus
Incus
Axis of rotation
Malleus longer
Incus shorter

▲ *Figure 7–5. Lever action of ossicles*


Length of long process of malleus
Length of long process of incus
= 1.3
2-Ossicular lever action
1.0
1.3
1
17
(56 sq mm)
14
45 sq mm
3.2 sq mm
Effective area ratio
14:1
Malleus and incus behave as 1st degree lever

▲ *Figure 7–6. Diagrammatic representation of ossicular lever mechanism*

3. Eustachian Tube

Functions

- ***Ventilates the middle ear***
- ***Equalizes pressure between the middle ear and atmosphere***
- ***Allows free movement of the tympanic membrane and ossicles***
- ***Drains middle ear secretions***


23 mm
6 mm
Isthmus
3 mm
5 mm
x
2 mm
Tympanic end
160°
Pharyngeal end

▲ *Figure 7–7. Structure and shape of Eustachian tube*


Ossicles
Tympanic membrane
Tympanic opening
Mucosa
Bony part
1/3
Isthmus
Nasopharyngeal
opening
Cartilaginous part
2/3
Tubal cartilage

▲ *Figure7–8. Eustachian tube anatomy. Horizontal section*


Bony portion of ET
Cartilaginous portion of ET
Isthmus of ET
Eustachian tube
Nasopharynx

▲ *Figure 7–9.* ***Eustachian tube (ET) connecting the middle ear cavity with the nasopharynx***


Elastin hinge
Greater wing of sphenoid
Tubal cartilage
Petrous temporal bone
Mandibular nerve
Auditory tube
Otic ganglion
Ostmann's pad of fat
Fibrous band
Tensor palati
Levator palati
Salpingopharyngeus

▲ *Figure 7–10.* ***Relations and structure of the cartilaginous part of eustachian tube.***


Elastin hinge in roof
Lateral lamina
Medial lamina
Fibrous membrane
Tubal cartilage
A
Tensor palati (Dilator tubae)
Elastin hinge in roof
Mucosa
Ostmann's pad of fat
Levator palati
B

▲ ***Figure 7–11. Eustachian tube (ET) physiology. Vertical section of ET. (A) Closed in resting position; (B) Contraction of tensor veli palatini (dilator tubae) opens the tube. Elastin roof helps in regaining resting closure position***



▲ ***Figure 7–12. (A) Lateral end of the Eustachian tube opening in the protympanum; (B) Medial end of Eustachian tube as seen by nasal endoscope (white arrows)***


Middle ear
Eustachian tube
Levator veli palatini muscle
Tensor veli palatini muscle
Soft palate

▲ *Figure 7–13. Muscles of Eustachian tube*


Cartilage
Lumen
Pull of tensor palati muscle

▲ *Figure 7–14. Mechanism of Eustachian tube opening*

Red Flag

Eustachian tube dysfunction leads to negative middle ear pressure and conductive hearing loss.

B. Sensory-Neural Part

(Conversion of mechanical energy into neural signals)

1. Cochlea

The cochlea is the sensory organ of hearing.

Mechanism of Sound Transduction

1. ***Stapes footplate vibrates at the oval window***
2. ***Cochlear fluids move***
3. ***Basilar membrane vibrates***
4. ***Hair cells in the organ of Corti bend***
5. ***Stereocilia deflection causes depolarization***
6. ***Mechanical energy is converted into electrical impulses***


Tectorial membrane
Inner hair cell
Stereocilia of outer hair cells
Shearing force
Basilar membrane
Nerve fibers
Auditory nerve

▲ ***Figure 7–15.The organ of Corti.***
The central axis of the spiraling cochlea is to the left of the drawing.
Eighth nerve fibers pass through a bony shelf (the spiral osseus lamina) on their way to the hair cells.


2,000 Hz
3,000 Hz
500 Hz
Apex
100 Hz
Base
Cochlear duct
4,000 Hz
1,000 Hz
20,000 Hz
Basilar membrane
10,000 Hz
5,000 Hz

▲ *Figure 2–16. Frequency of sound perception in cochlea*


Scala vestibuli
Stria vascularis
Scala media
Scala tympani

▲ *Figure 7–17. Cross section of the cochlea.*
There are 3 fluid-filled chambers: the scala vestibuli and scala tympani are connected at the apex of the cochlea and contain perilymph; the scala media contains endolymph.
The stria vascularis maintains the endolymphatic potential and drives the silent current (arrows) that provides the energy for hearing.


To vestibular system
Stapes
Acoustic energy
Reissner's membrane
Helicotrema
Scala vestibuli
Scala tympani (perilymph)
Round window
Cochlear partition
Scala media (endolymph)
A
Scala vestibuli
Section n + 1
Reissner's membrane
Pv
Scala media
Spring
mass
Friction
Pt
Scala tympani
Section n
B

▲ *Figure 7–18.The traveling wave.*
The basilar membrane varies in mass and stiffness along the length of the cochlea (here shown unrolled).
This creates a tonotopic organization in which different segments of the basilar membrane are most sensitive to different frequencies.
The pressure wave introduced from movement of the stapes propagates up the cochlea and is dissipated at its characteristic frequency place.
The cochlea can be modeled as having multiple sections,
each with a distinct mass and stiffness of the basilar membrane.


Plasma membrane
Cytoskeleton
Subsurface cisternae
Extracisternal space
Axial core
Cuticular plate
Particles
Pillar
Spectrin
Actin

▲ Figure 7–19. ***Anatomy of the outer hair cell.***
The outer hair cell is cylindrical and is divided in 3 parts.
The top part is capped with a cuticular plate (CP) into which the stereocilia are inserted. The base of the cell is hemispheric.
It contains the cell nucleus and synaptic structures (not shown).
The central part of the cell is cylindrical.



▲ Figure 7–20. ***Various forms of hair cell damage found with noise trauma and their effect on eighth nerve tuning curves.***
The dotted lines represent normal turning curves and the solid lines are pathologic.

2. Cochlear Nerve

- ***Transmits electrical impulses from hair cells***
- ***Signals travel to cochlear nuclei in the brainstem***
- ***Then to higher auditory centers in the temporal lobe***
- ***The brain interprets these signals as meaningful sound***

Developmental Physiology of Hearing

Auditory system development begins early in gestation:

- ***Inner ear structures form first***
- ***Cochlear morphology is largely complete before birth***
- ***Peripheral anatomy may appear "ready" long before function is mature***

However:

- ***Neural pathways continue to develop late into gestation***
- ***Synaptic organization is incomplete***
- ***Central auditory pathways are immature***

An anatomically formed ear does not equal a functionally mature system.

Neonatal Auditory Physiology

Physiological characteristics:

- ***Prolonged ABR latencies***
- ***Reduced neural efficiency***
- ***Limited temporal resolution***

Clinical implication:

- ***"Abnormal" results may reflect immaturity, not pathology***
- ***Interpretation must be age-appropriate***

Postnatal Maturation — Experience Shapes Physiology

Auditory physiology matures through:

- ***Sound exposure***
- ***Language input***
- ***Neural stimulation***

Key processes:

- ***Myelination***
- ***Synaptic refinement***
- ***Improved neural synchrony***
- ***Central auditory organization***

This maturation is experience-dependent.

Hearing develops by being used.

Impact of Early Hearing Loss on Development

Early auditory deprivation affects:

- ***Neural synchrony***
- ***Central auditory mapping***
- ***Speech and language development***
- ***Vestibular and balance integration***

Why Developmental Physiology Matters to Intervention Timing

Timing decisions (e.g., amplification, CI, monitoring) depend on:

- ***Physiological readiness***
- ***Neural plasticity***
- ***Developmental trajectory***

Otologic interventions aim to:

- ***Restore access before plasticity declines***
- ***Avoid intervention after irreversible central changes***

Good timing protects outcomes more than perfect thresholds.

Key Points

- ***Hearing consists of conductive and sensory-neural stages***
- ***The conductive system delivers and amplifies sound***
- ***The cochlea converts mechanical energy into neural signals***
- ***The auditory nerve carries these signals to the brain***

Golden Rule

Conductive mechanisms prepare the sound, but perception occurs only after neural transduction.

Chapter

8

Vestibular Physiology



How Balance Is Detected, Integrated, and Stabilized

Why This Chapter Matters

Balance is not a single sense.

It is an integrated physiological system that depends on precise timing, symmetry, and central integration.

Otologists worry about:

- *Stability*
- *Asymmetry*
- *Compensation*
- *Risk of irreversible imbalance*

Audiologists and audio-vestibular clinicians must therefore understand how balance works physiologically—before interpreting tests or recommending intervention.

In vestibular care, physiology explains symptoms better than labels.

1. The Vestibular System as a Functional Network

The vestibular system functions as a three-part network:

1. *Peripheral sensors*
2. *Central integration*

3. Motor output (eye, head, posture)

Dysfunction at any level may produce similar symptoms but very different clinical meanings.

2. Peripheral Vestibular Physiology — Motion Sensors, Not Balance Organs

Semicircular Canals

- Detect angular acceleration
- Three canals, orthogonally oriented
- Function in paired, push–pull fashion

Clinical relevance:

- Unilateral loss causes asymmetry
- Acute loss produces vertigo
- Chronic loss may appear “silent” after compensation


Eyes
External rectus muscles
Internal rectus muscles
To cerebral cortex
Oculomotor nucleus
Thalamus
Superior nucleus
Cerebellum
Lateral nucleus
Medial nucleus
Inferior nucleus
Vestibular labyrinth
Inferior cerebellar peduncle
Medial longitudinal bundle
Vestibular (Scarpa) ganglion
Vestibulospinal tract

▲ Figure 8–1. The central connection of the vestibular nerve


Superior semicircular canal
Bony labyrinth
Endolymph
Perilymph
Posterior semicircular canal
Membranous labyrinth
Hair cells
Ampullary crest
Crista ampullaris
Vestibular nerve (VIII)
Cupula
Ampullary sulcus

▲ *Figure 8–2.* ***The semicircular canals and ampullae.***

The hair cells sit on the crista, and their stereocilia are embedded in the gelatinous cupula.

Angular acceleration (head rotation) causes the endolymph within the semicircular canal to bend the cupula, resulting in stereociliary deflection.

The 3 semicircular canals (lateral, superior, and posterior) are perpendicular to one another, permitting detection of head rotation in any direction.


Semicircular canal
Endolymph
Ampulla
Cupula
Hair cell
Cristae
Vestibular nerve
Ampulla
Cupula
Kinocilium
Endolymph
Stereocilia
Support cells
Sensory hair cell
Vestibular nerve

▲ *Figure 8–3.* ***Structure of ampulla***



▲ Figure 8–4.Physiological pair of semicircular canals (SSC)

Otolith Organs (Utricle & Saccule)

- *Detect linear acceleration and gravity*
- *Respond to head tilt and translational movement*

Clinical relevance:

- *Otolith dysfunction affects orientation and posture*
- *Often under-recognized*
- *Explains imbalance without vertigo*

The vestibular system detects motion—not dizziness.


Utriculus
Striola
1
2
3
I
II
Sacculus
Striola
4
5
6

▲ *Figure 8–5.* ***Organization of the utricle and saccule.***

3. Vestibular Hair Cell Physiology

Vestibular hair cells:

- ***Convert mechanical motion into neural signals***
- ***Respond to acceleration, not position***

- *Adapt quickly to sustained stimuli*

Clinical implications:

- *Static tests may miss dynamic dysfunction*
- *Compensation masks peripheral loss*
- *Normal feeling does not equal normal physiology*


Otolithic membrane with otoconia
Hair cells
Supporting cells
Nerve cells
Force of gravity
Head erect resting membrane potential
Tilt toward the kinocilium: depolarization
Tilt away from the kinocilium: hyperpolarization

▲ *Figure 8–6. Organization and physiology of the maculae.*


Increased
Decreased
Saccule
C
D
Force of gravity
Resting tone
Decreased
A
B
Utricle
Endolymph

▲ *Figure 8–7. Nerve potential generation due to head positions and linear acceleration*

4. Vestibular Nerve and Neural Timing

Vestibular nerve function depends on:

- ***Neural synchrony***
- ***Bilateral symmetry***
- ***Accurate timing***

Clinical relevance:

- ***Asymmetry matters more than absolute loss***
- ***Central compensation alters symptom profile***
- ***Chronic patients may show minimal complaints despite deficit***

Otologists focus on risk of decompensation, not symptom intensity.

5. Central Vestibular Integration

Vestibular signals integrate with:

- ***Vision***
- ***Proprioception***
- ***Cerebellar processing***

Clinical implications:

- ***Central lesions mimic peripheral disorders***
- ***Visual dependence increases with vestibular loss***
- ***Poor central integration limits rehabilitation outcomes***

Balance is a central experience built on peripheral input.

6. Vestibulo-Ocular Reflex (VOR) — Stability Over Sensation

The VOR:

- ***Stabilizes vision during head movement***

- ***Operates subconsciously***
- ***Is the primary functional output of vestibular physiology***

Clinical relevance:

- ***VOR integrity predicts functional stability***
- ***Patients may complain of imbalance without vertigo***
- ***VOR deficits explain oscillopsia***

Otologists prioritize function, not symptom description.

7. Developmental Aspects of Vestibular Physiology

In children:

- ***Vestibular system matures early***
- ***Compensation is rapid***
- ***Deficits may present as motor delay, not dizziness***

Clinical implications:

- ***Vestibular loss may go unrecognized***
- ***Balance affects speech, posture, and attention***
- ***Early identification improves outcomes***

8. Why Vestibular Physiology Matters to Intervention

Vestibular physiology informs:

- ***Surgical risk assessment***
- ***CI planning***
- ***Labyrinthectomy considerations***
- ***Counseling and expectation setting***

Intervening without understanding vestibular physiology risks permanent imbalance.


No head rotation
MLF
III
III
Left
Right
VI
VI
VN
VN
Lateral semicircular canal
Lateral semicircular canal
Head rotates to right
Rotation
Slow phase
MLF
III
III
Left
Right
VI
VI
Endolymph flow
VN
VN
Lateral semicircular canal
Semicircular canal

▲ *Figure 8–8. The vestibuloocular reflex.*

Clinical Summary

Vestibular physiology is a dynamic, bilateral, and centrally integrated system.

Symptoms alone cannot define function or risk.

Equilibrium is the ability of the body to maintain balance, posture, and stable vision during movement.

Balance is not the function of the vestibular system alone.

It is the result of integration between:

- ***Vestibular system***
- ***Visual system***
- ***Proprioceptive (somatosensory) system***

Failure of any component, or poor integration between them, leads to dizziness and imbalance.

A. Vestibular System

(The sensory system for balance)

The vestibular system is located in the inner ear and detects head movement and position.

Main Functions

- ***Detect head movement***
- ***Detect head position relative to gravity***
- ***Stabilize gaze during movement***
- ***Maintain posture and balance***

B. Parts of the Vestibular System

1. Semicircular Canals

(Detection of angular or rotational movement)

There are three semicircular canals in each ear:

- ***Horizontal (lateral)***
- ***Anterior (superior)***
- ***Posterior***

They are arranged at right angles to each other.

Function

- ***Detect angular acceleration***
- ***Respond to head rotation, not to static position***

Mechanism

- ***Head rotation causes movement of endolymph***
- ***Endolymph deflects the cupula***
- ***Hair cells are stimulated or inhibited***
- ***Neural signals are generated***

Key Concept

Semicircular canals detect movement, not position.

2. Otolith Organs

(Detection of linear movement and gravity)

The otolith organs include:

- ***Utricle***
- ***Saccule***

Function

- ***Detect linear acceleration***

- ***Detect head position relative to gravity***

Mechanism

- ***Otoconia rest on a gelatinous membrane***
- ***Movement or tilt shifts the otoconia***
- ***Hair cells are deflected***
- ***Neural signals are produced***

Clinical Insight

Otolith dysfunction may cause dizziness even when semicircular canal tests are normal.

C. Vestibular Hair Cells and Signal Encoding

Vestibular hair cells:

- ***Have a constant resting firing rate***
- ***Increase firing when stimulated in one direction***
- ***Decrease firing when stimulated in the opposite direction***

This allows bidirectional signaling, which improves sensitivity and accuracy.

D. Vestibular Nerve and Central Connections

Vestibular Nerve

- ***Carries signals from semicircular canals and otolith organs***
- ***Joins the auditory nerve to form the vestibulocochlear nerve (CN VIII)***



▲ *Figure 8–8.* ***Deflection of the otoconial mass causes hair cell depolarization. This deflection is identical when the head is tilted backward (a) or when accelerating forward (b).***

Central Pathways

Vestibular signals project to:

- ***Vestibular nuclei in the brainstem***
- ***Cerebellum***
- ***Ocular motor nuclei***
- ***Spinal cord***


Otolithic membrane
Hair cells
Supporting cells
Nerve fibre

▲ *Figure 8–9. Hair cells of the otolith organs are located at the macula. The tips of these hair cells are embedded in the otolithic membrane (a gelatinous mass covered with calcium carbonate crystals, the so-called otoliths or otoconia).*

E. Vestibulo-Ocular Reflex (VOR)

Purpose

The VOR stabilizes vision during head movement.

Mechanism

- ***Head moves in one direction***
- ***Eyes move in the opposite direction***

- *Visual image remains stable on the retina*

Clinical Importance

- *Allows clear vision during walking or running*
- *Tested clinically by head impulse tests and caloric testing*

Red Flag

Abnormal VOR leads to blurred vision during head movement (oscillopsia).

F. Vestibulo-Spinal Reflexes

(Postural control)

Vestibular input helps:

- ***Maintain upright posture***
- ***Adjust muscle tone***
- ***Prevent falls***

These reflexes work with proprioceptive input from muscles and joints.

G. Central Integration and Compensation

The brain continuously integrates:

- ***Vestibular input***
- ***Visual input***
- ***Proprioceptive input***

Vestibular Compensation

- *Central nervous system adapts to vestibular loss*

Absence of dizziness does not mean normal vestibular function.



Chapter

8

Pharmacology of Otologic Drugs

How Medications Interact with Hearing and Balance Physiology

Why This Chapter Matters

Many changes in hearing and balance are not caused by disease progression.

They are caused by drugs.

Otologists constantly ask:

- ***Is this change pathological—or pharmacological?***
- ***Is it reversible—or toxic?***
- ***Is timing critical?***

Audiologists often detect the change first.

Understanding otologic pharmacology allows clinicians to:

- ***Recognize drug-related effects early***
- ***Prevent irreversible damage***
- ***Interpret audiological changes safely***

Some hearing losses are prescribed.

1. How Drugs Reach the Inner Ear

Otologic drugs reach the inner ear through:

- ***Systemic circulation***

- ***Blood–labyrinth barrier***
- ***Local diffusion (e.g., intratympanic routes)***

Key concepts:

- ***The inner ear is partially protected—but not isolated***
- ***Drug concentration may accumulate***
- ***Clearance is slow***

Clinical implication:

- ***Toxicity may appear delayed***
- ***Effects may persist after drug cessation***

2. Ototoxicity — A Pharmacological Reality

Ototoxic drugs affect:

- ***Cochlear hair cells***
- ***Vestibular hair cells***
- ***Neural elements***

Major ototoxic drug classes include:

- ***Aminoglycoside antibiotics***
- ***Platinum-based chemotherapeutics***
- ***Loop diuretics (dose-related)***
- ***Salicylates (usually reversible)***

Not all ototoxicity is predictable—but much is preventable.

3. Cochlear vs Vestibular Toxicity

Some drugs are predominantly:

- ***Cochleotoxic → hearing loss, tinnitus***
- ***Vestibulotoxic → imbalance, oscillopsia***

Clinical relevance:

- ***Absence of dizziness does not exclude toxicity***
- ***Vestibular loss may be masked by compensation***
- ***Hearing may decline silently***

Audiological monitoring is often the only early warning.

4. Dose, Duration, and Individual Vulnerability

Pharmacological risk depends on:

- ***Cumulative dose***
- ***Treatment duration***
- ***Renal function***
- ***Genetic susceptibility***

Clinical implications:

- ***"Safe dose" is not universal***
- ***Progressive loss may occur despite monitoring***
- ***High-risk patients require baseline and follow-up testing***

Ototoxicity is a risk curve—not an on/off switch.

5. Reversible vs Irreversible Drug Effects

Some pharmacological effects are:

- ***Temporary (e.g., salicylates)***
- ***Dose-dependent***
- ***Functionally reversible***

Others are:

- ***Progressive***
- ***Permanent***

- ***Cumulative***

Otologists prioritize early recognition over late confirmation.

6. Pharmacology and Vestibular Function

Drugs affecting vestibular physiology may:

- ***Suppress symptoms***
- ***Mask compensation***
- ***Delay central adaptation***

Common examples:

- ***Vestibular suppressants***
- ***Sedatives***
- ***Antiemetics***

Clinical implication:

- ***Symptom relief ≠ physiological recovery***
- ***Prolonged use may hinder rehabilitation***

7. Why Pharmacology Matters in Surgical and MDT Decisions

Drug effects may:

- ***Mimic disease progression***
- ***Confound candidacy assessment***
- ***Alter baseline measurements***
- ***Affect postoperative outcomes***

Audiologists must document:

- ***Medication history***
- ***Timing of changes***

- *Dose-related patterns*

Pharmacology shapes interpretation before it shapes treatment.

Clinical Summary

Otologic pharmacology influences hearing and balance through dose-dependent, time-dependent, and individual-specific mechanisms.

Key Points

- *Drugs can alter auditory and vestibular physiology*
- *Monitoring protects outcomes*
- *Symptom relief does not equal safety*

Common Mistakes

- *Ignoring medication history*
- *Delayed monitoring*

"Some medicines can affect hearing and balance.

That's why we monitor—not because something went wrong."

Teaching Pearl

Prevention is the only treatment for ototoxicity.

Final Clinical Message

Pharmacology must always be part of auditory interpretation.

Take-Home Messages

- *Ask about drugs*
- *Monitor early*
- *Act before damage accumulates*

SECTION 3

Outer & Middle Ear Pathologies and Management

Outer Ear Pathologies and Management

From Surface Findings to Clinical Consequences

Why This Part Matters

Outer ear disorders are often labeled as **minor** or **simple**.

They are rarely treated as decision-shaping conditions.

Otologists know better.

The outer ear is:

- The first point of access
- The first barrier to assessment
- Often the hidden cause of diagnostic error

Audiologists frequently encounter:

- Abnormal findings dismissed as incidental
- Test results distorted by external ear pathology
- Delayed or inappropriate referrals

This part reframes outer ear conditions as **foundational clinical modifiers**, not background noise.

If sound cannot enter properly, everything downstream becomes misleading.

How Otologists Think About the Outer Ear

Otologists do not see the outer ear as:

- A skin problem

- A canal issue
- A temporary inconvenience

They see it as:

- An access issue
- A stability issue
- A risk factor
- A potential contraindication

Audiological interpretation must align with that thinking.

What This Part Will Cover



PART II focuses on:

- Common and uncommon outer ear pathologies
- Their physiological impact on hearing assessment
- How they distort audiological data
- When they require treatment, monitoring, or referral
- How management decisions affect downstream care

This part is not about memorizing conditions.

It is about knowing when outer ear pathology changes meaning.

How to Read This Part

When reading each chapter, ask:

- Does this condition alter sound transmission?
- Does it affect test reliability?
- Is it stable or progressive?
- Does it require resolution before further assessment?

Otologists always ask:

Can I trust the data collected through this ear?

Each chapter in this part will follow:

- ***Pathophysiology overview***
- ***Clinical presentation***
- ***Audiological impact***
- ***Management principles***
- ***Decision points for referral or delay***

And will close with the Framework

focused on misinterpretation risk and clinical judgment.

Chapter



9

Disorders of the Auricle

When External Appearance Reflects Deeper Risk

Why This Chapter Matters

Disorders of the auricle are often viewed as:

- ***Cosmetic***
- ***Congenital curiosities***
- ***Minor findings***

In otologic thinking, this is incomplete.

The auricle can signal:

- ***Underlying developmental anomalies***
- ***Associated middle or inner ear pathology***
- ***Syndromic conditions***
- ***Increased surgical or audiological risk***

For audiologists, auricular findings are not background details.

They are clinical clues.

The ear you see may reveal the ear you cannot test.

1. Congenital Auricular Anomalies

Microtia

- ***Partial or complete underdevelopment of the auricle***

Clinical relevance:

- ***Frequently associated with external auditory canal atresia***
- ***Often accompanied by middle ear malformations***
- ***Inner ear involvement may coexist***

Audiological implications:

- ***Air-conduction results are unreliable***
- ***Bone-conduction must be interpreted carefully***
- ***Early referral and imaging are essential***



▲ ***Figure 9–1. Classification of microtia.***



▲ *Figure 9–2. Type 3 microtia.*



▲ *Figure 9–3. Multistaged auricular reconstruction. In the first stage, a large piece of cartilage is harvested from the rib (a), and the auricular framework is carved to resemble the normal landmarks of the pinna (b). This framework is placed under the skin (c). In a second stage, retroauricular sulcus and tragus are created using the auricular remnant (d).*



▲ Figure 9–4. Auricular prosthesis.

Anotia

- *Complete absence of the auricle*

Clinical relevance:

- *High likelihood of canal atresia*
- *Middle ear anomalies almost always present*

Audiological implications:

- *Conventional testing pathways are limited*
- *Early rehabilitation planning is required*

2. Preauricular Anomalies

Preauricular Tags and Pits

Often dismissed as isolated findings.

Clinical relevance:

- *May be associated with syndromes (e.g., branchio-oto-renal spectrum)*
- *Can indicate abnormal embryological development*

Audiological implications:

- *Baseline hearing assessment is mandatory*
- *Renal and genetic considerations may be relevant*

Small external findings can signal systemic involvement.

3. Acquired Auricular Disorders

Perichondritis

- *Inflammation or infection of auricular cartilage*

Clinical relevance:

- *Risk of cartilage necrosis and deformity*
- *May follow trauma or piercing*

Audiological implications:

- *Pain limits testing*
- *Infection must be managed before reliable assessment*



▲ *Figure 9–5. Relapsing perichondritis leading to cauliflower ear.*

Trauma and Hematoma

- Blunt trauma leading to auricular hematoma

Clinical relevance:

- Risk of "cauliflower ear" if untreated
- May mask deeper injury

Audiological implications:

- Delay testing until stabilization
- Document trauma history carefully

4. Syndromic Associations

Auricular abnormalities may be part of:

- Craniofacial syndromes
- Genetic hearing loss conditions
- Developmental anomalies affecting multiple systems

Audiological role:

- Identify patterns
- Prompt multidisciplinary referral
- Avoid isolated interpretation of hearing results

5. Why Auricular Disorders Matter to Management

Otologists assess auricular disorders as:

- Markers of developmental risk
- Indicators of possible surgical complexity
- Signals for further investigation

Audiologists contribute by:

- Documenting external findings

- *Adjusting testing strategies*
- *Communicating risk clearly*

Ignoring the auricle risks missing the diagnosis.

Clinical Summary

Auricular disorders often reflect deeper otologic or systemic abnormalities and must never be treated as purely cosmetic findings.

Diseases of the auricle can be classified into four main groups:

- *Congenital*
- *Traumatic*
- *Inflammatory*
- *Neoplastic*

This classification helps in systematic diagnosis and management.

A. Congenital Diseases

The auricle develops from six tubercles arising from the first and second branchial arches.

Any disturbance in development leads to congenital deformities.

1. Defects in Size

Anotia

- *Complete absence of the auricle*
- *May be unilateral or bilateral*
- *Commonly associated with:*
    - *External auditory canal atresia*
    - *Middle ear abnormalities*

Microtia

- *Small, malformed auricle*
- *May be associated with conductive hearing loss*

Macrotia

- *Excessively large auricle*

Treatment

- *Plastic reconstruction*
- *Preferably performed before school age*

2. Bat Ear (Protruding Ear)

- *Abnormally protruding auricle*
- *Large concha*
- *Poorly developed antihelix and scapha*

Treatment

- *Surgical correction before school age*



▲ *Figure 9–6. (A) Preoperative; (B) Postoperative photographs of two brothers with protruding ears*

3. Accessory Auricle

- *Skin covered tags*
- *Located along a line from:*
    - *Tragus*
    - *To the angle of the mouth*
- *May contain cartilage*

Treatment

- *Surgical removal if cosmetically undesirable*

4. Preauricular Pit or Sinus

- *Small depression anterior to the crus of helix or above the tragus*
- *Represents a blind epithelial tract*

Clinical Features

- *Recurrent infection*
- *Purulent discharge*
- *Pain and swelling*
- *May form a cyst or abscess*

Treatment

- *Acute infection:*
    - *Drainage and antibiotics*
- *Recurrent infection or cosmetic concern:*
    - *Surgical excision of the tract*



▲ Figure 9–7. Left preauricular sinus with inflammation of the overlying skin.



▲ Figure 9–8. Preauricular appendages. (a) Associated with a mildly dysplastic right pinna; (b) associated with microtia of the left pinna; (c) associated with an otherwise normally developed pinna.


[redacted]

▲ *Figure 9–9. Dysplastic left pinna in a girl with CHARGE syndrome.*


[redacted]

▲ *Figure 9–10. Unilateral right microtia in a boy with Goldenhar syndrome. He has right hemifacial microsomia and a left epibulbar dermoid.*



▲ *Figure 9–11. Child with Treacher Collins syndrome and microtia with CCA. The child is wearing a BAHA on a softband and has a tracheostomy.*


1
2
3
A
B
Processor
Abutment
Implant
C

▲ *Figure 9–12. Bone-anchored hearing aid (BAHA). (A) BAHA system; (B) BAHA parts; and (C) BAHA processor in position*

B. Traumatic Diseases

1. Auricular Hematoma

Definition

- ***Collection of blood between:***
    - ***Auricular cartilage***
    - ***Perichondrium***

Cause

- ***Blunt trauma, commonly in boxers***

Clinical Features

- ***Fluctuant swelling***
- ***Painful but not tender***

Complications

- ***Organization of clot leading to cauliflower ear***
- ***Secondary infection causing perichondritis***

Treatment

- ***Antibiotics***
- ***Aspiration under aseptic conditions***
- ***Pressure dressing***
- ***Incision and drainage if aspiration fails***

2. Lacerations of the Auricle

- ***Range from minor cuts to complete avulsion***

Causes

- ***Sharp trauma***
- ***Bites***
- ***Forceful pulling***

Management

- ***Early repair***
- ***Preservation of perichondrium***
- ***Antibiotics to prevent infection and deformity***

3. Frostbite

Cause

- ***Exposure to very cold temperatures***

Pathology

- ***Vasoconstriction followed by ischemia and necrosis***

Clinical Features

- ***Erythema and edema***
- ***Skin necrosis in severe cases***
- ***Pale, numb auricle***

Treatment

- ***Gradual rewarming***
- ***Hot compresses***
- ***Medical treatment with antibiotics and analgesics***
- ***Surgical debridement if necessary***

C. Inflammatory Diseases

1. Perichondritis

Definition

- ***Inflammation of the perichondrium of auricular cartilage***

Causes

- ***Infected hematoma***
- ***Infected lacerations***
- ***Spread from otitis externa***

Clinical Features

- ***Red, hot, painful auricle***

- *Abscess formation*
- *Lobule usually spared*

Complications

- *Cartilage necrosis*
- *Cauliflower ear deformity*

Treatment

- *Systemic and local antibiotics*
- *Analgesics*
- *Incision and drainage if abscess forms*

2. Relapsing Polychondritis

- *Rare autoimmune disease*
- *Involves auricular cartilage and other cartilages:*
    - *Nasal*
    - *Laryngeal*
    - *Tracheal*

Clinical Feature

- *Painful inflamed auricle with sparing of the lobule*

Treatment

- *High dose systemic steroids*

D. Neoplastic Diseases

Benign Tumors

- *Papilloma*
- *Nevus*

- *Chondroma*

Malignant Tumors

Basal Cell Carcinoma

- ***Common in elderly males***
- ***Presents as an ulcer with rolled edges***
- ***Rare lymph node metastasis***

Treatment

- ***Local excision with safety margin***
- ***Local flap reconstruction***

Squamous Cell Carcinoma

- ***Presents as:***
    - ***Painless nodule***
    - ***Ulcer with raised everted edges***
- ***Late lymph node metastasis***
- ***Common in sun exposed individuals***

Treatment

- ***Surgical excision with adequate margins***
- ***Postoperative radiotherapy***
- ***Neck dissection if lymph nodes are involved***

Golden Rule

Early recognition and proper management of auricular diseases prevent permanent deformity and functional loss.

Chapter



10

External Auditory Canal Disorders

When Access Determines Accuracy

Why This Chapter Matters

The external auditory canal is often treated as a simple passage for sound. In reality, it is a critical modifier of every audiological test.

Otologists never ask first:

- *How bad is the hearing loss?*

They ask:

- *Can sound reach the tympanic membrane reliably?*
- *Is the canal stable, inflamed, narrowed, or obstructed?*

Audiological data collected through a compromised canal is conditionally valid at best—and misleading at worst.

If access is compromised, interpretation must pause.

1. Cerumen (Ear Wax) — The Most Underestimated Variable

Cerumen is not a disease.

But it is one of the most common causes of:

- *Conductive hearing loss*

- *Test variability*
- *False asymmetry*

Clinical Characteristics

- *Partial or complete obstruction*
- *May fluctuate over time*
- *Often asymptomatic*

Audiological Implications

- *Artificial air–bone gaps*
- *Reduced speech clarity*
- *Unreliable high-frequency thresholds*

Even partial occlusion can:

- *Distort results*
- *Mask underlying pathology*

Cerumen converts physiology into artifact.



▲ *Figure 10–1. Ear syringing. Water jet of ear syringe taking out lump of wax*


Suction
Cerumen
A
Blunt curet
Cerumen
B
C
Irrigating syringe
Cerumen

▲ *Figure 10–2. (A) Wax removal by suction; (Bl Cleaning by hooking; (C) Syringing*

2. Otitis Externa — Inflammation Alters Interpretation

Otitis externa involves:

- *Canal skin inflammation*
- *Edema*
- *Pain*
- *Discharge*

Clinical Impact

- *Sound attenuation*
- *Patient discomfort limiting cooperation*
- *Risk of spread in severe cases*

Audiological Implications

- *Testing during active infection is unreliable*
- *Pain alters behavioral responses*
- *Tympanometry may be impossible or misleading*

Otologists prioritize treating inflammation before interpreting hearing.



▲ *Figure 10–3. Acute otitis externa. Erythema, edema, and copious purulent/fibrinous debris are seen in the left image (A). After otoendoscopic debridement, canal edema and erythema are seen extending to the tympanic membrane (B), and the ear drum is intact (C,arrow).*

3. External Auditory Canal Stenosis

Canal stenosis may be:

- *Congenital*
- *Acquired (post-infectious, post-traumatic)*

Clinical Relevance

- *Reduced canal diameter*
- *Chronic debris accumulation*
- *Increased infection risk*

Audiological Implications

- *Persistent conductive components*
- *Difficulty with probe placement*
- *Long-term instability*

Stenosis is a structural limitation, not a temporary finding.

4. Exostosis and Osteoma

Exostosis

- *Broad-based bony growths*
- *Often bilateral*
- *Associated with cold water exposure*

Osteoma

- *Solitary, pedunculated lesion*
- *Usually unilateral*

Clinical Implications

- *Progressive canal narrowing*
- *Trapped moisture and debris*
- *Increased infection risk*

Audiological Implications

- *Fluctuating conductive loss*
- *Progressive access limitation*
- *Inconsistent results over time*

Bony canal disease causes dynamic—not static—hearing changes.



▲ ***Figure 10–4. Coronal view of superior and inferior exostoses of the external auditory canal.***



▲ *Figure 10–5. (a) Exostoses. (b) Exostoses coexisting with osteoma (red stars).*



▲ *Figure 10–6. External ear cholesteatoma*

5. Foreign Bodies and Canal Trauma

Foreign bodies may be:

- ***Organic***
- ***Inorganic***
- ***Iatrogenic***

Clinical relevance:

- ***Acute conductive loss***
- ***Risk of infection or trauma***

Audiological implications:

- ***Testing should be deferred***
- ***Removal precedes interpretation***
- ***Document pre- and post-removal status***



▲ ***Figure 10–7. Removal of foreign body by blunt curet***

6. Why Canal Disorders Matter to Management Decisions

Otologists consider canal pathology when deciding:

- *Timing of assessment*
- *Surgical feasibility*
- *Risk of recurrence*
- *Reliability of audiological data*

Audiologists support this by:

- *Documenting canal status clearly*
- *Avoiding definitive conclusions through compromised access*
- *Recommending reassessment after resolution*

Management decisions depend on stable access, not rushed data.

Clinical Summary

External auditory canal disorders distort sound transmission and compromise the reliability of audiological assessment.

Diseases of the external auditory canal can be classified into four main groups:

- *Congenital*
- *Traumatic*
- *Inflammatory*
- *Neoplastic*

This classification helps in systematic understanding and management.

A. Congenital Diseases

Atresia of the External Auditory Canal

Definition

Congenital atresia of the external auditory canal may occur:

- ***Alone***
- ***Or associated with microtia***

When isolated, it results from failure of canalization of the ectodermal core of the first branchial cleft.

Features

- ***Usually unilateral***
- ***May be bony or fibrous***
- ***Conductive hearing loss is present***
- ***Sensorineural or mixed loss may occur if inner ear anomalies coexist***

Investigations

- ***CT scan of the temporal bone to assess:***
    - ***Middle ear condition***
    - ***Inner ear status***
    - ***Length of the atretic segment***
- ***Full audiological evaluation***

Treatment

- ***Bilateral cases:***
    - ***Hearing aids to preserve hearing***
- ***Surgical:***
    - ***Canalomeatoplasty***
    - ***Auricular reconstruction if microtia is present***
- ***Alternative option:***

  - ***Bone-anchored hearing aid (BAHA)***

B. Traumatic Diseases

1. Lacerations of the Canal Skin

Minor Lacerations

- ***Caused by scratching with:***
  - ***Hair pins***
  - ***Needles***
  - ***Matchsticks***
- ***Or unskilled instrumentation***
- ***Usually heal without sequelae***

Major Lacerations

- ***Caused by:***
  - ***Gunshot injuries***
  - ***Road traffic accidents***
  - ***Severe trauma***
- ***May lead to canal stenosis as a complication***

2. Foreign Body in the EAC

Common Patients

- ***Children***
- ***Mentally challenged adults***

Types

- ***Animate, insects***

- ***Inanimate:***
    - ***Vegetable, beans, seeds***
    - ***Non-vegetable, cotton, beads, paper***

Clinical Features

- ***History of foreign body insertion***
- ***Irritation or noise sensation***
- ***Conductive hearing loss if the canal is obstructed***

Complications

- ***Injury to EAC or tympanic membrane***
- ***Otitis externa***
- ***Otitis media if the drum is perforated***

Management

- ***Animate foreign body:***
    - ***Kill first with oil drops, then irrigation***
- ***Vegetable foreign body:***
    - ***Removal by hook or suction***
    - ***No irrigation***
- ***Non-vegetable foreign body:***
    - ***Hook, suction, or gentle irrigation***
- ***Uncooperative child:***
    - ***Removal under general anesthesia***



▲ **Figure 10–8.** Right ear, foreign body of the EAC; the patient was an adult who lost the adapter after taking out his hearing aid.

C. Inflammatory Diseases

(Otitis Externa)

Otitis externa is inflammation of the skin of the external auditory canal.

I. Infective Otitis Externa

A. Viral

Herpes Simplex

- *Occurs during febrile illnesses*
- *Vesicles in EAC, tympanic membrane, or nose*
- *Rupture and heal without scarring*

Herpes Zoster Oticus (Ramsay Hunt Syndrome)

- ***Severe otalgia***
- ***Vesicular eruption around the ear***
- ***Facial nerve palsy***
- ***Sensorineural hearing loss***
- ***Vertigo***

B. Fungal Otitis Externa (Otomycosis)

Causes

- ***Aspergillus niger***
- ***Candida albicans***

Predisposing Factors

- ***Hot, humid climate***
- ***Excessive sweating***
- ***Prolonged topical antibiotic use***

Clinical Features

- ***Intense itching***
- ***Otorrhea***
- ***Conductive hearing loss***
- ***Fungal debris resembling wet newspaper***

Treatment

- ***Removal of fungal debris***
- ***Topical antifungal drops***



▲ Figure 10–9. fungal OE (otomycosis) showing fungal hyphae of Aspergillus niger.

C. Bacterial Otitis Externa

1. Acute Localized Otitis Externa (Furuncle)

- *Staphylococcal infection of a hair follicle*
- *Occurs in the outer cartilaginous part*

Symptoms:

- *Severe pain, increased by chewing*
- *Local swelling and tenderness*

Treatment:

- *Systemic antibiotics and analgesics*
- *Local heat*
- *Incision and drainage if abscess forms*

2. Acute Diffuse Otitis Externa

- *Diffuse inflammation of the EAC*

Predisposing factors:

- ***Swimming***
- ***Scratching***
- ***Diabetes***
- ***Hot humid environment***

Signs:

- ***Diffuse edema and redness***
- ***Canal may be blocked***
- ***Tympanic membrane usually normal***

Treatment:

- ***Systemic antibiotics and analgesics***
- ***Local cleaning by dry methods***
- ***Topical antibiotic-steroid preparations***

3. Malignant Otitis Externa

(Skull Base Osteomyelitis)

- ***Occurs mainly in elderly uncontrolled diabetics***
- ***Caused by Pseudomonas aeruginosa***

Clinical Features:

- ***Severe persistent deep ear pain***
- ***Granulation tissue in the canal***
- ***Cranial nerve involvement***

- ***Facial nerve paralysis***

Treatment:

- ***Hospitalization***
- ***Strict diabetic control***
- ***Long-term systemic antibiotics***
- ***Surgical debridement if needed***

D. Neoplastic Diseases

Benign Tumors

- ***Exostosis***
- ***Osteoma***

Exostosis:

- ***Multiple broad-based bony swellings***
- ***Associated with cold water exposure***
- ***Causes canal obstruction***

Osteoma:

- ***Solitary, pedunculated lesion***

Treatment:

- ***Surgical removal if symptomatic***



▲ Figure 10–11. (C and D) Diagrammatic representation of osteoma arising from bony canal wall

Malignant Tumors

Squamous Cell Carcinoma

- *Originates deep in the canal*
- *Presents with:*
    - *Persistent discharge*
    - *Hearing loss*
- *Difficult to differentiate from middle ear carcinoma*

Treatment:

- *Surgical excision*
- *Radiotherapy when indicated*

Golden Rule

Persistent ear pain or discharge in the external canal must never be ignored.

Ear Wash (Aural Irrigation)

Ear wash is a simple clinical procedure used to remove material from the external auditory canal using warm water.

When done correctly, it is safe.

When done improperly, it may cause serious complications

Indications

Ear wash is indicated in the following situations:

1. Impacted Wax

- *Excessive or impacted cerumen causing:*
    - *Hearing loss*
    - *Ear fullness*
    - *Obstructed view of the tympanic membrane*
- *Note*

  Wax should be softened first using glycerin or sodium bicarbonate drops.



▲ *Figure 10–12. Method of syringing*



▲ *Figure 10–13. Right ear; ear wax partially occluding the EAC. The patient was referred with a scratchy noise over his right ear, symptoms were relieved after removal.*

2. Foreign Body in the EAC

Ear wash can be used only if the foreign body is:

- *Not impacted*
- *Not vegetable matter*
- *Animate foreign body, after killing it first with oil*

3. Otomycosis

- *Used to remove fungal debris*
- *Dry methods are preferred, but gentle ear wash may be used if needed*

4. Caloric Test

- *Ear wash using warm or cold water*
- *Used to assess vestibular function*



▲ *Figure 10–14. Caloric test*

Contraindications

Ear wash must NOT be done in the following conditions:

1. Tympanic Membrane Perforation

- *Traumatic*
- *Pathological, such as chronic suppurative otitis media*

2. Foreign Body

- *Vegetable foreign body, because it swells with water*

- *Impacted foreign body*

3. After Ear Surgery

- *Risk of middle or inner ear complications*

4. Otitis Externa

- *Bacterial*
- *Viral*

5. Fistula Between Middle and Inner Ear

- *May cause labyrinthitis and severe vertigo*

Technique of Ear Wash

1. *The patient is seated comfortably*
2. *A towel is placed to protect clothes*
3. *A kidney dish or basin is placed under the ear*
4. *The ear syringe is filled with sterile warm water*
    - *Temperature around 37°C*
    - *To avoid vertigo*
5. *The auricle is pulled:*
    - *Upward*
    - *Backward*
    - *Laterally*

    To straighten the external auditory canal
6. *The syringe nozzle is directed toward the posterosuperior wall of the canal*
    - *Not directly at the tympanic membrane*
7. *Water is injected gently and steadily*

Complications

1. Traumatic Rupture of the Tympanic Membrane

- ***Caused by:***
  - ***Wrong direction of water stream***
  - ***Excessive pressure***

Symptoms

- ***Sudden severe pain***
- ***Vertigo***
- ***Bleeding***
- ***Tinnitus***
- ***Hearing loss***
- ***Sensation of water trickling in the throat***

2. Trauma to the Canal Skin

- ***Caused by the syringe nozzle***

3. Vertigo

- ***Due to:***
  - ***Too hot water***
  - ***Too cold water***

4. Reflex Cough or Vasovagal Attack

- ***Due to stimulation of the auricular branch of the vagus nerve***

5. Otitis Externa

- ***From contaminated instruments or water***

6. Otitis Media

- ***If ear wash is done with an undetected TM perforation***

7. Further Impaction of Foreign Body

- *Caused by faulty technique*

Key Points

- *Ear wash is useful but not always safe*
- *Proper indication and contraindication must be respected*
- *Warm water and correct direction are essential*
- *Dry methods are preferred in many situations*

Golden Rule

Never perform ear wash unless the tympanic membrane is known to be intact.

11

Aural Atresia

When the Canal Does Not Exist — and Assumptions Become Dangerous

Why This Chapter Matters

Aural atresia is not simply:

- ***A closed ear canal***
- ***A conductive hearing loss***
- ***A cosmetic or surgical problem***

It is a developmental condition with implications that extend beyond the external auditory canal.

Otologists view aural atresia as:

- ***A marker of embryological disruption***
- ***A predictor of middle ear anatomy***
- ***A determinant of surgical risk and outcome***

Audiologists must view it as:

- ***A limitation of access***
- ***A modifier of testing strategy***

- *A condition that demands cautious interpretation*

In aural atresia, what you cannot access matters more than what you measure.

1. Definition and Developmental Basis

Aural atresia refers to:

- *Congenital absence or severe narrowing of the external auditory canal*

It results from:

- *Failure of canalization during embryologic development*
- *Often associated with auricular anomalies (e.g., microtia)*

Developmental implications:

- *External canal, middle ear, and ossicles often share embryological disruption*
- *Inner ear is frequently normal, but not guaranteed*

2. Types of Aural Atresia

Complete Atresia

- *No visible external auditory canal*
- *Usually associated with microtia*

Partial Atresia / Severe Stenosis

- *Narrow canal*
- *Often unstable, debris-retaining, and infection-prone*

Clinical relevance:

- *Partial patency does not equal functional access*
- *Stenotic canals behave like dynamic obstructions*

3. Audiological Characteristics

Typical audiological profile:

- *Significant conductive hearing loss*
- *Bone conduction often within normal limits*
- *Masking challenges in unilateral cases*

However:

- *Bone conduction thresholds may not reflect true cochlear reserve*
- *Cross-hearing and vibrotactile responses must be considered*
- *Behavioral responses may be misleading in children*

In atresia, "good bone conduction" is not a diagnosis—it is a hypothesis.

4. Middle Ear and Ossicular Considerations

Otologists assume that:

- *Middle ear anomalies are common*
- *Ossicular discontinuity or fixation is likely*
- *Facial nerve course may be aberrant*

Audiological implication:

- *Air–bone gap magnitude does not predict surgical success*
- *Stability matters more than degree*

5. Imaging and Multidisciplinary Assessment

High-resolution CT is essential to:

- *Assess middle ear space*
- *Identify ossicular anatomy*
- *Evaluate facial nerve position*
- *Determine surgical candidacy*

Audiologists must:

- ***Avoid definitive recommendations without imaging***
- ***Frame reports as functional assessments, not surgical opinions***

6. Unilateral Aural Atresia — The Silent Trap

Unilateral atresia is often underestimated.

Clinical risks include:

- ***Delayed speech-in-noise difficulties***
- ***Localization deficits***
- ***Academic and attentional impact***

Audiological challenges:

- ***Masking errors***
- ***Overestimation of "normal ear compensation"***

One normal ear does not guarantee normal auditory development.

7. Management Pathways — What Audiology Should and Should Not Do

Audiology should:

- ***Document functional impact***
- ***Recommend early access to sound (e.g., bone conduction devices)***
- ***Monitor development longitudinally***

Audiology should not:

- ***Predict surgical outcomes***
- ***Label candidacy prematurely***
- ***Frame atresia as "just conductive"***

Otologic decisions require MDT input and imaging.

Clinical Summary

Aural atresia is a developmental condition affecting access, anatomy, and decision-making—not merely a conductive hearing loss.

Congenital aural atresia is a developmental absence or severe narrowing of the external auditory canal, usually associated with malformation of the middle ear structures.

It is present at birth and may be unilateral or bilateral.

Epidemiology

- ***Incidence approximately 1 in 10,000 to 20,000 births***
- ***More common in males***
- ***Unilateral cases are more common than bilateral***
- ***Right ear is affected more frequently***

Embryology and Pathogenesis

- ***Results from abnormal development of the first and second branchial arches***
- ***External ear malformation is often associated***
- ***Middle ear ossicles may be malformed or fused***
- ***Inner ear is usually normal***

Classification

According to Canal Development

- ***Canal stenosis***
- ***Partial atresia***
- ***Complete atresia***

According to Laterality

- *Unilateral congenital aural atresia*
- *Bilateral congenital aural atresia*

Associated Anomalies

- *Microtia*
- *Ossicular chain malformations*
- *Facial nerve anomalies*
- *Craniofacial syndromes*

Inner ear anomalies are uncommon but must be excluded.

Audiologic Features

- *Hearing loss is typically conductive*
- *Degree of loss usually moderate to severe*
- *Bone conduction thresholds are usually normal*
- *Unilateral cases may be missed without screening*

Audiologic Assessment

Infants and Young Children

- *Auditory Brainstem Response (bone conduction)*
- *Otoacoustic emissions if feasible*
- *Behavioral observation audiometry*

Older Children and Adults

- *Pure tone audiometry*
- *Speech audiometry*
- *Bone conduction testing is essential*

Imaging

- ***High resolution CT of the temporal bone***
- ***Assesses:***

– Middle ear space

– Ossicular development

– Facial nerve course

– Cochlear integrity

Imaging is essential before any surgical decision.

Management Principles

Management depends on:

- ***Laterality***
- ***Degree of hearing loss***
- ***Age of patient***
- ***Middle ear anatomy***

Audiologic Management

Early Hearing Rehabilitation

- ***Bone conduction hearing devices***
- ***Soft band bone conduction aids in infants***
- ***Early amplification is critical for language development***

Unilateral Atresia

- ***Hearing rehabilitation is still recommended***
- ***Educational and listening difficulties may occur***
- ***Classroom accommodations advised***

Surgical Management

- ***Canalplasty or atresiaplasty***
- ***Usually delayed until older childhood***
- ***Surgical candidacy depends on CT findings***
- ***Hearing outcome is variable***

Audiology must provide realistic expectations.

Prognosis

- ***Hearing outcome varies***
- ***Bone conduction devices provide reliable benefit***
- ***Surgical hearing improvement is unpredictable***

Audiology Role

Audiology is responsible for:

- ***Early diagnosis***
- ***Objective hearing assessment***
- ***Selection and fitting of hearing devices***
- ***Long term monitoring***
- ***Counseling of family and patient***

Audiology does not decide surgical candidacy.

Medico-Legal Considerations

- ***Delay in amplification may cause language delay***
- ***Unilateral cases still require documentation and follow up***
- ***Counseling and rehabilitation plans must be documented***

Safe wording:

"Hearing rehabilitation was recommended to support auditory development in congenital aural atresia."

Key Points

- ***Congenital aural atresia causes conductive hearing loss***
- ***Inner ear is usually normal***
- ***Early rehabilitation is essential***
- ***Surgery is optional and anatomy dependent***

Golden Rule

In congenital aural atresia, hearing access matters more than canal appearance.

Audiology Management Flowchart

Congenital Aural Atresia

Newborn or child diagnosed with congenital aural atresia

↓

Step 1 Confirm laterality

• Unilateral

• Bilateral

↓

Step 2 Initial audiologic assessment

• ABR with bone conduction

• OAE if feasible

↓

***Step 3** Determine hearing status*

• Conductive hearing loss confirmed

• Bone conduction thresholds normal

↓

***Step 4** Early intervention decision*

↓ ***Bilateral***	↓ ***Unilateral***
↓	↓
Immediate hearing access	***Recommend hearing access***
• Bone conduction device	***• Bone conduction device***
• Soft band in infants	***• Classroom accommodations***

↓

***Step 5** Ongoing monitoring*

• Speech and language development

• Educational performance

↓

***Step 6** Imaging and surgical consideration*

• CT temporal bone when age appropriate

• ENT evaluates surgical candidacy

↓

***Step 7** Long term rehabilitation*

• Upgrade device as child grows

• Reassess hearing and outcomes

Flowchart Rule

Early hearing access is mandatory, even when surgery is planned later.

Audiology Note

Bone conduction devices provide more predictable hearing benefit than surgery.

Hospital Policy

Management of Congenital Ear Anomalies

ENT and Audiology Joint Policy

1. Purpose

To ensure early identification, hearing rehabilitation, and safe multidisciplinary management of patients with congenital ear anomalies, including aural atresia.

2. Scope

This policy applies to:

- **Audiology Department**
- **ENT Department**
- **Pediatrics and Neonatology**
- **Speech and Language Therapy**

3. Policy Statement

All patients with congenital ear anomalies must receive early audiologic assessment and hearing rehabilitation, regardless of later surgical plans.

4. Definitions

Congenital Ear Anomalies

Structural abnormalities of the external or middle ear present at birth that may affect hearing.

5. Audiology Responsibilities

Audiology must:

- ***Perform early objective hearing assessment***
- ***Confirm type and degree of hearing loss***
- ***Initiate hearing rehabilitation promptly***
- ***Monitor auditory and language development***
- ***Counsel family regarding realistic expectations***

6. ENT Responsibilities

ENT must:

- ***Assess anatomical severity***
- ***Request imaging when appropriate***
- ***Determine surgical candidacy and timing***
- ***Coordinate care with Audiology***

7. Hearing Rehabilitation Standards

- ***Bilateral atresia***

– Immediate bone conduction hearing device

– Urgent speech and language referral

- *Unilateral atresia*

– Hearing rehabilitation recommended

– Educational support advised

8. Surgical Considerations

- *Surgery is elective*
- *Decision is anatomy dependent*
- *Hearing rehabilitation must not be delayed for surgery*

9. Documentation Standards

Mandatory documentation includes:

- *Laterality and severity*
- *Audiologic test results*
- *Rehabilitation plan*
- *Counseling provided to family*

Mandatory wording:

"Hearing rehabilitation was initiated to support auditory and language development in congenital ear anomalies."

10. Medico Legal Considerations

- *Delay in hearing access increases risk of language delay*
- *Unilateral cases still require documentation and follow up*
- *Compliance with this policy is mandatory*

Policy Key Statement

In congenital ear anomalies, hearing access is a developmental emergency, not a cosmetic issue.

Common Mistakes

- *Treating atresia as simple conductive loss*
- *Over-relying on bone conduction results*

What I Would Explain to the Family

"The ear canal did not form normally.

We assess hearing carefully, but decisions require imaging and a team approach."

Congenital Canal Stenosis (Non-Atretic)

Congenital canal stenosis refers to:

- *A narrowed but present external auditory canal*
- *Without complete atresia*

It is a developmental condition, not a transient finding.

Clinical Characteristics

- *Narrow canal diameter*
- *Reduced self-cleaning*
- *Tendency for debris accumulation*
- *Increased infection risk*

Unlike atresia:

- *Access exists*
- *But it is functionally unstable*

Audiological Implications

- *Conductive component may be mild or fluctuating*
- *Tympanometry may be technically difficult*
- *Probe placement variability affects results*

Key danger:

- ***Progressive worsening mistaken for cochlear decline***

Stenosis behaves like partial atresia over time.

Otologic Perspective

Otologists worry about:

- ***Chronic inflammation***
- ***Canal skin breakdown***
- ***Surgical access limitations***

Audiological reports should emphasize:

- ***Stability vs progression***
- ***Reliability of access***
- ***Need for longitudinal monitoring***

Exostosis — Broad-Based Progressive Narrowing

Definition

Exostosis is:

- ***A benign bony overgrowth***
- ***Broad-based***
- ***Often bilateral***

Strongly associated with:

- ***Cold water exposure***
- ***Repetitive irritation***

Clinical Characteristics

- ***Slowly progressive canal narrowing***
- ***Moisture trapping***
- ***Recurrent otitis externa***

Often asymptomatic until advanced.

Audiological Implications

- *Fluctuating conductive loss*
- *Increasing test variability*
- *Progressive access compromise*

Audiograms may show:

- *Inconsistent air–bone gaps*
- *Sudden "improvement" after cleaning*
- *Apparent progression without cochlear cause*

Exostosis distorts data before it causes symptoms.

Osteoma — Focal Obstruction with Big Consequences

Definition

Osteoma is:

- *A solitary, pedunculated bony lesion*
- *Usually unilateral*
- *Slow-growing*

Clinical Characteristics

- *Partial obstruction*
- *Debris entrapment behind lesion*
- *Infection risk*

Often discovered incidentally.

Audiological Implications

- *Localized conductive effects*
- *Sudden threshold changes after minor inflammation*

- ***High-frequency distortion***

Osteoma is deceptive:

- ***Looks small***
- ***Acts big***

Why Progressive EAC Disease Confuses Audiology

Common patterns of misinterpretation:

- ***Fluctuation labeled as “test variability”***
- ***Gradual decline labeled as SNHL progression***
- ***Improvement after cleaning misread as recovery***

Otologists think longitudinally.

Audiology must do the same.

Management Principles — What Audiology Should Do

Audiology should:

- ***Document canal diameter and access quality***
- ***Note debris accumulation tendency***
- ***Avoid definitive conclusions from single sessions***
- ***Recommend ENT review when progression is suspected***

Audiology should not:

- ***Attribute progression to cochlear pathology prematurely***
- ***Over-interpret small changes***
- ***Ignore access limitations in reports***

What I Would Explain to the Family

"The ear canal is narrowing over time, which affects how sound enters the ear.

We monitor this carefully before making bigger decisions."

EAC Decision Algorithm

Clinical Flowchart for Safe Audiological Interpretation

Why This Algorithm Exists

Most audiological errors related to the outer ear occur because clinicians:

- ***Test first***
- ***Interpret quickly***
- ***Inspect late***

Otologists reverse this order.

This algorithm provides a step-by-step clinical pathway to ensure that:

- ***Data is valid***
- ***Interpretation is safe***
- ***Decisions are defensible***

In EAC pathology, sequence matters more than speed.

STEP 1 — Inspect the External Auditory Canal

Before any test, assess:

- ***Canal patency***
- ***Presence of cerumen***
- ***Inflammation or discharge***
- ***Pain or tenderness***
- ***Structural narrowing or bony lesions***

If canal is not fully visible or stable → STOP.

STEP 2 — Is the Canal Clear and Stable?

YES

Proceed to audiological assessment with confidence.

NO

Do not finalize interpretation.

Classify the limitation:

- ***Obstruction (cerumen, foreign body)***
- ***Inflammation (otitis externa)***
- ***Structural narrowing (stenosis, exostosis, osteoma)***
- ***Developmental absence (atresia)***

Document clearly and proceed to Step 3.

STEP 3 — Determine the Nature of the Limitation

Temporary / Reversible

Examples:

- ***Cerumen***
- ***Acute otitis externa***

Action:

- ***Treat or refer***
- ***Defer definitive testing***
- ***Reassess after resolution***

Structural / Progressive

Examples:

- ***Congenital canal stenosis***
- ***Exostosis***

- ***Osteoma***

Action:

- ***Interpret results with caution***
- ***Focus on trends, not absolutes***
- ***Recommend ENT review***
- ***Avoid labeling progression as cochlear***

Developmental / Absent

Examples:

- ***Aural atresia***

Action:

- ***Shift testing strategy (BC-focused)***
- ***Avoid surgical predictions***
- ***Require imaging + MDT discussion***

STEP 4 — Can Audiological Data Be Trusted?

Ask:

- ***Is access consistent between visits?***
- ***Are results repeatable?***
- ***Is there fluctuation suggesting instability?***

If NO:

- ***Qualify every conclusion***
- ***Avoid irreversible recommendations***

Unstable access produces unstable truth.

STEP 5 — Reporting Language (Critical Step)

Use protective phrasing:

Acceptable:

"Findings are interpreted with caution due to external auditory canal pathology."

Dangerous:

"Hearing loss is progressive."

(without addressing canal status)

STEP 6 — Decide the Next Action

Canal Status	*Audiology Action*
Clear & stable	*Proceed normally*
Obstructed	*Clear → retest*
Inflamed	*Treat → retest*
Narrowed	*Monitor trends*
Atretic	*MDT + imaging*

STEP 7 — Patient & Family Counseling

Explain simply:

"Before we trust hearing tests, we need to make sure sound can enter the ear reliably."

This prevents:

- ***False reassurance***
- ***Premature anxiety***
- ***Medico-legal exposure***

Medico-Legal Protection Tip

Always document canal status before interpreting results.

Teaching Pearl

The cochlea should never be blamed before the canal is cleared.

Take-Home Messages

- ***Inspect first***

EAC–Hearing Aid Interaction

When the Canal Shapes Amplification

Hearing aids do not work in isolation.

They work inside an ear canal.

When the external auditory canal is abnormal, unstable, or diseased, amplification may:

- ***Fail technically***
- ***Harm physiologically***
- ***Or worsen symptoms***

Otologists often see the consequences.

Audiologists often see the complaints.

This chapter explains when the canal—not the device—is the problem.

1. Narrow or Stenotic Canals

Clinical impact:

- ***Difficulty achieving acoustic seal***
- ***Increased feedback***
- ***Limited gain before instability***

Audiological implications:

- ***Over-amplification attempts***
- ***Distorted fitting outcomes***
- ***False assumption of "poor cochlear reserve"***

Some canals physically cannot tolerate amplification.

2. Exostosis & Osteoma

Clinical impact:

- ***Moisture trapping***
- ***Poor ventilation***
- ***Recurrent otitis externa***

Audiological implications:

- ***Recurrent device intolerance***
- ***Frequent repairs or refitting***
- ***Patient non-compliance***

Key principle:

A canal prone to infection is not a safe amplification environment.

3. Otitis Externa & Chronic Canal Inflammation

Clinical impact:

- ***Pain***
- ***Edema***
- ***Skin breakdown***

Audiological implications:

- ***Absolute contraindication to fitting***
- ***Need for medical management first***

Amplification in an inflamed canal is not treatment—it is provocation.

4. Aural Atresia & Severe Stenosis

Audiological implications:

- *Conventional air-conduction hearing aids are inappropriate*
- *Bone-conduction solutions are preferred*
- *MDT planning is mandatory*

Audiology should never:

- *Force air-conduction fitting*
- *Promise benefit*

Some external auditory canal findings are not "ENT problems later."

They are audiology stop signs now.

Missing them exposes:

- *Patients to harm*
- *Clinicians to medico-legal risk*

Absolute Red Flags

Immediate ENT referral required if any of the following are present:

- *Persistent otorrhea*
- *Bleeding from the canal*
- *Severe pain disproportionate to findings*

Final EAC Message (Before Moving On)

Clear access protects clear interpretation.

Tympanic Membrane & Middle Ear Pathologies

Where Transmission Becomes Interpretation

Why This Part Matters

The tympanic membrane and middle ear are often treated as:

- Simple mechanical systems
- Causes of “just conductive” hearing loss

In otologic decision-making, this is incomplete.

Otologists view the tympanic membrane and middle ear as:

- Indicators of chronicity
- Predictors of surgical complexity
- Modifiers of long-term outcomes

Audiologists must therefore approach middle ear pathology not as isolated findings, but as dynamic systems that evolve over time.

The membrane does not just vibrate — it tells a story.

How Otologists Think About the Tympanic Membrane

Otologists do not ask:

- Is there a perforation?
- Is the tympanogram flat?

They ask:

- *How long has this been present?*
- *Is it stable or progressive?*
- *What does it imply about middle ear health?*

Audiological interpretation must follow this logic.

What This Part Will Cover

PART III will address:

- *Tympanic membrane integrity and pathology*
- *Middle ear mechanics and ventilation*
- *Acute vs chronic disease*
- *Audiological consequences*
- *Decision points for referral and intervention*

Each chapter will emphasize:

- *Stability vs instability*
- *Reversibility vs chronic change*
- *Interpretation risk*

How to Read This Part

Before interpreting thresholds, always ask:

- *Is the membrane intact?*
- *Is the middle ear aerated?*
- *Is pathology active or quiescent?*

Middle ear disease is rarely a snapshot — it is a timeline.

Chapter



12

Tympanic Membrane

The Most Interpreted — and Most Misunderstood Structure

Why This Chapter Matters

The tympanic membrane (TM) is often reduced to:

- “Intact” or “perforated”
- “Normal” or “abnormal”

For otologists, this is insufficient.

The TM is:

- A biomechanical structure
- A historical record of middle ear health
- A predictor of stability and surgical outcome

Audiological interpretation becomes unsafe
when the TM is treated as a checkbox instead of a system.

The membrane does not simply transmit sound — it reflects disease behavior.

1. Anatomical Structure of the Tympanic Membrane

The TM consists of:

- Pars tensa (major functional component)
- Pars flaccida (pressure-sensitive region)

Layered structure:

1. ***Outer epithelial layer***
2. ***Middle fibrous layer***
3. ***Inner mucosal layer***

Clinical relevance:

- ***The fibrous layer determines mechanical integrity***
- ***Loss or thinning alters sound transmission***
- ***Structural damage predicts chronicity***

2. Orientation and Mechanical Function

The TM is:

- ***Obliquely oriented***
- ***Cone-shaped***
- ***Attached to the manubrium of the malleus***

Physiological implications:

- ***Converts acoustic energy into mechanical motion***
- ***Distributes energy across ossicles***
- ***Maintains impedance matching***

Small structural changes produce disproportionate functional effects.

3. Tympanic Membrane Compliance and Mobility

Normal TM function depends on:

- ***Elasticity***
- ***Tension***
- ***Pressure balance***

Factors affecting mobility:

- ***Middle ear pressure***
- ***Scarring***
- ***Fluid***
- ***Ossicular coupling***

Audiological relevance:

- ***Reduced mobility ≠ fixed loss***
- ***Hypermobility ≠ normal function***

4. Vascularity and Healing Capacity

The TM has:

- ***Limited but adequate blood supply***
- ***Capacity for spontaneous healing***

Clinical implications:

- ***Acute changes may resolve***
- ***Recurrent insults reduce healing potential***
- ***Chronic disease alters tissue quality***

Otologists assess tissue quality, not just closure.

5. Tympanic Membrane as a Disease Timeline

The TM records:

- ***Past infections***
- ***Ventilation problems***
- ***Pressure dysregulation***

Signs of chronicity:

- ***Tympanosclerosis***
- ***Retraction***

- *Atrophy*

A "closed" membrane may still be a diseased membrane.

6. Audiological Implications of TM Physiology

TM pathology can:

- ***Alter impedance***
- ***Distort tympanometry***
- ***Produce inconsistent thresholds***

Audiologists must:

- ***Correlate visual findings with test behavior***
- ***Avoid absolute conclusions from curves alone***
- ***Document observed TM features***

7. Why Otologists Respect the TM So Much

Otologists see the TM as:

- ***A window into middle ear health***
- ***A predictor of surgical success***
- ***A determinant of recurrence risk***

Audiology strengthens MDT decisions
when it communicates TM condition accurately and descriptively.

The Sentence That Changes Everything

"The tympanic membrane appears intact but shows features suggestive of chronic middle ear stress."



Chapter

13

Trauma of the Middle Ear &Tympanic Membrane Perforations

More Than a Hole in the Membrane

Why This Chapter Matters

A tympanic membrane perforation is often reduced to:

- ***Size***
- ***Location***
- ***Presence or absence***

In otologic thinking, this is dangerously simplistic.

Otologists assess perforations as:

- ***Evidence of disease behavior***
- ***Indicators of middle ear environment***
- ***Predictors of healing and surgical success***

Audiological interpretation becomes unsafe

when perforations are treated as static findings.

A perforation is not a defect — it is a diagnosis in context.

1. Definition and Basic Classification

A tympanic membrane perforation is a:

- ***Disruption of membrane integrity***

- ***Resulting in altered sound transmission and ventilation***

Basic descriptive elements:

- ***Size (small / medium / large)***
- ***Location (central, marginal)***
- ***Chronicity (acute vs chronic)***

However:

Description alone does not define impact.

2. Acute vs Chronic Perforations

Acute Perforations

Common causes:

- ***Infection***
- ***Trauma***
- ***Barotrauma***

Characteristics:

- ***Active inflammation may be present***
- ***Healing potential is high***
- ***Audiological findings may fluctuate***

Audiological implication:

- ***Avoid permanent labeling***
- ***Monitor resolution before final conclusions***

Chronic Perforations

Characteristics:

- ***Long-standing***
- ***Altered tissue quality***

- *Reduced healing capacity*

Otologic implications:

- *Higher risk of recurrent infection*
- *Surgical planning considerations*

Audiological implication:

- *Stability matters more than magnitude*

Chronicity changes everything.



▲ *Figure 13–1. Types of central perforation (A) Small; (B) Medium; (C) Large; (D) Sub-total*



▲ *Figure 13–2. Types of perforations of tympanic membrane in cases of chronic suppurative otitis media (CSOM).*
(A) Small central perforation
in anterosuperior quadrant;
(B) Medium size kidney-shaped central perforation;
(C) Subtotal central perforation;
(D) Total perforation with destruction of fibrous annulus;
(E) Attic perforation of pars flaccida; and
(F) Posterosuperior marginal perforation.
Note: A, B and C occur in tubotympanic CSOM. D, E and F seen in CSOM with cholesteatoma


Incudostapedial joint
Handle of malleus
Eustachian tube orifice
Round window niche
Pro-montory

▲ *Figure 13–3. Showing sub-total central perforation*



▲ *Figure 13–4. (A) Tympanic membrane (TM) perforation with ossicular discontinuity. (B) Right-sided conductive hearing loss of TM perforation with ossicular discontinuity.*



▲ Figure 13–5. Tympanic membrane perforation with chronic inflammation.

3. Location Matters — Not All Perforations Behave the Same

Central Perforations

- *Surrounded by TM remnant*
- *Often safer*
- *Lower risk of cholesteatoma*

Audiological pattern:

- *Conductive loss proportional to size*
- *Relatively predictable*

Marginal Perforations

- *Involve annulus*
- *Higher risk of epithelial migration*

Otologic concern:

- *Cholesteatoma risk*
- *Progressive disease*

Audiological implication:

- ***Treat as high-risk***
- ***Do not downplay***

4. Size vs Function — A Common Misconception

Large perforations may:

- ***Produce moderate loss***
- ***But remain stable***

Small perforations may:

- ***Cause disproportionate symptoms***
- ***Be associated with chronic disease***

Function does not scale linearly with size.

5. Middle Ear Environment and Ventilation

Perforations alter:

- ***Middle ear aeration***
- ***Pressure regulation***
- ***Infection susceptibility***

Audiological consequences:

- ***Fluctuating thresholds***
- ***Variable speech understanding***
- ***Inconsistent tympanometric patterns***

Otologists assess:

- ***Dry vs wet ear***
- ***Stability over time***

6. Audiological Assessment Considerations

Key principles:

- ***Tympanometry may be misleading***
- ***Large ear canal volume is expected***
- ***Masking assumptions may fail***

Audiology should:

- ***Document perforation characteristics***
- ***Correlate with behavioral responses***
- ***Avoid over-interpretation***

7. Why Otologists Care About Perforations

Perforations inform:

- ***Disease chronicity***
- ***Surgical timing***
- ***Risk of recurrence***

Audiology supports decisions by:

- ***Describing stability***
- ***Tracking functional impact***
- ***Communicating uncertainty clearly***

The audiogram informs function — the perforation explains behavior.

Clinical Summary

Tympanic membrane perforations reflect underlying disease behavior and middle ear environment, not just mechanical defects.

Middle ear trauma may result from direct injury, pressure changes, or severe head trauma.

The consequences range from simple conductive hearing loss to life-threatening complications.

1. Traumatic Perforation of the Tympanic Membrane

Definition

A traumatic perforation is a tear in the tympanic membrane caused by direct or indirect trauma.

Causes

- ***Slap over the ear***
- ***Penetrating injury***
- ***Unskilled ear instrumentation***
- ***Ear syringing***
- ***Sudden pressure changes***
- ***Temporal bone trauma***

Clinical Picture

Symptoms

- ***Sudden severe ear pain***
- ***Hearing loss***
- ***Tinnitus***
- ***Bleeding from the ear***
- ***Vertigo may occur***

Signs

- ***Fresh perforation of the tympanic membrane***
- ***Blood clots in the external canal***
- ***Middle ear mucosa may be visible***

Audiologic Findings

- ***Conductive hearing loss***
- ***Mixed loss if inner ear is involved***

Treatment

- ***Keep ear dry***
- ***Avoid ear drops unless infected***
- ***Most perforations heal spontaneously***
- ***Surgical repair if perforation persists beyond 3 months***

Clinical Note

Sudden SNHL or vertigo suggests inner ear injury and needs urgent ENT referral.

2. Otitic Barotrauma

Definition

Otitic barotrauma is injury to the middle ear caused by failure of pressure equalization across the tympanic membrane.

Causes

- ***Air travel***
- ***Diving***
- ***Hyperbaric exposure***
- ***Rapid altitude changes***

Predisposing factors

- ***Upper respiratory infection***
- ***Eustachian tube dysfunction***

Pathology

- *Negative middle ear pressure*
- *Congestion and hemorrhage of middle ear mucosa*
- *TM retraction or rupture*

Clinical Picture

Symptoms

- *Ear pain during ascent or descent*
- *Ear fullness*
- *Hearing loss*
- *Tinnitus*
- *Vertigo in severe cases*

Signs

- *Congested or retracted TM*
- *Hemotympanum*
- *TM perforation in severe cases*

Audiologic Findings

- *Conductive hearing loss*
- *Tympanometry shows negative pressure or flat curve*

Treatment

- *Decongestants*
- *Analgesics*
- *Avoid pressure exposure until recovery*
- *Myringotomy in severe cases*

Golden Rule

Never dive or fly with an upper respiratory infection.



▲ Figure 13–6. Increased pressure causing tympanic perforation and inner ear barotrauma resulting in round window perforation and perilymph leak.

3. Temporal Bone Fracture

Definition

A temporal bone fracture occurs as part of severe head injury and commonly affects the middle and inner ear.

Types

Longitudinal fracture

- More common
- Involves external canal and middle ear
- Causes conductive hearing loss

Transverse fracture

- Less common
- Involves inner ear
- Causes sensorineural hearing loss and vertigo

Clinical Picture

Symptoms

- ***Ear bleeding***
- ***Hearing loss***
- ***Vertigo***
- ***Facial nerve paralysis***
- ***CSF otorrhea***

Signs

- ***Hemotympanum***
- ***TM perforation***
- ***Battle's sign***

Audiologic Findings

- ***Conductive loss in longitudinal fractures***
- ***Sensorineural loss in transverse fractures***

Management

- ***Hospital admission***
- ***Multidisciplinary management***
- ***Audiologic testing after stabilization***

Red Flag

CSF otorrhea indicates skull base fracture.


Transverse fracture line
Long axis of temporal bone
Longitudinal fracture line

▲ *Figure 13–7. Temporal bone fracture*


Longitudinal fracture
Foramen ovale
Geniculate ganglion
External auditory canal
Clivus
A
Jugular foramen
Semicircular canals
Transverse fracture
Foramen spinosum
Foramen lacerum
Clivus
Foramen magnum
B
External auditory canal

▲ *Figure 13–8. Orientations of temporal bone fractures. (A) Longitudinal fractures begin at the squamous portion of the temporal bone, run through the external auditory canal, and then turn anteriorly toward the foramen lacerum. (B) Transverse fractures begin from the foramen magnum, run petrous pyramid, and then turn anteriorly toward the foramen lacerum.*

4. Ossicular Dislocation

Definition

Ossicular dislocation is disruption of the ossicular chain following trauma.

Causes

- ***Head injury***
- ***Barotrauma***
- ***Penetrating ear trauma***

Clinical Picture

Symptoms

- ***Persistent hearing loss after trauma***
- ***Minimal pain***

Signs

- ***Normal or healed TM***
- ***No active infection***

Audiologic Findings

- ***Marked conductive hearing loss***
- ***Large air bone gap***
- ***Speech discrimination usually preserved***

Diagnosis

- ***Pure tone audiometry***
- ***CT scan of temporal bone***

Treatment

- ***Observation initially***
- ***Ossiculoplasty if hearing loss persists***

Clinical Insight

A large air bone gap with an intact TM after trauma strongly suggests ossicular discontinuity.

Golden Rule

In ear trauma, always correlate hearing findings with mechanism of injury before testing aggressively.

Middle ear trauma may result from direct injury, pressure changes, or severe head trauma.

The consequences range from simple conductive hearing loss to life-threatening complications.

Mandatory Otoscopic Examination

Before audiologic testing, the audiologist must document:

☐ Condition of the tympanic membrane

☐ Presence of perforation

☐ Presence of blood or clots

☐ Hemotympanum

☐ Suspicion of CSF leak

Audiologic testing without prior otoscopy is not permitted in traumatic cases.

Mandatory wording

"Audiologic testing was limited due to recent middle ear trauma. Findings should be interpreted with caution and correlated clinically."

Red Flags Requiring Immediate ENT Referral

☐ ***Sudden sensorineural hearing loss***

☐ ***Persistent vertigo***

☐ ***Facial nerve weakness***

☐ ***CSF like ear discharge***

☐ ***Severe headache***

☐ ***Progressive hearing loss after trauma***

ENT Referral Pathway

- ***Routine referral for stable traumatic perforation without red flags***
- ***Urgent referral for any red flag finding***
- ***Emergency referral for suspected skull base fracture or CSF leak***

Referral documentation must include audiologic findings and safety concerns.

Flowchart

ENT–Audiology Joint Management of Middle Ear Trauma

Patient with Suspected Middle Ear Trauma

(slapping injury, barotrauma, head injury, ear bleeding)

↓

Audiology Safety Screening

↓

Is the patient clinically stable?

(No vertigo, no facial weakness, no CSF leak, no severe head injury)

↓ YES

Proceed with limited audiologic assessment

↓ NO

STOP audiologic testing
Urgent ENT / Emergency referral

↓

Mandatory Otoscopic Examination

↓

Findings documented?
(TM status, blood, hemotympanum)

↓

Select Audiologic Tests Safely

- Pure tone audiometry
- Speech audiometry
- Tympanometry ONLY if TM intact

↓

Any Red Flags Identified?
(Sudden SNHL, vertigo, facial palsy, CSF otorrhea, severe headache)

↓ YES

Urgent ENT referral

↓ NO

Routine ENT referral

↓

ENT Evaluation

• Otologic diagnosis

• Imaging if indicated

• Medical / surgical management

↓

ENT Clearance for Reassessment

↓

Audiology Follow Up

(Post healing or post surgery)

Immediate ENT or Emergency referral if any of the following are present:

- ***Sudden sensorineural hearing loss***
- ***Persistent or severe vertigo***
- ***Facial nerve weakness***
- ***Clear watery ear discharge (suspected CSF)***
- ***Severe headache or altered consciousness***

14

Retraction Pockets & Atelectasis

When the Membrane Signals Progressive Middle Ear Disease

Why This Chapter Matters

Retraction of the tympanic membrane is often described as:

- ***"Mild"***
- ***"Shallow"***
- ***"Non-significant"***

In otologic thinking, this is misleading.

Retraction pockets and atelectasis represent:

- ***Chronic ventilation failure***
- ***Progressive middle ear pathology***
- ***Potential precursors to cholesteatoma***

Audiologists must recognize these conditions as dynamic risk states, not static findings.

Retraction is not a position — it is a process.

1. Pathophysiological Basis

Retraction occurs due to:

- ***Chronic negative middle ear pressure***
- ***Eustachian tube dysfunction***

- *Loss of TM structural integrity*

Over time, this leads to:

- ***Progressive membrane collapse***
- ***Adhesion to middle ear structures***
- ***Loss of self-cleaning capacity***

Atelectasis represents the end-stage of this process.

2. Retraction Pockets — Localized Disease

Characteristics

- ***Localized inward displacement***
- ***Commonly affects:***
    - ***Pars flaccida***
    - ***Posterior pars tensa***

Clinical relevance:

- ***Keratin accumulation risk***
- ***Limited visualization of pocket depth***

Audiological implications:

- ***Mild or fluctuating conductive loss***
- ***Often underestimated functional impact***

A small pocket can hide a big problem.

3. Atelectasis — Global Membrane Collapse

Characteristics

- ***Generalized TM collapse***
- ***Adhesion to ossicles or promontory***

Clinical relevance:

- ***High-risk middle ear environment***
- ***Poor ventilation and clearance***

Audiological implications:

- ***Variable conductive loss***
- ***Poor test repeatability***
- ***Tympanometry often misleading***

4. Stability vs Progression — The Critical Question

Otologists ask:

- ***Is the retraction stable?***
- ***Is it deepening?***
- ***Is debris accumulating?***

Audiology must support this by:

- ***Documenting changes over time***
- ***Avoiding single-session conclusions***
- ***Communicating uncertainty***

Progression matters more than appearance.

5. Audiological Assessment Considerations

Key points:

- ***Tympanometry may appear “near-normal”***
- ***Conductive loss may be mild***
- ***Speech performance may fluctuate***

Audiology should:

- ***Correlate otoscopy with results***
- ***Flag retraction as a risk condition***

- ***Recommend ENT follow-up***

6. Why Otologists Take Retraction Seriously

Retraction and atelectasis predict:

- ***Chronic disease behavior***
- ***Surgical complexity***
- ***Recurrence risk***



▲ ***Figure 14–1. Middle ear atelectasis with effusion.***



▲ ***Figure 14–2. Adhesive otitis media with ossicular erosion.***

When NOT to Perform Tympanometry

Do NOT perform tympanometry when:

- ***Active otitis externa is present***
- ***Severe canal pain prevents proper seal***
- ***External auditory canal is obstructed***
- ***Large TM perforation is already visible***
- ***Recent ear surgery (unless cleared)***
- ***Profuse otorrhea is present***
- ***Patient cannot tolerate probe pressure***

Why Tympanometry Becomes Dangerous

Tympanometry assumes:

- ***Intact membrane or controlled perforation***
- ***Stable canal conditions***
- ***Predictable pressure transmission***

When these conditions fail:

- ***Results become misleading***
- ***False reassurance occurs***
- ***Clinical decisions may be unsafe***

A curve is not evidence if the conditions are invalid.

Medico-Legal Protection Sentence

"Tympanometry was deferred due to conditions limiting validity and patient safety."

Teaching Pearl

Knowing when not to test is as important as knowing how to test.

Final Safety Message

No tympanogram is better than a misleading one.

Atelectatic Otitis Media

Atelectatic otitis media is a condition characterized by marked retraction of the tympanic membrane onto the promontory and ossicles, with partial or complete obliteration of the middle ear space.

The tympanic membrane is thinned and weakened but not adherent to the medial wall.

Etiology

- ***Long standing Eustachian tube dysfunction***
- ***Chronic otitis media with effusion***
- ***Recurrent attacks of acute otitis media***
- ***Progressive thinning and loss of elasticity of the tympanic membrane***

Pathophysiology

- ***Persistent negative middle ear pressure leads to progressive retraction of the tympanic membrane***
- ***The middle ear air space becomes reduced***
- ***The ossicular chain may be contacted by the retracted tympanic membrane***
- ***Middle ear ventilation becomes progressively impaired***
- ***The condition is potentially reversible if treated early***

Clinical Picture

Symptoms

- *Deafness, usually conductive*
- *Tinnitus*
- *Ear fullness or blockage sensation*

Signs

- *Tympanic membrane is markedly retracted*
- *Retraction may be seen over*

– *Promontory*

– *Ossicles*

- *Tympanic membrane is thin and transparent*
- *Tympanic membrane is not adherent to the medial wall*
- *Reduced or absent mobility on pneumatic otoscopy*
- *Tuning fork tests show conductive hearing loss*

Audiologic Findings

- *Pure tone audiometry*

– *Conductive hearing loss*

– *Variable air bone gap*

- *Tympanometry*

– *Type C tympanogram indicating Eustachian tube dysfunction*

- *Speech audiometry*

– *Usually normal speech discrimination*

Differential Diagnosis

Condition	*Key Difference*
Otitis media with effusion	*Presence of middle ear fluid*

Condition	Key Difference
Adhesive otitis media	TM is adherent to medial wall
Cholesteatoma	Presence of keratin debris and retraction pocket

Treatment

Preventive

- Early and adequate treatment of
  - Acute otitis media
  - Otitis media with effusion
- Management of nasal and nasopharyngeal pathology

Conservative

- Repeated Eustachian tube inflation exercises
- Treatment of nasal allergy and sinus disease

Surgical

- Insertion of ventilation tubes, usually T tubes
- Restores middle ear aeration
- Prevents progression to adhesive otitis media
- Condition is usually reversible if managed early

Adhesive Otitis Media

Definition

Adhesive otitis media is a condition in which the tympanic membrane becomes permanently adherent to the medial wall of the middle ear due to

long-standing Eustachian tube dysfunction and persistent negative middle ear pressure.

Unlike atelectatic otitis media, the condition is irreversible.

Etiology

- ***Long standing Eustachian tube dysfunction***
- ***Untreated or recurrent otitis media with effusion***
- ***Progression from atelectatic otitis media***
- ***Recurrent acute otitis media***
- ***Chronic nasal or nasopharyngeal pathology***

Pathophysiology

- ***Persistent negative middle ear pressure causes progressive retraction of the tympanic membrane***
- ***The tympanic membrane loses elasticity and thickness***
- ***Eventually, the membrane becomes adherent to the promontory and ossicles***
- ***Middle ear air space is completely obliterated***
- ***Ossicular movement becomes restricted***

Clinical Picture

Symptoms

- ***Conductive hearing loss, often progressive***
- ***Ear fullness***
- ***Tinnitus***
- ***Usually no pain or discharge***

Signs

- ***Tympanic membrane is severely retracted***
- ***Tympanic membrane is plastered to the medial wall***
- ***Ossicles and promontory are clearly visible***
- ***No middle ear space***
- ***Tympanic membrane does not move on pneumatic otoscopy***

Audiologic Findings

- ***Pure tone audiometry***

– Conductive hearing loss

– Variable but often significant air bone gap

- ***Tympanometry***

– Flat curve, Type B

– No compliance

- ***Speech audiometry***

– Usually normal speech discrimination

Differential Diagnosis

Condition	*Key Feature*
Otitis media with effusion	*Presence of middle ear fluid*
Atelectatic otitis media	*TM retracted but not adherent*
Cholesteatoma	*Retraction pocket with keratin debris*

Treatment

Conservative Treatment

- ***Usually ineffective due to irreversible changes***

Surgical Treatment

- ***Tympanoplasty***
- ***Ossiculoplasty if ossicular fixation is present***
- ***Ventilation tubes are generally ineffective***

Prognosis

- ***Hearing improvement is often limited***
- ***Risk of recurrence exists***
- ***Early treatment of earlier stages improves outcome***

Golden Rule

Treat otitis media with effusion and atelectatic otitis media early to prevent adhesive otitis media.



Chapter

15

Tympanometry Revisited — Beyond Type A / B / C

From Curves to Clinical Meaning

Why This Chapter Matters

Tympanometry is one of the most commonly performed tests in audiology.

It is also one of the most misunderstood.

Too often, tympanograms are reduced to:

- ***Type A → Normal***
- ***Type B → Fluid***
- ***Type C → Eustachian tube dysfunction***

Otologists do not think this way.

They use tympanometry to ask deeper questions:

- ***Is the system stable or unstable?***
- ***Is this an acute or chronic process?***
- ***Does the curve match the otoscopic story?***

This chapter reframes tympanometry as a context-dependent tool, not a diagnostic label.

A tympanogram is not a diagnosis. It is a clue.

1. What Tympanometry Actually Measures

Tympanometry measures:

- ***Acoustic admittance of the middle ear system***
- ***As pressure is varied in the ear canal***

It reflects:

- ***Tympanic membrane mobility***
- ***Ossicular coupling***
- ***Middle ear pressure***

It does not directly measure:

- ***Presence of fluid***
- ***Disease severity***
- ***Hearing level***

Tympanometry measures mechanics, not pathology.

2. Type A — The Most Dangerous Curve

Why?

Because it is often interpreted as "normal".

A Type A tympanogram only means:

- ***The system moves at atmospheric pressure***

It does not exclude:

- ***Ossicular discontinuity***
- ***Early otosclerosis***
- ***Retraction pockets***
- ***Atelectasis with residual mobility***

Normal mobility does not guarantee a healthy middle ear.

3. Type B — Flat Does Not Mean Simple

A Type B tympanogram indicates:

- ***Severely reduced mobility***

Possible causes include:

- ***Middle ear effusion***
- ***Thickened TM***
- ***Tympanosclerosis***
- ***Large perforation (with large ECV)***

Audiological danger:

- ***Assuming fluid without otoscopy***
- ***Missing chronic disease***

A flat curve needs eyes, not assumptions.

4. Type C — Pressure Without Context Is Meaningless

Type C reflects:

- ***Negative middle ear pressure***

It does not automatically equal:

- ***Active Eustachian tube dysfunction***
- ***Pathology requiring treatment***

Type C may be:

- ***Transient***
- ***Compensated***
- ***Clinically insignificant***

Otologists ask:

- ***Is it persistent?***
- ***Is it symptomatic?***
- ***Is it progressing?***

Pressure alone does not define disease.

5. Tympanometric Width, Peak Height & Shape

Beyond A/B/C, otologists look at:

- *Peak height (compliance)*
- *Tympanometric width*
- *Curve symmetry*

Clinical implications:

- *Narrow, sharp peak → stiff but responsive system*
- *Broad peak → instability*
- *Hypermobility → atrophy or ossicular issues*

The shape of the curve often matters more than the letter.

6. Ear Canal Volume — The Forgotten Variable

ECV helps differentiate:

- *True Type B with effusion*
- *Type B with perforation*
- *Canal obstruction or probe error*

Audiology mistakes happen when:

- *ECV is ignored*
- *Results are interpreted blindly*

Always read the volume before reading the curve.

7. Correlation Is Mandatory

Tympanometry must be correlated with:

- *Otoscopy*
- *Audiogram*

- *Symptoms*
- *History*

If findings do not align:

Do not force interpretation

Otologists respect mismatch.

Audiology must document it.

8. Tympanometry as a Monitoring Tool

Its strongest value is in:

- *Tracking change over time*
- *Assessing stability*
- *Supporting MDT decisions*

One curve is information.

A series of curves is insight.

Trends reveal truth.

Clinical Summary

Tympanometry assesses middle ear mechanics, not disease labels, and must be interpreted within full clinical context.

The Golden Rule

Never interpret a tympanogram without otoscopy and audiometric correlation.

Common Mistakes

- *Ignoring ECV*

Red Flags

- *Mismatch between curve and otoscopy*
- *Persistent abnormal curves*

Pitfalls

- *False reassurance from Type A*



▲ *Figure 15–1. Various types of tympanograms*

The Sentence That Changes Everything

"Tympanometric findings are interpreted in correlation with otoscopic and audiometric data."

Clinical Decision-Making

Use tympanometry to support—not replace—clinical judgment.

The Uncomfortable Truth

Most tympanometric errors are interpretive, not technical.

Take-Home Messages

- ***Look beyond A/B/C***
- ***Correlate always***



Chapter

16

Otitis Media with Effusion (Foundations)

Fluid Without Fire — Disease Without Infection

Why This Chapter Matters

Otitis Media with Effusion (OME) is often misunderstood as:

- “Mild otitis”
- “Silent infection”
- “Temporary fluid that will surely resolve”

Otologists know OME as:

- A ventilation disorder, not an infection
- A time-dependent condition
- A major modifier of developmental outcomes, especially in children

Audiological misinterpretation of OME can lead to:

- Delayed language development
- Inappropriate reassurance
- Premature intervention

OME is quiet—but its consequences are not.

1. Definition and Core Concept

OME is defined as:

- Presence of non-purulent fluid in the middle ear

- *Without signs of acute infection*

Key distinction:

- ***No fever***
- ***No acute pain***
- ***No inflammatory “fire”***

OME reflects ventilation failure, not bacterial disease.

2. Pathophysiology — Why Fluid Accumulates

OME develops due to:

- ***Eustachian tube dysfunction***
- ***Negative middle ear pressure***
- ***Transudation of fluid***

Contributing factors:

- ***Upper respiratory infections***
- ***Adenoid hypertrophy***
- ***Allergic inflammation***
- ***Immature ET function in children***

Fluid is the result—not the cause—of dysfunction.

3. Acute vs Chronic OME

Acute / Transient OME

- ***Often follows URI***
- ***May resolve spontaneously***
- ***Short duration***

Audiological implication:

- ***Temporary conductive loss***

- ***Monitor, don't label***

Chronic OME

- ***Persists ≥ 3 months***
- ***Indicates ongoing ventilation failure***

Clinical implications:

- ***Structural TM changes***
- ***Hearing fluctuation***
- ***Developmental risk***

Duration transforms OME from nuisance to disease.

4. Audiological Characteristics

Typical findings:

- ***Conductive hearing loss (often mild–moderate)***
- ***Flat or reduced-mobility tympanogram***
- ***Fluctuating thresholds***
- ***Reduced speech audibility in noise***

Important:

- ***Severity of loss does not equal impact***
- ***Fluctuation disrupts learning more than stable loss***

5. Tympanic Membrane Clues in OME

Otoscopy may reveal:

- ***Dull or opaque TM***
- ***Air–fluid levels***
- ***Reduced mobility***
- ***Mild retraction***

Audiologists should:

- ***Describe findings***
- ***Avoid "normal TM" labeling***
- ***Correlate with tympanometry***

The membrane often whispers the diagnosis.



▲ ***Figure 16–1. Otoscopic examination of otitis media with effusion.***

6. OME in Children — Why Timing Matters

Children with OME may show:

- ***Speech delay***
- ***Poor attention***
- ***Behavioral concerns***
- ***Academic difficulty***

Risk increases when:

- ***OME is bilateral***

- ***OME is chronic***
- ***Child has additional vulnerabilities***

Children do not “adapt” to fluctuating access.

7. Management Principles — Audiology’s Role

Audiology should:

- ***Document degree and stability of loss***
- ***Track changes over time***
- ***Highlight developmental risk***
- ***Avoid overconfidence in single tests***

Audiology should not:

- ***Assume infection***
- ***Promise spontaneous resolution***
- ***Push irreversible decisions without context***

Otologic decisions depend on trajectory, not snapshots.

Clinical Summary

OME is a ventilation disorder causing fluctuating conductive hearing loss, with significant developmental implications when persistent.

Chronic Secretory Otitis Media

(Otitis Media with Effusion, Glue Ear)

Definition

Chronic secretory otitis media is the presence of serous, mucoid, or seromucinous fluid in the middle ear cleft behind an intact tympanic membrane, without signs of acute infection.

It is the most common cause of hearing loss in school-age children.

Types

1. Serous Otitis Media

- ***Caused by transudation of plasma from blood vessels due to negative middle ear pressure.***

2. Mucoid Otitis Media

- ***Caused by active mucus secretion from glands and cysts in the middle ear mucosa following long-standing irritation.***

Etiology

- ***Incomplete resolution of acute otitis media, especially with inadequate antibiotic treatment***
- ***Eustachian tube dysfunction due to***

– Adenoid hypertrophy

– Nasopharyngeal tumors, e.g. angiofibroma

– Cleft palate with abnormal action of levator palati

– Chronic rhinitis, sinusitis, nasal allergy

- ***Allergy of middle ear mucosa***
- ***Viral infections***
- ***Otitic barotrauma***
- ***Radiation therapy to head and neck causing fibrosis of the Eustachian tube***



▲ Figure 16–2. Secretory otitis media

Pathology

- *Prolonged Eustachian tube obstruction leads to persistent negative pressure in the middle ear*
- *Metaplasia of middle ear mucosa occurs*
- *Simple columnar epithelium changes into pseudostratified columnar epithelium rich in goblet cells*
- *Excess mucus secretion accumulates in the middle ear causing persistent effusion*

Clinical Picture

Symptoms

- *Deafness*

– Main and sometimes the only symptom

– In children presents as poor school performance or inattention

– Adults complain of ear fullness or blockage

- *Occasional bubbling sensation in the ear, mainly in adults*
- *Tinnitus*

Signs

- *Tympanic membrane is intact and usually retracted*
- *Otoscopic findings include*

– Foreshortened handle of malleus

– Prominent lateral process of malleus

– Exaggerated malleolar folds

– Disturbed or absent cone of light

– Reduced or absent mobility on pneumatic otoscopy

- *Tympanic membrane color varies*

– Amber yellow in serous effusion

– Dull gray in mucoid effusion

- *Fluid level or air bubbles may be visible behind the tympanic membrane*
- *Tuning fork tests show conductive hearing loss*

Investigations

- *Pure tone audiometry*

– Mild to moderate conductive hearing loss with air bone gap

- *Tympanometry*

– Flat curve, Type B

- *Lateral X-ray nasopharynx or nasopharyngoscopy*

– To detect adenoids or nasopharyngeal masses

Important clinical note

Any adult with unilateral secretory otitis media must be evaluated carefully to exclude nasopharyngeal carcinoma.

Treatment

Conservative Management

- ***Treatment of predisposing factors***
  - ***Control nasal and sinus infections***
  - ***Adenoidectomy when indicated***
- ***Systemic antibiotics to prevent secondary infection***
- ***Corticosteroids***
- ***Nasal decongestants to improve Eustachian tube function***
- ***Mucolytics***
- ***Eustachian tube ventilation exercises, e.g. Valsalva maneuver***

Surgical Management

- ***Myringotomy with insertion of tympanostomy ventilation tube***
  - ***Normalizes middle ear pressure***
  - ***Allows drainage of effusion***
  - ***Produces immediate hearing improvement***
  - ***Reduces recurrence***

Indications for surgery

- ***Middle ear effusion lasting more than 3 months despite medical treatment***
- ***Significant conductive hearing loss***
- ***Presence of nasopharyngeal tumor or before radiotherapy***
- ***Myringotomy incision is usually done in the anteroinferior quadrant of the tympanic membrane***

- ***Ventilation tubes***

– Short term tubes, extrude spontaneously after 3 to 6 months

– Long term tubes, T-tubes, preferred in adults

Myringotomy and grommet insertion



▲ *Figure 16–3. Myringotomy with grommet insertion*


PF
PT
Anterior
Posterior
A
B
Inferior

▲ Figure 16–4. Incisions for myringotomy: (A) Radial incision; (B) Circumferential Incision



▲ Figure 16–5. Shepard type of grommet



▲ Figure 16–6. Goode T–tube (Ventilation tube)



▲ *Figure 16–7. Grommet in situ*

Key Points

- ***OME is not infection***
- ***Chronicity defines risk***

The Golden Rule

In OME, duration matters more than degree.

The Uncomfortable Truth

Many language delays begin with "just fluid."

Chapter

17

Acute Otitis Media — Why Audiology Should Be Careful

When Active Disease Makes Testing Unsafe

Why This Chapter Matters

Acute Otitis Media (AOM) is not an audiological diagnosis.

Yet audiologists frequently encounter patients during or shortly after an acute episode.

The danger is not missing AOM.

The danger is testing through it.

Otologists treat AOM as:

- ***An active inflammatory state***
- ***A temporary distortion of middle ear mechanics***
- ***A condition where interpretation must wait***

Audiology must treat AOM as a clinical pause, not a data opportunity.

Active disease distorts data before it distorts hearing.



▲ **Figure 17–1.** Otoscopic examination of acute otitis media.

1. What Acute Otitis Media Really Is

AOM is characterized by:

- ***Acute inflammation of the middle ear***
- ***Rapid symptom onset***
- ***Pain, fever, irritability***
- ***Hyperemic or bulging tympanic membrane***

Key distinction from OME:

- ***AOM = inflammation + symptoms***
- ***OME = fluid without inflammation***

This distinction defines audiology's role.

2. Why AOM Invalidates Audiological Testing

During AOM:

- ***TM mobility is severely altered***
- ***Middle ear pressure is unstable***

- *Pain affects behavioral responses*
- *Patient cooperation is unreliable*

Audiological findings may show:

- *Marked conductive loss*
- *Flat tympanograms*
- *Inconsistent thresholds*

None of these reflect baseline hearing.

AOM creates temporary mechanics, not permanent hearing loss.

3. Tympanometry During AOM — A False Sense of Certainty

Tympanometry during AOM often yields:

- *Type B curves*
- *Reduced compliance*

Clinical danger:

- *Mislabeling fluid severity*
- *Over-documenting "loss"*
- *Triggering unnecessary concern*

Otologists do not base decisions on tympanograms during acute inflammation.

Audiology should follow the same logic.

4. Pain, Pressure, and Behavioral Bias

Pain changes patient behavior:

- *Elevated thresholds*
- *Poor response consistency*

- ***Early fatigue***

In children:

- ***Crying and movement invalidate testing***
- ***Results often overestimate loss***

Pain biases results upward.

5. The Post-Acute Window — Still Not Baseline

Even after symptoms improve:

- ***Residual effusion may persist***
- ***TM stiffness may remain***
- ***Pressure may normalize slowly***

Audiological implication:

- ***Early post-AOM testing still reflects recovery, not baseline***
- ***Serial follow-up is essential***

Resolution of pain ≠ resolution of physiology.

6. When Audiology SHOULD Be Involved During AOM

Audiology may:

- ***Document presence of conductive impact (qualitatively)***
- ***Educate families about temporary effects***
- ***Plan follow-up timing***

Audiology should not:

- ***Label permanent loss***
- ***Fit hearing aids***
- ***Predict outcomes***

7. Counseling and Documentation — The Protective Layer

Families often hear:

“He failed the hearing test.”

Audiology must reframe:

“Testing today reflects active infection effects, not long-term hearing.”

Proper documentation protects:

- ***Patients***
- ***Clinicians***
- ***MDT decisions***

Clinical Summary

Acute otitis media creates temporary, inflammatory changes that invalidate audiological interpretation of baseline hearing.

Acute Otitis Media (AOM)

Definition

Acute otitis media is an acute inflammation of the mucosal lining of the middle ear cleft, which includes:

Middle ear cavity
Eustachian tube
Mastoid air cells

Routes of Infection

A. Through the Eustachian Tube (Most Common)

Infection spreads from:

Nasopharynx
    Adenoids
Nose

Rhinitis of common cold

Influenza

Paranasal sinuses

Sinusitis

Oropharynx

Acute tonsillitis or pharyngitis

***Other causes*:**

Passage of infected material such as vomitus

Post-nasal packing

B. Through the Tympanic Membrane (Rare)

Through a pre-existing perforation

Through ventilation tubes

C. Hematogenous Spread (Very Rare)

Seen in severe septicemia or toxemia

Causative Organisms

Common bacteria include:

Streptococcus pneumoniae

Group A beta-hemolytic streptococci

Haemophilus influenzae

Moraxella catarrhalis

Pathological Stages of AOM

1. Tubal Catarrh

Eustachian tube obstruction due to congestion

Absorption of air from middle ear

Development of negative middle ear pressure

Result

Retraction of the tympanic membrane

2. Catarrhal Otitis Media

Congestion and edema of middle ear mucosa

Accumulation of serous fluid

3. Suppurative Otitis Media (Before Perforation)

Serous fluid becomes infected

Accumulation of pus under tension

Increased middle ear pressure



▲ *Figure 17–2. Stage of suppuration: Congested tympanic membrane with a bulge with impending rupture*

4. Perforation Stage

Necrosis of a part of the tympanic membrane

Escape of mucopurulent discharge

Relief of pain

5. Resolution

Tympanic membrane heals

Fluid becomes thin and serous

May occur at any stage depending on:

- *Virulence of organism*
- *Host resistance*
- *Adequacy of treatment*



▲ *Figure 18–3. Stage of resolution: Tympanic membrane perforation following AOM (Inferior division)*

Clinical Picture

1. Stage of Tubal Catarrh

Symptoms

- ***Ear fullness***
- ***Mild hearing loss***
- ***Tinnitus***

Signs

- ***Retracted tympanic membrane***
- ***Prominent lateral process of malleus***
- ***Disturbed cone of light***
- ***Reduced mobility on pneumatic otoscopy***

2. Catarrhal Stage

Symptoms

- ***Fever, headache, malaise***
- ***Ear pain***
- ***Hearing loss***

Signs

- ***Congested tympanic membrane***
- ***Loss of luster and cone of light***

3. Suppurative Stage

Symptoms

- ***Severe throbbing ear pain***
- ***Increased deafness***

Marked systemic symptoms, especially in children

Signs

Bulging congested tympanic membrane

4. Perforation Stage

Symptoms

Sudden relief of pain

Appearance of ear discharge

Signs

Mucopurulent pulsating discharge

Central or anteroinferior perforation

Fate of Acute Otitis Media

Complete resolution

Incomplete resolution, leading to:

- *Residual deafness due to secretory otitis media*
- *Persistent perforation*
- *Persistent discharge due to adenoids or mastoiditis*

Complications

- *Intratemporal*
- *Extracranial*
- *Intracranial*

Investigations

Culture and sensitivity of ear discharge

Pure tone audiometry

Treatment

A. Medical Treatment

(If TM is intact)

Bed rest and fluids

Systemic antibiotics for 10–14 days

Analgesics

Antihistamines

Nasal decongestant drops



▲ *Figure 18–4. Drop instillation in adults*



▲ *Figure 18–5. Drop instillation in children*

B. Surgical Treatment

(Myringotomy and Drainage)

Indications

Failure of medical treatment for more than 48 hours

Bulging tympanic membrane

Development of complications

Infants, neonates, or immunocompromised patients

Procedure

Incision in the posteroinferior quadrant of the tympanic membrane

Drainage of pus

Sample sent for culture and sensitivity

Post-procedure

Continue systemic antibiotics

Use non-ototoxic topical ear drops

Avoid water entry into the ear

Special Types of AOM

A. AOM in Infancy and Early Childhood

More common due to:

Short, wide, horizontal Eustachian tube

Frequent upper respiratory infections

Immature immunity

Clinical Notes

General symptoms are more severe

Vomiting and diarrhea may occur

Bulging and perforation are less common due to thick TM

Key Points

Early treatment prevents complications

Myringotomy should not be delayed when indicated

Do not wait for bulging TM

Myringotomy is indicated if pain and fever persist for more than 48hours

Acute Otitis Media

Children vs Adults

Feature	Children	Adults
Incidence	Very common	Less common
Severity	Usually more severe	Usually milder
Eustachian Tube	Short, wide, horizontal	Long, narrow, oblique
Immune Status	Immature	Fully developed
Predisposing Factors	URTI, adenoids, bottle feeding, teething	URTI, smoking, sinusitis
General Symptoms	High fever, irritability, excessive crying	Fever, malaise
Gastrointestinal Symptoms	Vomiting and diarrhea common	Rare
Ear Pain	May be difficult to localize	Clearly localized
Otoscopic Findings	Bulging TM less common	Bulging TM more common
TM Perforation	Less common (thick TM)	More common
Hearing Loss	Conductive hearing loss, may affect speech development	Conductive hearing loss
Risk of Complications	High	Lower
Response to Treatment	Slower	Faster

Feature	*Children*	*Adults*
Indication for Myringotomy	*Persistent pain and fever > 48 hours even without bulging TM*	*Bulging TM or complications*
Audiology Impact	*Risk of speech and language delay*	*Usually temporary*
Follow-up Importance	*Mandatory*	*Recommended*

Key Points

- ***AOM is an active disease state***
- ***Testing reflects inflammation, not baseline***
- ***Tympanometry is unreliable during AOM***
- ***Timing matters more than data***

Red Flags

- ***Severe pain during testing***
- ***Bulging or hyperemic TM***

The Sentence That Changes Everything

"Audiological findings during acute otitis media reflect temporary inflammatory changes and do not represent baseline hearing."

What I Would Explain to the Family

"Because there's an active ear infection, today's test can't show true hearing.
We'll reassess once the ear heals."

Chapter

18

When to Test, When to Wait — Middle Ear Edition

Clinical Timing Rules for Safe Audiological Decisions

Why This Chapter Matters

Middle ear pathology is time-sensitive.

The same ear can produce:

- Misleading results today
- Reliable results weeks later

Errors in middle ear audiology rarely come from:

- Poor technique

They come from:

- Poor timing

Otologists think in phases of disease.

Audiologists must do the same.

Testing at the wrong time creates confident misinformation.

TEST NOW — When Audiology Is Appropriate

Proceed with audiological testing when:

- Ear is pain-free
- No active otorrhea
- Tympanic membrane is stable (even if abnormal)

- ***Middle ear condition is quiescent or chronic***
- ***Patient cooperation is reliable***
- ***Results are expected to reflect baseline physiology***

Typical examples:

- ***Chronic OME under monitoring***
- ***Stable TM perforation***
- ***Healed AOM (after recovery window)***
- ***Long-standing retraction without acute inflammation***

Stable disease allows meaningful measurement.

WAIT — When Audiology Must Pause

Defer definitive audiological testing when:

- ***Acute otitis media is present***
- ***Significant ear pain or fever exists***
- ***Active otorrhea is ongoing***
- ***Canal or TM inflammation prevents safe testing***
- ***Recent ear surgery without clearance***
- ***Patient distress compromises cooperation***

In these situations:

- ***Testing reflects disease activity, not hearing status***
- ***Results exaggerate loss***
- ***Documentation becomes risky***

Waiting is not delay — it is accuracy.

GRAY ZONE — Test With Caution

Situations requiring qualified interpretation:

- *Early post-AOM recovery*
- *Residual effusion after symptom resolution*
- *Fluctuating OME*
- *Mild TM retraction with variable pressure*

Audiology actions:

- *Qualify all conclusions*
- *Avoid irreversible recommendations*
- *Plan serial follow-up*

If you must test, you must qualify.

Mandatory Documentation Language

Use protective phrasing:

"Audiological findings are interpreted within the context of current middle ear status and may not reflect baseline hearing."

Avoid:

- *"Permanent"*
- *"Progressive"*
- *"Definitive"*

Red Flags to WAIT

- *Pain*
- *Fever*
- *Active discharge*
- *Bulging TM*



Chapter

19

Chronic Suppurative Otitis Media (CSOM)

Chronic suppurative otitis media is a chronic inflammation of the mucosal lining of the middle ear cleft caused by pyogenic organisms.

It is characterized by

- ***Persistent or intermittent ear discharge***
- ***Permanent tympanic membrane perforation***
- ***Conductive or mixed hearing loss***

The pathological changes are irreversible.

Classification of CSOM

CSOM is classified into two main types:

1. Tubotympanic Type

(Safe type)

2. Atticoantral Type

(Unsafe type)

This classification is clinically and surgically very important.

I. Tubotympanic Type (Safe Type)

Etiology

Usually follows inadequately treated acute otitis media due to

- ***Inadequate antibiotic dose or duration***
- ***Inadequate drainage due to small or high perforation***
- ***Repeated middle ear infections through***

– Eustachian tube

– Persistent tympanic membrane perforation

- ***Low host resistance***

– Malnutrition

– Anemia

– Diabetes mellitus

Clinical Picture

Symptoms

- ***Persistent or intermittent ear discharge***
- ***Hearing loss***
- ***Tinnitus***

Signs

Ear Discharge

- ***Mucoid or mucopurulent***
- ***Profuse***
- ***Usually non offensive***
- ***May be pulsatile during acute exacerbations***

Tympanic Membrane

- Central perforation in pars tensa
- Any size or shape
- A rim of tympanic membrane present all around
- Never reaches the bony annulus

Middle Ear Mucosa

- Edematous and congested during activity
- Granulations may be present
- Aural polyp may be seen protruding through the perforation

Important Clinical Warning

Aural polyp should never be pulled blindly, as it may be attached to the ossicles or facial nerve.

Hearing Loss

- Conductive hearing loss of variable degree
- Ossicular chain is usually intact
- Ossicular erosion may occur in long standing cases

Investigations

- Pure tone audiometry

– Variable degree of conductive hearing loss

- Culture and sensitivity of ear discharge
- CT scan of temporal bone if surgery is planned or complications suspected

Treatment

Conservative Treatment

Aim is to obtain a safe dry ear.

- ***Systemic antibiotics according to culture and sensitivity***
- ***Topical antibiotic ear drops***

– Avoid ototoxic drops

- ***Regular aural toilet***
- ***Avoid water entry into the ear***
- ***Control nasal and nasopharyngeal infections***

Surgical Treatment

- ***Tympanoplasty***
- ***Tympanoplasty with cortical mastoidectomy if mastoid reservoir is suspected***
- ***Grafting materials***

– Temporalis fascia

– Tragal cartilage

II. Atticoantral Type (Unsafe Type)

Pathology

- ***Disease involves attic, antrum, and mastoid***
- ***Lining changes to keratinizing squamous epithelium***
- ***Associated with cholesteatoma***
- ***Bone erosion is common***

Clinical Picture

Symptoms

- ***Scanty persistent ear discharge***
- ***Hearing loss***

- ***Tinnitus***
- ***Symptoms of complications may appear***

Signs

Ear Discharge

- ***Purulent***
- ***Scanty***
- ***Foul smelling***

Tympanic Membrane

- ***Attic perforation***
- ***Marginal perforation***
- ***Retraction pocket***
- ***Cholesteatoma flakes may be seen***

Granulations and Polyps

- ***Common***
- ***Fleshy and bleed easily***

Hearing Loss

- ***Conductive hearing loss***
- ***May become mixed hearing loss***

Investigations

- ***Pure tone audiometry***
- ***CT scan of temporal bone***

– To detect cholesteatoma

– Bone erosion

– Ossicular status

- ***Culture and sensitivity of discharge***

Treatment

Surgery is mandatory

- ***Canal wall down mastoidectomy***
- ***Canal wall up mastoidectomy with second look surgery***
- ***Medical treatment is only temporary and preparatory***

Comparison Between Safe and Unsafe CSOM

Feature	Safe CSOM	Unsafe CSOM
Site	Tubotympanic	Atticoantral
Perforation	Central	Attic or marginal
Discharge	Profuse, non offensive	Scanty, foul smelling
Bone erosion	Rare	Common
Cholesteatoma	Absent	Present
Complications	Rare	Common
Treatment	Tympanoplasty	Mastoid surgery

Golden Rule

Any ear with foul smelling discharge and attic or marginal perforation is unsafe until proven otherwise.

Chronic Otitis Media (Non-Cholesteatomatous)

Stable Disease, Unstable Consequences

Chronic Otitis Media (COM) without cholesteatoma is often underestimated.

It is described as:

- ***"Dry ear"***
- ***"Simple perforation"***
- ***"Just conductive loss"***

Otologists know better.

Non-cholesteatomatous COM is:

- ***A chronic inflammatory state***
- ***A marker of long-standing middle ear dysfunction***
- ***A condition where stability—not severity—guides decisions***

Audiology plays a critical role in:

- ***Determining functional impact***
- ***Assessing stability***
- ***Preventing false reassurance***

Chronic does not mean harmless.

1. What Defines Non-Cholesteatomatous COM

Key features:

- ***Persistent TM perforation or chronic middle ear changes***
- ***Absence of keratinizing epithelial disease***
- ***Variable middle ear ventilation***
- ***History of recurrent infection or long-standing effusion***

This is disease memory, not active infection.

2. Pathophysiology — The Legacy of Recurrent Disease

Chronic inflammation leads to:

- *Altered mucosal lining*
- *Reduced middle ear compliance*
- *Ossicular erosion or fixation (without cholesteatoma)*
- *TM scarring and loss of elasticity*

The middle ear becomes:

- *Functionally compromised*
- *Structurally altered*
- *Less predictable*

The ear remembers every insult.



▲ *Figure 19–1. Congenital cholesteatoma with intact tympanic membrane, seen as a small pearl of cholesteatoma (C) in the anterosuperior quadrant.*



▲ *Figure 19–2. Congenital cholesteatoma.*



▲ *Figure 19–3. Acquired cholesteatoma. (a) Attic cholesteatoma. (b) Pars tensa retraction with cholesteatoma visible in the posterosuperior quadrant.*


Bone
Fibrous stroma
Matrix of stratified squamous epithelium
Keratin mass

▲ Figure 19–4. Cholesteatoma structure: Stroma, matrix and keratin mass

3. Audiological Characteristics

Typical findings include:

- *Conductive hearing loss (often moderate)*
- *Mixed loss in long-standing cases*
- *Reduced speech audibility*
- *Variable thresholds between visits*

Important nuance:

- *Degree of loss does not equal disease severity*
- *Stability over time is the key audiological signal*

4. Tympanometry and COM — Use with Caution

Possible findings:

- *Flat or reduced-mobility curves*
- *Large ear canal volume with perforation*

- *Irregular patterns with scarring*

Audiological danger:

- *Over-interpreting curves*
- *Ignoring otoscopic correlation*

In COM, the tympanogram supports the story—it never leads it.

5. The “Dry Ear” Myth

A dry ear may still:

- *Harbor chronic mucosal disease*
- *Be prone to recurrent breakdown*
- *Carry long-term functional consequences*

Otologists ask:

- *How long has it been dry?*
- *How stable is it?*
- *What is the tissue quality?*

Audiology must reflect this thinking.

6. Stability vs Progression — The Core Question

Otologic decision-making hinges on:

- *Stability over months, not days*
- *Predictability, not optimism*

Audiology contributes by:

- *Tracking thresholds longitudinally*
- *Monitoring speech outcomes*
- *Flagging subtle deterioration*

Progression whispers before it declares itself.

7. When Non-Cholesteatomatous COM Becomes Dangerous

Red flags suggesting evolution toward higher risk:

- ***Increasing discharge frequency***
- ***Progressive conductive or mixed loss***
- ***New asymmetry***
- ***Poor response to expected amplification benefit***

Audiology must escalate—not normalize—these patterns.

8. Audiology's Role in Management

Audiology should:

- ***Describe function clearly***
- ***Emphasize stability or change***
- ***Avoid surgical predictions***
- ***Support MDT decisions with longitudinal data***

Audiology should not:

- ***Reassure based on one "good" test***
- ***Label disease as benign***
- ***Promise outcomes***

Audiology guards the timeline.

The Golden Rule

In chronic otitis media, stability over time is more important than degree of loss.

What I Would Explain to the Family

“The ear isn’t infected now, but it has a long history.

We monitor it carefully because chronic ears can change quietly.”

Cholesteatoma — Foundations & Red Flags

The Quiet Disease That Destroys Without Pain

Cholesteatoma is one of the most dangerous conditions in otology

not because it is rare,

but because it is often silent.

It is frequently missed because:

- ***Hearing loss may be mild***
- ***Pain may be absent***
- ***Tympanic membrane may appear “almost normal”***

Otologists treat cholesteatoma as:

- ***A progressive destructive disease***
- ***A time-sensitive surgical condition***
- ***A non-negotiable diagnosis***

Audiology’s role is not to diagnose cholesteatoma —

but to never miss the warning signs.

Cholesteatoma is not an infection.

It is a behavior.

1. What Cholesteatoma Really Is

Cholesteatoma is:

- ***Keratinizing squamous epithelium***
- ***Located in the middle ear or mastoid***
- ***Capable of progressive expansion and destruction***

It is not:

- ***A tumor***
- ***A simple retraction***
- ***A benign finding***

Its danger lies in:

- ***Bone erosion***
- ***Ossicular destruction***
- ***Labyrinthine involvement***
- ***Facial nerve injury***
- ***Intracranial complications***

Cholesteatoma destroys by pressure and enzymes — quietly.



▲ ***Figure 19–5. A. The typical spread of a pars flaccida cholesteatoma lateral to the incus body. B. The typical spread of a pars tensa cholesteatoma medial to the incus body.***



▲ **Figure 19–6. *Primary acquired cholesteatoma in the pars flaccida portion of the left tympanic membrane.***
The arrowhead points to retraction. The arrow points to the cholesteatoma sac behind the tympanic membrane.



▲ **Figure 19–7. *The exposure obtained during a canal wall-up mastoidectomy of the right ear.***
A cholesteatoma arising from the pars flaccida and extending into the mesotympanum, the epitympanum, and the mastoid is depicted.

2. Pathophysiology — How Damage Occurs

Cholesteatoma causes damage through:

- ***Continuous keratin accumulation***
- ***Chronic inflammation***

- ***Enzymatic bone resorption***

Structures at risk:

- ***Ossicular chain***
- ***Facial nerve canal***
- ***Lateral semicircular canal***
- ***Tegmen tympani***

Audiological relevance:

- ***Destruction may precede symptoms***
- ***Audiograms often underestimate severity***

3. Types of Cholesteatoma (Clinically Relevant)

Acquired Cholesteatoma

- ***Often arises from retraction pockets***
- ***Commonly pars flaccida origin***

Congenital Cholesteatoma

- ***Occurs behind intact TM***
- ***Often detected incidentally***

Audiological danger:

An intact TM does not exclude cholesteatoma.

4. Audiological Characteristics — Why They Mislead

Typical findings may include:

- ***Conductive hearing loss***
- ***Mixed loss in advanced disease***
- ***Relatively preserved thresholds early***

Critical point:

- ***Degree of hearing loss does not reflect disease extent***

Cholesteatoma severity is anatomical, not audiometric.

5. Otoscopic Red Flags — The Audiologist's Lifeline

Never ignore the following:

- ***Deep pars flaccida retraction***
- ***Retraction pocket with debris***
- ***Foul-smelling discharge***
- ***Marginal perforation***
- ***White mass behind TM***
- ***Recurrent unilateral disease***

If debris cannot be visualized fully, assume risk.

6. Audiological Red Flags — Subtle but Deadly

High-risk audiological patterns:

- ***Progressive conductive loss***
- ***Sudden change in stable COM***
- ***New asymmetry***
- ***Poor speech understanding disproportionate to thresholds***
- ***Unexpected mixed component***

Audiology must escalate, not normalize.

7. Tympanometry and Cholesteatoma — Limited but Informative

Possible patterns:

- ***Flat or irregular curves***
- ***Apparent normal curves in early disease***

Danger:

- ***False reassurance***

Tympanometry cannot rule out cholesteatoma.

8. What Audiology Must NEVER Do

Audiology must never:

- ***Reassure without ENT review***
- ***Delay referral for repeat testing***
- ***Fit hearing aids in suspicious ears***
- ***Downplay debris or odor***

Cholesteatoma is not a "monitor and see" disease.

9. Audiology's Correct Role

Audiology should:

- ***Recognize red flags***
- ***Document findings precisely***
- ***Trigger urgent ENT referral***
- ***Avoid definitive functional labeling***

Audiology is the early warning system.

Clinical Summary

Cholesteatoma is a progressive, destructive disease whose severity is often underestimated by audiological findings.

Key Points

- ***Silent progression is common***
- ***Hearing loss underestimates disease***
- ***Otoscopic red flags are critical***

- ***Urgency saves structure***

The Golden Rule

When cholesteatoma is suspected, audiology stops measuring and starts escalating.

Absolute Red Flags

- ***Retraction with debris***
- ***Marginal perforation***
- ***Foul discharge***
- ***Unilateral progressive disease***

The Sentence That Saves You

"Findings raise concern for possible cholesteatoma, and urgent otologic evaluation is recommended."

What I Would Explain to the Family

"There's a growth behind the eardrum that can damage the ear quietly.
It needs specialist treatment quickly."

Chapter

20

Complications of Otitis Media

When a Treatable Disease Becomes a Threat

Why This Chapter Matters

Otitis media is often perceived as:

- ***Common***
- ***Treatable***
- ***Benign***

This perception is dangerous.

Otitis media—acute or chronic—can progress to life-threatening complications

when disease behavior is underestimated or intervention is delayed.

Otologists fear complications not because they are frequent,

but because they are preventable.

Audiology's role is critical:

- ***Recognizing warning patterns***
- ***Escalating early***
- ***Avoiding false reassurance***

Complications do not announce themselves loudly.

They evolve quietly.

1. How Complications Develop

Complications arise due to:

- *Persistent infection*
- *Chronic inflammation*
- *Bone erosion*
- *Poor ventilation and drainage*

Pathways of spread:

- *Direct bony erosion*
- *Preformed anatomical pathways*
- *Vascular channels*

The middle ear has no respect for boundaries.

2. Extracranial Complications

Mastoiditis

- *Most common complication*

Clinical features:

- *Postauricular pain or swelling*
- *Fever*
- *Ear protrusion*

Audiological implication:

- *Rapid deterioration of hearing*
- *Pain-limited testing*

Mastoiditis is failure of containment.

Subperiosteal Abscess

- *Extension of mastoid infection*

Clinical urgency:

- *Surgical emergency*

Audiology role:

- *Recognize severity*
- *Do not delay referral*

Facial Nerve Paralysis

- *Due to inflammation or erosion of facial canal*

Key point:

- *May occur without massive hearing loss*

Audiology must:

- *Treat this as urgent*
- *Avoid focusing on thresholds*

Facial weakness outweighs any audiogram.

Labyrinthitis / Labyrinthine Fistula

- *Inner ear involvement*

Clinical features:

- *Sudden SNHL*
- *Vertigo*
- *Imbalance*

Audiological danger:

- *Mislabeling as idiopathic SNHL*

Middle ear disease can cross into the inner ear.

3. Intracranial Complications

Meningitis

- ***Most feared complication***

Clinical signs:

- ***Headache***
- ***Fever***
- ***Neck stiffness***

Audiology implication:

- ***Immediate escalation***
- ***Testing is irrelevant***

Brain Abscess

- ***Temporal lobe or cerebellar***

Audiological relevance:

- ***Hearing loss may be minimal***
- ***Neurological signs dominate***

The ear can be the doorway to the brain.

Lateral Sinus Thrombosis

- ***Venous complication***

Clinical features:

- ***Persistent fever***
- ***Headache***
- ***Papilledema***

Audiology role:

- ***Recognize systemic red flags***


Subdural abscess
Brain abscess
Meningitis
Petrositis
Extradural abscess
Labyrinthitis
Facial paralysis
Lateral sinus thrombophlebitis
Mastoiditis

▲ *Figure 20–1. Complications of OM*


Temporal lobe abscess
Suppurative labyrinthitis
Epidural abscess
Cerebellar abscess
Subdural abscess
Zygomatic abscess
Lateral sinus thrombosis
Postauricular abscess
Infectious facial neuritis
Bezold abscess

▲ *Figure 20–2. Intracranial complications of OM.*


Thrombophlebitis
Erosion of bone by granulation tissue
Preformed pathway fracture line in tegmen

▲ *Figure 20–3. Mode of spread of infection*


Skull
Subperiosteal abscess
Extradural abscess
Subdural abscess

▲ *Figure 20–4. Extradural abscess, subdural abscess, and subperiosteal abscess*

4. Audiological Red Flags That Predict Complications

Never ignore:

- ***Sudden worsening of hearing in COM***
- ***New vertigo***
- ***Facial weakness***
- ***Severe headache***
- ***Persistent fever***
- ***Neurological symptoms***

Audiology often sees the first crack before collapse.

5. Why Mild Hearing Loss Does NOT Equal Mild Disease

Key misconception:

- ***"If hearing loss is mild, disease is mild."***

Reality:

- ***Complications relate to anatomical spread, not thresholds***

Severity of disease is spatial, not audiometric.

6. Audiology's Correct Response to Suspected Complications

Audiology must:

- ***Stop routine testing***
- ***Document red flags***
- ***Trigger urgent referral***
- ***Avoid reassurance***

Audiology must NOT:

- ***Schedule routine follow-up***

- ***Repeat testing "to be sure"***
- ***Downplay systemic symptoms***

Clinical Summary

Otitis media can progress beyond the middle ear, causing extracranial and intracranial complications with serious morbidity.

Complications of Chronic Suppurative Otitis Media (CSOM)

Definition

Complications of CSOM occur when infection spreads beyond the mucoperiosteal lining of the middle ear cleft to involve bone, soft tissues, or intracranial structures.

They are much more common in unsafe CSOM (cholesteatoma).

Causes

- ***Long-standing CSOM***
- ***Cholesteatoma, the commonest cause***
- ***Acute exacerbation of CSOM***
- ***Inadequate or irregular treatment***

Classification of Complications

Complications of CSOM are classified into three main groups:

I. Intratemporal (Cranial) Complications

(Within the temporal bone)

1. Mastoiditis

- *Infection spreads to mastoid air cells*
- *Pain and tenderness behind the ear*
- *Post-auricular swelling*
- *Loss of post-auricular groove*

2. Mastoid Abscess

- *Collection of pus beneath periosteum*
- *Ear is pushed forward and downward*
- *Requires urgent surgical drainage*


Zygomatic
Postauricular
Bezold

▲ Figure 20–5. Type of mastoid abscesses


Mastoid abscess bursting through medial side of its tip
Under the sternocleido-mastoid muscle
Styloid process
Facial nerve
Submandibular triangle
Mandible
Digastric muscle
Posterior and anterior bellies

▲ Figure 20–6. Bezold's abscess. Through the mastoid tip abscess extends either deeper to sternocleidomastoid muscle or into the submandibular triangle.

3. Petrositis

- ***Infection spreads to petrous apex***
- ***Classical Gradenigo syndrome***

– Deep facial pain (5th nerve)

– Abducens nerve palsy (6th nerve)

– Persistent ear discharge

4. Otitic Labyrinthitis

- ***Infection spreads to inner ear***
- ***Severe vertigo***
- ***Nystagmus***
- ***Sudden sensorineural hearing loss***
- ***Nausea and vomiting***

5. Labyrinthine Fistula

- ***Erosion of lateral semicircular canal***
- ***Vertigo induced by pressure or sound***
- ***Positive fistula test***

6. Facial Nerve Paralysis

- ***Due to erosion of facial canal***
- ***More common in cholesteatoma***
- ***Lower motor neuron facial palsy***

Red Flag

Facial paralysis in CSOM is always pathological until proven otherwise.

II. Intracranial Complications

1. Extradural Abscess

- Collection of pus between bone and dura
- Often silent
- Headache and fever

2. Subdural Abscess

- Severe headache
- Fever
- Neck rigidity
- Altered consciousness

3. Brain Abscess

Common sites

- Temporal lobe abscess
- Cerebellar abscess

Clinical features

- Persistent headache
- Vomiting
- Papilledemau

4. Meningitis

- High fever
- Neck stiffness
- Altered sensorium
- Rapidly progressive and life-threatening


Cranial bone
Extradural space
Intracranial dural venous sinus
Dura mater
Cerebral cortex
Rupturing in subarachnoid space
Brain abscess
Rupturing in ventricle
Ventricle

▲ *Figure 20–7.* ***Brain abscess rupture in ventricle and subarachnoid space. Rupture usually occurs in the ventricle because encapsulation of abscess is more on outer side.***

5. Lateral Sinus Thrombosis

- ***Septic thrombosis of sigmoid sinus***
- ***Spiking fever***
- ***Rigors***
- ***Headache***
- ***Papilledema***

III. Extracranial Complications

1. Post-auricular Abscess

- ***Extension of mastoid infection***
- ***Fluctuant swelling behind ear***

2. Zygomatic Abscess

- Swelling over zygomatic root
- Seen in extensive attic disease

3. Neck Abscess

- Spread through mastoid tip
- Presents as upper neck swelling

Investigations in Complicated CSOM

- CT scan of temporal bone
- MRI brain if intracranial complication suspected
- Culture and sensitivity of ear discharge
- Audiological assessment

Treatment Principles

- Hospital admission
- High-dose intravenous antibiotics
- Urgent surgical management
  - Mastoidectomy
  - Drainage of abscess
- Management of intracranial complications in coordination with neurosurgery

Key Points

- Complications are common in unsafe CSOM
- Cholesteatoma is the main risk factor
- Intracranial complications are life-threatening
- Early diagnosis and surgery save life and hearing

Golden Rule

Any CSOM patient with headache, vertigo, facial weakness, or altered consciousness must be treated as an emergency.

Flowchart

From CSOM to Complications

Chronic Suppurative Otitis Media (CSOM)

↓

Persistent middle ear infection

↓

Bone erosion (especially with cholesteatoma)

↓

| Intratemporal Spread |

↓

Mastoiditis

↓

Mastoid abscess

↓

- ***Facial nerve paralysis***
- ***Labyrinthitis***

- *Labyrinthine fistula*
- *Petrositis*

↓

| *Intracranial Spread* |

↓

- *Extradural abscess*
- *Subdural abscess*
- *Brain abscess*
- *Meningitis*
- *Lateral sinus thrombosis*

↓

Life-threatening condition

Clinical Message

Progression is usually slow but relentless, especially in unsafe CSOM.

Absence of pain does not mean absence of danger.

Audiology Red Flags in CSOM

These are warning signs that every audiologist must recognize immediately.

Major Red Flags

1. Persistent Foul-Smelling Ear Discharge

- ***Suggests cholesteatoma***
- ***Indicates unsafe CSOM***

Immediate ENT referral required

2. Sudden Worsening of Hearing

- ***Especially sensorineural component***
- ***May indicate:***

– Labyrinthitis

– Labyrinthine fistula

Audiologic emergency

3. Vertigo in a CSOM Patient

- ***Never normal***
- ***Suggests inner ear involvement***

Stop audiologic testing and refer urgently

4. Facial Weakness or Asymmetry

- ***Indicates facial nerve involvement***
- ***Always pathological in CSOM***

Emergency referral

5. Severe or Persistent Headache

- ***Especially with ear discharge***
- ***May indicate intracranial complication***

Immediate hospital referral

6. Tinnitus with Neurological Symptoms

- *May indicate inner ear or intracranial spread*

7. Fever with CSOM

- *Suggests acute exacerbation or complication*

Audiologic Test Red Flags

Pure Tone Audiometry

- *Mixed or sudden sensorineural hearing loss*
- *Rapid progression of thresholds*

Speech Audiometry

- *Poor speech discrimination not explained by PTA*

Tympanometry

- *Flat curve with active discharge is expected*
- *BUT change in pattern with worsening symptoms is suspicious*

Vestibular Tests

- *Presence of spontaneous nystagmus*
- *Positive fistula test*

Golden Audiology Rule

CSOM with new symptoms is never "stable disease".

Any change equals danger until proven otherwise.

Quick Audiology Action Box

If a CSOM patient presents with:

Vertigo

Facial weakness

Sudden SNHL

Severe headache

Do not proceed with routine audiologic testing

Document findings clearly

Refer urgently to ENT or emergency services

Audiology Checklist

Before Testing a Patient with CSOM

This checklist helps the audiologist decide

Is it safe to test, what to test, and when to stop and refer.

1. History Checklist

Before any audiologic test, confirm the following:

☐ *Duration of ear discharge*

☐ *Nature of discharge*

- *Profuse or scanty*
- *Foul smelling or non offensive*

☐ *Presence of ear pain*

☐ *Recent increase in discharge or pain*

☐ *History of vertigo or dizziness*

☐ *History of tinnitus*

☐ *History of facial weakness*

☐ *History of headache or fever*

☐ *Previous ear surgery*

Red Flag

Recent change in symptoms in a known CSOM patient.

2. Otoscopic Examination Checklist

Always examine the ear before testing.

☐ Type of perforation identified

- ***Central***
- ***Attic***
- ***Marginal***

☐ Condition of middle ear mucosa

- ***Pale***
- ***Congested***
- ***Granulations***

☐ Presence of cholesteatoma flakes

☐ Presence of aural polyp

☐ Active discharge present

Red Flags

- ***Attic or marginal perforation***
- ***Foul smelling discharge***
- ***Granulations or polyp***

Unsafe CSOM suspected.

3. Safety Checklist

Decide if Testing Is Allowed

☐ ***No vertigo***

☐ ***No facial weakness***

☐ ***No severe headache***

☐ ***No fever***

☐ ***Patient clinically stable***

If any item is positive

Stop routine testing and refer urgently.

4. Audiologic Test Selection Checklist

Pure Tone Audiometry

☐ ***Safe to perform***

☐ ***Expect conductive or mixed hearing loss***

☐ ***Monitor for sudden SNHL***

Stop test if vertigo or discomfort occurs.

Speech Audiometry

☐ ***Allowed***

☐ ***Compare speech discrimination with PTA***

☐ ***Poor score may indicate inner ear involvement***

Tympanometry

☐ ***Usually NOT reliable with active discharge***

☐ ***Avoid pressurization in suspected fistula***

Acoustic Reflexes

☐ ***Not routinely indicated***

☐ ***Avoid in active unsafe CSOM***

Vestibular Tests

☐ ***Generally CONTRAINDICATED***

☐ ***Especially if vertigo or suspected labyrinthine fistula***

5. During Testing Checklist

☐ ***Monitor patient comfort continuously***

☐ ***Ask about dizziness during testing***

☐ ***Stop immediately if vertigo or nausea occurs***

6. Documentation Checklist

☐ ***Clearly document presence of CSOM***

☐ ***Document active discharge***

☐ ***Document any red flags***

☐ ***Use safe language***

Example of safe wording

"Audiologic testing was limited due to active chronic suppurative otitis media.

Findings should be interpreted with caution and correlated clinically."

7. Immediate Referral Checklist

Refer urgently if any of the following are present:

☐ ***Vertigo***

☐ ***Facial nerve weakness***

☐ ***Sudden sensorineural hearing loss***

☐ ***Severe headache***

☐ ***Fever***

☐ ***Foul smelling discharge with attic or marginal perforation***

Golden Audiology Rule

In CSOM, safety comes before data.

If the ear is unstable, testing can wait but complications cannot.

Audiology Medico-Legal Documentation Template

Chronic Suppurative Otitis Media (CSOM)

Reason for Referral

Audiologic assessment requested for a patient with a known or suspected history of chronic suppurative otitis media.

Relevant History

The patient reports the following:

- ☐ ***Chronic ear discharge***
- ☐ ***Duration of discharge: __________***
- ☐ ***Nature of discharge:***
  - ☐ ***Profuse***
  - ☐ ***Scanty***
  - ☐ ***Foul smelling***
  - ☐ ***Non offensive***
- ☐ ***Ear pain***
- ☐ ***Vertigo or dizziness***
- ☐ ***Tinnitus***
- ☐ ***Facial weakness***
- ☐ ***Headache or fever***
- ☐ ***Previous ear surgery***

Otoscopic Findings

Otoscopic examination revealed:

- ☐ ***Tympanic membrane perforation***
  - ☐ ***Central***
  - ☐ ***Attic***
  - ☐ ***Marginal***
- ☐ ***Active discharge present***
- ☐ ***Granulation tissue present***

☐ *Aural polyp present*

☐ *Suspicion of cholesteatoma*

Clinical note

Findings are consistent with chronic middle ear disease.

Audiologic Testing Performed

Testing was selected based on patient safety and clinical stability.

- ☐ *Pure tone audiometry*
- ☐ *Speech audiometry*
- ☐ *Tympanometry*
    - ☐ *Not performed due to active discharge or safety concerns*
- ☐ *Acoustic reflex testing*
    - ☐ *Not indicated*
- ☐ *Vestibular testing*
    - ☐ *Contraindicated in the current clinical condition*

Audiologic Results Summary

Hearing loss pattern:

- ☐ *Conductive*
- ☐ *Mixed*
- ☐ *Sensorineural component suspected*

Degree of hearing loss: __________

Speech discrimination:

- ☐ *Consistent with pure tone findings*
- ☐ *Poorer than expected*

Safety and Limitation Statement

Audiologic testing was limited due to the presence of chronic suppurative otitis media and associated clinical findings.

Certain tests were deferred to avoid patient discomfort or potential risk.

Red Flag Assessment

During assessment, the following red flags were:

- ☐ *Absent*
- ☐ *Present, specify:*
  - ☐ *Vertigo*
  - ☐ *Facial nerve weakness*
  - ☐ *Sudden worsening of hearing*
  - ☐ *Severe headache*
  - ☐ *Fever*

Interpretation and Clinical Impression

Audiologic findings demonstrate hearing impairment in the context of chronic middle ear disease.

Results should be interpreted within the limitations imposed by active or unsafe CSOM.

Audiologic findings alone do not establish medical diagnosis.

Recommendation

- ☐ *Routine ENT follow-up*
- ☐ *Urgent ENT referral recommended due to red flag findings*
- ☐ *Medical or surgical management as per ENT evaluation*
- ☐ *Repeat audiologic assessment after ear becomes dry and stable*

Medico-Legal Safe Language Statement

Audiologic findings reflect functional assessment at the time of testing and are influenced by the presence of chronic suppurative otitis media.

Results should be correlated clinically and interpreted in conjunction with otologic evaluation.

No causal medical diagnosis is inferred from audiologic testing alone.

ENT Referral Letter

Chronic Suppurative Otitis Media (CSOM)

Date: __________

To: ENT Consultant

From: Audiology Department

Patient Information

Name: ________________________

Age: ________________________

Medical Record Number: ________________________

Reason for Referral

Referral for otologic evaluation and management of chronic suppurative otitis media, following audiologic assessment.

Relevant History

The patient reports:

Chronic ear discharge of __________ duration

Nature of discharge:

- ☐ Foul smelling
- ☐ Non offensive

Hearing difficulty

☐ Vertigo

☐ Tinnitus

- ☐ Ear pain
- ☐ Facial weakness
- ☐ Headache or fever

Past history includes:

- ☐ Recurrent ear infections
- ☐ Previous ear surgery

Otoscopic Findings

Audiologic otoscopic examination revealed:

- Tympanic membrane perforation:
  - ☐ Central
  - ☐ Attic
  - ☐ Marginal
- ☐ Active discharge
- ☐ Granulation tissue
- ☐ Aural polyp
- ☐ Suspicion of cholesteatoma

Audiologic Assessment Summary

Audiologic testing was performed with patient safety considerations.

- Pure tone audiometry:
  - ☐ Conductive hearing loss
  - ☐ Mixed hearing loss
  - ☐ Sensorineural component suspected

Degree of hearing loss: ___________________

Speech audiometry:

- ☐ ***Consistent with pure tone findings***
- ☐ ***Poorer than expected for thresholds***

Tympanometry:

- ☐ ***Not performed due to active discharge / safety concerns***

Red Flag Assessment

The following red flags were identified:

- ☐ ***Vertigo***
- ☐ ***Facial nerve weakness***
- ☐ ***Sudden worsening of hearing***
- ☐ ***Severe headache***
- ☐ ***Fever***

Clinical Impression

Audiologic findings indicate hearing impairment in the context of chronic middle ear disease.

The pattern and associated findings raise concern for unsafe CSOM and possible complications, warranting ENT evaluation.

Reason for Urgency

- ☐ ***Routine ENT assessment***
- ☐ ***Urgent ENT assessment recommended due to red flag findings***

Medico-Legal Safe Statement

Audiologic findings represent functional assessment at the time of testing and are influenced by active chronic middle ear disease.

Results should be interpreted in conjunction with otologic examination and imaging as deemed appropriate.

No definitive medical diagnosis is inferred from audiologic testing alone.

Recommendations

ENT evaluation and management

Consider imaging and surgical planning if indicated

Repeat audiologic assessment after ear is dry and clinically stable

The referral letter should explain why you are concerned,
not what the diagnosis is.

Red Flags

- ***Facial paralysis***
- ***Vertigo with COM***
- ***Severe headache or fever***
- ***Neurological signs***



Chapter

21

Otitic Barotrauma

When Pressure Overcomes Physiology

Why This Chapter Matters

Otitic barotrauma is not rare.

What is rare is correct audiological interpretation of barotrauma.

It occurs in:

- ***Air travel***
- ***Diving***
- ***Hyperbaric exposure***
- ***Rapid altitude or pressure changes***

Otologists view barotrauma as:

- ***A pressure–ventilation mismatch***
- ***A time-limited injury***
- ***A risk amplifier for middle ear disease***

Audiology must view it as:

A temporary mechanical insult, not a baseline hearing condition.

1. What Otitic Barotrauma Really Is

Otitic barotrauma results from:

- ***Failure of the Eustachian tube to equalize pressure***

- *Rapid ambient pressure change*

Leading to:

- *Tympanic membrane stress*
- *Middle ear mucosal injury*
- *Fluid or hemorrhage*
- *TM rupture in severe cases*

Pressure acts faster than physiology can adapt.

2. Pathophysiology — Why Damage Occurs

When pressure is unequal:

- *TM retracts or bulges*
- *Middle ear pressure becomes extreme*
- *Vascular leakage occurs*

Possible outcomes:

- *Hyperemia*
- *Serous or hemorrhagic effusion*
- *TM perforation*

Severity depends on:

- *Rate of pressure change*
- *ET function*
- *Pre-existing middle ear disease*

3. Clinical Spectrum of Barotrauma

Mild Barotrauma

- *Fullness*
- *Discomfort*

- *Transient conductive loss*

Audiological implication:

- *Often self-limited*
- *Testing may exaggerate loss*

Moderate Barotrauma

- *Pain*
- *Effusion or hemotympanum*
- *Reduced TM mobility*

Audiological implication:

- *Flat or stiff tympanograms*
- *Temporary CHL*

Severe Barotrauma

- *TM rupture*
- *Vertigo*
- *Sudden hearing loss*

Audiological implication:

- ***Urgent ENT referral***
- *Testing deferred or limited*

Severity is defined by anatomy, not symptoms alone.

4. Audiological Findings — Why They Mislead

Common findings:

- *Conductive hearing loss*
- *Type B or C tympanograms*
- *Fluctuating thresholds*

Danger:

- ***Labeling loss as permanent***
- ***Treating as chronic ETD***
- ***Overreacting to early results***

Barotrauma audiograms reflect injury phase, not outcome.

5. Tympanometry in Barotrauma — Use Carefully

Tympanometry may show:

- ***Reduced compliance***
- ***Negative pressure***
- ***Large ECV if TM ruptured***

Audiology must:

- ***Correlate with otoscopy***
- ***Avoid serial testing too early***

Pressure injuries evolve over days, not minutes.

6. Red Flags That Change Everything

Immediate ENT referral if:

- ***Severe vertigo***
- ***Sudden SNHL***
- ***Facial weakness***
- ***Persistent severe pain***
- ***Suspected perilymph fistula***

Barotrauma can cross into the inner ear.

Chapter

22

Otosclerosis



The Silent Fixation That Mimics Many Diseases

Why This Chapter Matters

Otosclerosis is often introduced as:

- A cause of conductive hearing loss
- A surgical disease
- A “good prognosis” condition

This framing is incomplete—and sometimes dangerous.

Otologists see otosclerosis as:

- A progressive bone remodeling disorder
- A disease that may involve:
    - The stapes
    - The cochlea
    - Or both
- A condition where timing and interpretation matter more than thresholds

Audiology errors in otosclerosis are common—and costly.

Otosclerosis does not announce itself loudly.

It convinces quietly.

1. What Otosclerosis Really Is

Otosclerosis is:

- ***Abnormal bone remodeling of the otic capsule***
- ***Most commonly affecting the stapes footplate***
- ***Leading to fixation and impaired sound transmission***

Key truth:

It is not purely a middle ear disease.

Cochlear involvement (otospongiosis) changes everything.

2. Pathophysiology — Why Hearing Changes

Two main mechanisms:

Stapedial Otosclerosis

- ***Fixation of stapes footplate***
- ***Impaired mechanical transmission***

Audiological result:

- ***Progressive conductive hearing loss***

Cochlear Otosclerosis

- ***Involvement of cochlear capsule***
- ***Direct inner ear damage***

Audiological result:

- ***Sensorineural or mixed hearing loss***
- ***Reduced speech discrimination***

Same disease. Very different futures.



▲ *Figure 22–1. Types of footplate*

3. Clinical Presentation — The Classic Pattern (and Its Traps)

Typical history:

- ***Slowly progressive hearing loss***
- ***Often bilateral, but asymmetric***
- ***Tinnitus common***
- ***Vertigo uncommon***

Classic audiological clues:

- ***Conductive or mixed loss***
- ***Carhart notch (often misunderstood)***

Danger:

Classic patterns are absent in many real patients.

4. Audiological Findings — What Helps and What Lies

Audiogram

- *Progressive CHL or mixed loss*
- *Bone thresholds may worsen over time*

Carhart notch:

- *Mechanical artifact*
- *Not true SNHL*
- *Disappears after stapes surgery*

Not every notch is cochlear disease.



▲ *Figure 22–2. Carhart's notch is dip in the bone conduction at 2000 Hz*

Tympanometry

- ***Often Type A***
- ***Sometimes shallow***

Danger:

- ***"Normal tymp" falsely reassures***

Normal mobility does not exclude fixation.

Acoustic Reflexes

- ***Often absent or elevated***

One of the strongest audiological clues.

5. Differential Diagnosis — Where Mistakes Happen

Otosclerosis mimics:

- ***Chronic otitis media***
- ***Ossicular discontinuity***
- ***Congenital stapes fixation***
- ***Superior canal dehiscence (pseudo-conductive loss)***



▲ Figure 22–3. Superior semicircular canal dehiscence syndrome

Audiology must:

- ***Flag suspicion***
- ***Avoid definitive labeling***

The wrong label leads to the wrong surgery.

6. Otosclerosis and Surgery — Where Audiology Must Be Careful

Audiology must NEVER:

- ***Promise surgical success***
- ***Predict hearing gain***
- ***Declare "ideal candidate" alone***

Audiology SHOULD:

- ***Describe functional status***
- ***Document stability/progression***
- ***Flag cochlear involvement risk***

Stapes surgery is irreversible.

7. When Hearing Aids Are the Better First Choice

Audiological considerations:

- ***Mild to moderate loss***
- ***Good speech discrimination***
- ***Patient preference***
- ***Medical contraindications to surgery***

Audiology protects patients by:

- ***Presenting non-surgical pathways honestly***

Surgery is an option—not a default.

8. Red Flags That Change the Conversation

High-risk features:

- ***Rapid progression***
- ***Disproportionately poor speech scores***
- ***Prominent SNHL component***
- ***Vertigo***

These suggest:

- ***Cochlear involvement***
- ***Or alternative diagnoses***

9. Counseling & Documentation — High-Stakes Language

Safe counseling:

"This condition can affect the sound transmission bones, and sometimes the inner ear as well. Different management options exist, each with benefits and risks."

Dangerous counseling:

"Surgery will fix it."

Words create expectations—and litigation.

Clinical Summary

Otosclerosis is a progressive otic capsule disorder that may affect both mechanical transmission and cochlear function.

Otosclerosis is a localized bone dystrophy of the otic capsule characterized by abnormal bone remodeling, most commonly affecting the stapes footplate, leading to progressive conductive hearing loss.

Epidemiology

Common in young and middle aged adults

More common in females

Often bilateral

Positive family history in many cases

Etiology

The exact cause is unknown, but contributing factors include:

Genetic predisposition

Hormonal influence, symptoms may worsen during pregnancy

Abnormal bone metabolism of the otic capsule

Pathology

Site of Lesion

Most commonly at the fissula ante fenestram

Leads to fixation of the stapes footplate

Types of Lesions

Otospongiotic phase, active vascular bone

Otosclerotic phase, dense sclerotic bone

Classification

1. Fenestral Otosclerosis

Involves stapes footplate

Causes conductive hearing loss

2. Cochlear Otosclerosis

Involves cochlear capsule

Causes sensorineural hearing loss

3. Mixed Otosclerosis

Combination of conductive and sensorineural hearing loss

Clinical Picture

Symptoms

- *Progressive hearing loss*
- *Hearing loss is usually bilateral*
- *Tinnitus is common*
- *Paracusis Willisii, hearing improves in noisy surroundings*
- *Vertigo is uncommon*

Otoscopic Findings

- *Tympanic membrane usually normal*
- *Pinkish hue over promontory, Schwartze sign, may be seen in active phase*

Audiological Findings

Pure Tone Audiometry

- *Conductive hearing loss with air bone gap*
- *Carhart notch at 2000 Hz*
- *Mixed loss in advanced cases*

Speech Audiometry

- *Speech discrimination usually good*
- *Reduced in cochlear involvement*

Tympanometry

- *Type As tympanogram*
- *Reduced compliance*

Acoustic Reflexes

- *Absent stapedial reflex*



▲ *Figure 22–4. Audiogram shows a 21-dB air-bone gap and depressed bone conduction threshold at 2 kHz.*

Radiological Findings

- *CT scan of temporal bone*
- *Shows hypodense lesions near oval window or cochlear capsule*

Differential Diagnosis

Condition	*Key Difference*
Ossicular discontinuity	*Large ABG with hypercompliant TM*
Serous otitis media	*Fluid behind TM*
Congenital stapes fixation	*Non progressive*

Condition	Key Difference
Tympanosclerosis	White plaques on TM

Treatment

Medical Treatment

- *Sodium fluoride may be used to slow cochlear otosclerosis*
- *Hearing aids for non surgical candidates*

Surgical Treatment

- *Stapedotomy or stapedectomy*
- *Indicated in conductive hearing loss with good cochlear reserve*


Stapedius muscle
Stapes bone and foot plate
Otosclerotic lesion
A
B
C
D
E

▲ *Figure 22–5. Types of otosclerotic lesions causing fixation of stapes footplate.*
(A) Anterior focus "fissula ante fenestram";
(B) Posterior focus;
(C) Circumferential; (D) Biscuit-type lesion;
(E) Obliterative lesion


Tympanomeatal flap
Fibrous annulus
Incudostapedial joint
Stapedius
Facial canal
Stapes footplate
Chorda tympani nerve
Posterosuperior bony canal wall overhang

▲ *Figure 22–6. Left ear stapedectomy.*
After elevating the tympanomeatal flap, posterosuperior bony canal wall is removed to get better view of stapes


Long process of incus
Stapes superstructure removed
Piston
Fenestra in stapes footplate

▲ *Figure 22–6. Stapedectomy piston. After creating a fenestra in the stapes footplate, piston is placed from long process of incus to the footplate fenestra. Note stapes superstructure is removed*


Tympanomeatal flap
Chorda tympani nerve
Stapes footplate
Facial nerve
Bony scutum removed
Incus
Stapedial tendon
Canal bone
Round window
Incision
Wire wrapped around incus
Piston with end in footplate
Base of stapes posterior crus
Chorda tympani nerve
Piston prosthesis

▲ *Figure 22–7.*
(A) An intraoperative view
(B) After removal of the stapes superstructure


Long process of incus
Facial nerve
Shea teflon piston
Gelfoam
Tissue seal of oval window

▲ *Figure 22–8. Shea teflon piston*


TM flap
Incus
VII
N
TM flap
Incus
VII
N

▲ Figure 22–9. Fenestra formation and insertion of piston



▲ Figure 22–10. Teflon piston crimped to long process of incus


Teflon piston
Nitinol piston
Soft clip titanium piston

▲ *Figure 22–11. Pistons for stapes surgery*

Prognosis

- ***Excellent hearing improvement after surgery***
- ***Risk of sensorineural hearing loss exists but is low***

Key Points

- ***Otosclerosis is a common cause of adult conductive hearing loss***
- ***Tympanic membrane is usually normal***
- ***Audiology plays a key role in diagnosis***
- ***Surgery provides excellent results***

Golden Rule

A normal tympanic membrane with progressive conductive hearing loss should always raise suspicion of otosclerosis.

Audiology Checklist

Before Stapes Surgery (Stapedotomy / Stapedectomy)

1. Patient History Checklist

☐ Progressive hearing loss

☐ Usually bilateral

☐ No history of chronic ear discharge

☐ No previous ear surgery

☐ Family history of otosclerosis

☐ Tinnitus present

☐ No active vertigo

Red Flag

History of vertigo or sudden SNHL requires re evaluation before surgery.

2. Otoscopic Examination

☐ Tympanic membrane intact

☐ Tympanic membrane appears normal

☐ No retraction or middle ear effusion

☐ No tympanosclerosis

3. Pure Tone Audiometry

☐ ***Conductive hearing loss confirmed***

☐ ***Air bone gap ≥ 20 dB***

☐ ***Bone conduction thresholds within acceptable limits***

☐ ***Presence of Carhart notch at 2000 Hz***

4. Speech Audiometry

☐ ***Good speech discrimination score***

☐ ***SDS consistent with pure tone thresholds***

Poor SDS suggests cochlear involvement.

5. Tympanometry

☐ ***Type As tympanogram***

☐ ***Reduced compliance***

☐ ***Normal middle ear pressure***

6. Acoustic Reflexes

☐ ***Absent stapedial reflex***

7. Radiology Correlation

☐ ***CT temporal bone reviewed if available***

☐ ***No alternative ossicular pathology identified***

8. Ear Selection Checklist

☐ ***Worse hearing ear selected for first surgery***

☐ ***Better cochlear reserve***

☐ ***Patient informed about bilateral nature***

9. Medico Legal Documentation

☐ ***Hearing thresholds documented clearly***

☐ ***Pre operative counseling documented***

☐ ***Risk of SNHL explained***

☐ ***Informed consent verified***

Golden Rule

Never proceed to stapes surgery without complete audiologic confirmation of otosclerosis.

Comparison Table

Otosclerosis vs Ossicular Discontinuity

Feature	*Otosclerosis*	*Ossicular Discontinuity*
Onset	*Gradual, progressive*	*Sudden or after trauma*
Laterality	*Usually bilateral*	*Usually unilateral*
History	*Family history common*	*Trauma or surgery*
TM Appearance	*Normal*	*Normal or healed*
Hearing Loss	*Progressive conductive*	*Sudden conductive*
Air Bone Gap	*Moderate, stable*	*Large, often > 40 dB*
Carhart Notch	*Present*	*Absent*
Tympanometry	*Type As*	*Type Ad*
Acoustic Reflex	*Absent*	*May be present*

Feature	*Otosclerosis*	*Ossicular Discontinuity*
Speech Discrimination	***Usually good***	***Usually excellent***
Imaging	***Fenestral lesion***	***Ossicular gap***
Surgery	***Stapedotomy***	***Ossiculoplasty***

Clinical Take Home Message

Not every conductive hearing loss with normal TM is otosclerosis.

Audiology differentiates stapes fixation from ossicular disruption.

SECTION 4

Neuro-Otology
Inner Ear
& Otological symptoms



Chapter

23

Tumors of the Middle Ear (Glomus Tumors)

When Conductive Hearing Loss Is Not Benign

Why This Chapter Matters

Middle ear tumors are uncommon.

But when missed, the consequences are severe.

They are frequently mistaken for:

- ***Chronic otitis media***
- ***Otitis media with effusion***
- ***Tympanic membrane abnormalities***

Otologists fear middle ear tumors not because of frequency,

but because delay changes outcome.

Audiology's role is not diagnosis.

It is early suspicion and escalation.

Not every conductive loss belongs to inflammation.

Glomus Tumors

Glomus tumors are benign, highly vascular tumors arising from paraganglionic cells associated with the autonomic nervous system. In the middle ear, they most commonly originate from the jugular bulb or the promontory.



▲ *Figure 23–1. Glomus tumor arising from jugular bulb*

Types

Glomus Tympanicum

- ***Arises from the cochlear promontory***
- ***Most common primary tumor of the middle ear***

Glomus Jugulare

- ***Arises from the jugular bulb***
- ***Extends upward into the middle ear and mastoid***

Pathology

- *Highly vascular tumors*
- *Slow growing but locally aggressive*
- *Supplied mainly by branches of the external carotid artery*
- *Can erode bone and invade adjacent structures*

Clinical Presentation

Symptoms

- *Pulsatile tinnitus, classic and often the first symptom*
- *Conductive hearing loss*
- *Aural fullness*
- *Otalgia, less common*
- *Vertigo or SNHL if inner ear is involved*
- *Cranial nerve symptoms in advanced cases, especially IX, X, XI*

Signs

- *Reddish pulsatile mass behind an intact tympanic membrane*
- *Brown sign, blanching of the mass on pneumatic otoscopy*
- *Possible visible mass in the external auditory canal in advanced cases*

Audiological Findings

- *Conductive hearing loss is the most common*
- *Mixed or sensorineural loss in advanced disease*
- *Speech discrimination usually preserved early*

Investigations

Imaging

- *CT temporal bone*

– *Bony erosion*

– *Extent within middle ear and mastoid*

- *MRI with contrast*

– *Salt and pepper appearance*

– *Defines soft tissue extension*

- *Angiography*

– *Identifies feeding vessels*

– *Useful before surgery or embolization*



▲ *Figure 23–2. Angiogram showing highly vascular glomus tumor*



▲ *Figure 23–3. The rising sun appearance of glomus tumor*

Differential Diagnosis

- ***High jugular bulb***
- ***Aberrant internal carotid artery***
- ***Hemangioma***
- ***Cholesteatoma, non pulsatile***

Management

Options

- ***Surgical excision, treatment of choice in fit patients***
- ***Preoperative embolization to reduce bleeding***
- ***Radiotherapy, for elderly patients or inoperable tumors***
- ***Observation in very small, asymptomatic tumors***

Audiology Red Flags

- ***Pulsatile tinnitus with conductive hearing loss***
- ***Reddish mass behind intact TM***

- ***Sudden bleeding during myringotomy or ear surgery***
- ***Do NOT attempt biopsy or middle ear surgery without imaging***

Medico Legal Note

Any suspected glomus tumor must be clearly documented and referred to ENT before audiological intervention. Failure to recognize a vascular middle ear mass carries a high medico legal risk.

Clinical Summary

Middle ear tumors often masquerade as benign conductive pathologies, making early suspicion critical.

Tumors of the Middle Ear

Glomus Tumors

Definition

Glomus tumors are benign, highly vascular tumors arising from paraganglionic cells associated with the autonomic nervous system. In the middle ear, they most commonly originate from the jugular bulb or the promontory.

Types

Glomus Tympanicum

- ***Arises from the cochlear promontory***
- ***Most common primary tumor of the middle ear***

Glomus Jugulare

- ***Arises from the jugular bulb***
- ***Extends upward into the middle ear and mastoid***

Pathology

- *Highly vascular tumors*
- *Slow growing but locally aggressive*
- *Supplied mainly by branches of the external carotid artery*
- *Can erode bone and invade adjacent structures*

Clinical Presentation

Symptoms

- *Pulsatile tinnitus, classic and often the first symptom*
- *Conductive hearing loss*
- *Aural fullness*
- *Otalgia, less common*
- *Vertigo or SNHL if inner ear is involved*
- *Cranial nerve symptoms in advanced cases, especially IX, X, XI*

Signs

- *Reddish pulsatile mass behind an intact tympanic membrane*
- *Brown sign, blanching of the mass on pneumatic otoscopy*
- *Possible visible mass in the external auditory canal in advanced cases*



▲ *Figure 23–4. Rising-sun sign seen in case of glomus tympanicus*

Audiological Findings

- ***Conductive hearing loss is the most common***
- ***Mixed or sensorineural loss in advanced disease***
- ***Speech discrimination usually preserved early***

Investigations

Imaging

- ***CT temporal bone***

– Bony erosion

– Extent within middle ear and mastoid

- ***MRI with contrast***

– Salt and pepper appearance

– Defines soft tissue extension

- ***Angiography***

– Identifies feeding vessels

– Useful before surgery or embolization

Differential Diagnosis

- ***High jugular bulb***
- ***Aberrant internal carotid artery***
- ***Hemangioma***
- ***Cholesteatoma, non pulsatile***

Management

Options

- ***Surgical excision, treatment of choice in fit patients***
- ***Preoperative embolization to reduce bleeding***

- ***Radiotherapy, for elderly patients or inoperable tumors***
- ***Observation in very small, asymptomatic tumors***

Audiology Red Flags

- ***Pulsatile tinnitus with conductive hearing loss***
- ***Reddish mass behind intact TM***
- ***Sudden bleeding during myringotomy or ear surgery***
- ***Do NOT attempt biopsy or middle ear surgery without imaging***

Medico Legal Note

Any suspected glomus tumor must be clearly documented and referred to ENT before audiological intervention. Failure to recognize a vascular middle ear mass carries a high medico legal risk.

Chapter



24

Diseases of the Inner Ear

Inner ear diseases primarily affect the cochlea and vestibular labyrinth, leading to sensorineural hearing loss, vestibular symptoms, or both. Accurate audiologic interpretation is critical for localization, counseling, and referral.

Classification of Inner Ear Diseases

Congential

Traumatic

Inflammatory

Tumors

Miscellaneous

I. Congenital Inner Ear Disorders

1. Inner Ear Aplasia

Complete absence of inner ear structures.

Clinical features

- ***Profound congenital SNHL***

- *Absent vestibular function*
- *Delayed motor milestones*

Audiology

- *No response on ABR*
- *Absent OAE*
- *No cochlear nerve response*

Key audiology note

Hearing aids are ineffective, cochlear implantation is contraindicated.

2. Mondini Syndrome

Incomplete development of the cochlea with reduced turns.

Clinical features

- *Congenital or early onset SNHL*
- *May be progressive*
- *Possible vestibular symptoms*

Audiology

- *Mild to profound SNHL*
- *Fluctuating thresholds possible*
- *Poor speech discrimination in advanced cases*

Clinical importance

High risk of CSF leak and meningitis.

II. Traumatic Inner Ear Disorders

1. Transverse Temporal Bone Fracture

Usually associated with high velocity trauma.

Clinical features

- *Sudden profound SNHL*
- *Severe vertigo*
- *Facial nerve involvement may occur*

Audiology

- *Profound SNHL*
- *Absent speech discrimination*
- *Vestibular areflexia*

2. Acoustic Trauma

Damage due to excessive noise exposure.

Types

- *Acute acoustic trauma*
- *Chronic noise induced hearing loss*

Audiology

- *High frequency SNHL*
- *Classic 4000 Hz notch*
- *OAEs reduced or absent in affected frequencies*

Audiology red flag

Sudden hearing loss after noise exposure requires urgent ENT referral.

III. Inflammatory Inner Ear Disorders

Labyrinthitis

Inflammation of the membranous labyrinth.

Types

- ***Serous labyrinthitis***
- ***Suppurative labyrinthitis***

Clinical features

- ***Acute vertigo***
- ***Nausea and vomiting***
- ***Hearing loss, usually SNHL***

Audiology

- ***SNHL, often unilateral***
- ***Poor speech discrimination***
- ***Vestibular testing shows reduced or absent responses***

Important note

Suppurative labyrinthitis is a medical emergency.

IV. Tumors of the Inner Ear

Acoustic Neuroma (Vestibular Schwannoma)

Benign tumor arising from the vestibular division of the eighth nerve.

Clinical features

- ***Unilateral progressive SNHL***
- ***Tinnitus***
- ***Poor speech discrimination disproportionate to PTA***
- ***Imbalance rather than true vertigo***

Audiology

- ***Asymmetric SNHL***

- *Poor speech discrimination*
- *Abnormal ABR, prolonged interpeak latencies*

Audiology red flag

Asymmetric SNHL always requires MRI referral.



▲ *Figure 24–1. Vestibular schwannoma tumour*
**Cerebellopontine angle or cerebellopontine cistern is a three-dimensional triangular space bounded by brainstem medially, petrous part of temporal laterally, cerebellum superiorly and posteriorly and the arachnoid space containing lower cranial nerves IX to XII inferiorly.*


7,8
IAC
CO
CG
ME
EAC
M
SCC
CPA
SS
4V
5
P
Cb
Superior
Vertical crest
(Bill's bar)
Transverse
crest
Anterior
Posterior
Inferior
7
SV
C
IV

▲ *Figure 24–2.The anatomy of the CPA and its relationship to the temporal*

V. Miscellaneous Inner Ear Disorders

Meniere Disease

Disorder of endolymphatic hydrops.

Clinical triad

- ***Episodic vertigo***
- ***Fluctuating SNHL***
- ***Tinnitus with aural fullness***

Audiology

- ***Fluctuating low frequency SNHL early***
- ***Progressive flat SNHL later***
- ***Reduced speech discrimination during attacks***

Vestibular findings

- ***Reduced caloric response on affected side***

Audiology Focus Summary

- ***Inner ear diseases primarily cause SNHL***
- ***Speech discrimination often guides retrocochlear suspicion***
- ***Asymmetry equals pathology until proven otherwise***
- ***Audiology findings must always be correlated with imaging***

Red Flags for Immediate ENT Referral

- ***Sudden SNHL***
- ***Asymmetric SNHL***
- ***Severe vertigo with hearing loss***
- ***Poor speech discrimination inconsistent with PTA***

Golden Rule

Inner ear pathology is diagnosed by pattern recognition, not by a single audiologic test.

Diseases of the Inner Ear

Inner ear diseases primarily affect the cochlea and vestibular labyrinth, leading to sensorineural hearing loss, vestibular symptoms, or both. Accurate audiologic interpretation is critical for localization, counseling, and referral.

I. Congenital Inner Ear Disorders

1. Inner Ear Aplasia

Complete absence of inner ear structures.

Clinical features

- ***Profound congenital SNHL***
- ***Absent vestibular function***
- ***Delayed motor milestones***

Audiology

- ***No response on ABR***
- ***Absent OAE***
- ***No cochlear nerve response***

Key audiology note

Hearing aids are ineffective, cochlear implantation is contraindicated.

2. Mondini Syndrome

Incomplete development of the cochlea with reduced turns.

Clinical features

- ***Congenital or early onset SNHL***
- ***May be progressive***
- ***Possible vestibular symptoms***

Audiology

- ***Mild to profound SNHL***
- ***Fluctuating thresholds possible***
- ***Poor speech discrimination in advanced cases***

Clinical importance

High risk of CSF leak and meningitis.

II. Traumatic Inner Ear Disorders

1. Transverse Temporal Bone Fracture

Usually associated with high velocity trauma.

Clinical features

- ***Sudden profound SNHL***
- ***Severe vertigo***
- ***Facial nerve involvement may occur***

Audiology

- ***Profound SNHL***
- ***Absent speech discrimination***
- ***Vestibular areflexia***


Longitudinal
Transverse

▲ Figure 24–3. Types of temporal bone fractures

2. Acoustic Trauma

Damage due to excessive noise exposure.

Types

- ***Acute acoustic trauma***
- ***Chronic noise induced hearing loss***

Audiology

- ***High frequency SNHL***

- *Classic 4000 Hz notch*
- *OAEs reduced or absent in affected frequencies*

Audiology red flag

Sudden hearing loss after noise exposure requires urgent ENT referral.

III. Inflammatory Inner Ear Disorders

Labyrinthitis

Inflammation of the membranous labyrinth.

Types

- *Serous labyrinthitis*
- *Suppurative labyrinthitis*

Clinical features

- *Acute vertigo*
- *Nausea and vomiting*
- *Hearing loss, usually SNHL*

Audiology

- *SNHL, often unilateral*
- *Poor speech discrimination*
- *Vestibular testing shows reduced or absent responses*

Important note

Suppurative labyrinthitis is a medical emergency.

IV. Tumors of the Inner Ear

Acoustic Neuroma (Vestibular Schwannoma)

Benign tumor arising from the vestibular division of the eighth nerve.

Clinical features

- ***Unilateral progressive SNHL***
- ***Tinnitus***
- ***Poor speech discrimination disproportionate to PTA***
- ***Imbalance rather than true vertigo***

Audiology

- ***Asymmetric SNHL***
- ***Poor speech discrimination***
- ***Abnormal ABR, prolonged interpeak latencies***

Audiology red flag

Asymmetric SNHL always requires MRI referral.


Extracanalicular
VIII nerve schwannoma
Internal auditory
meatus erosion
A
B
Cerebellopontine angle tumor

▲ *Figure 24–4. Site of origin of acoustic tumor*

Audiology Focus Summary

- *Inner ear diseases primarily cause SNHL*
- *Speech discrimination often guides retrocochlear suspicion*
- *Asymmetry equals pathology until proven otherwise*
- *Audiology findings must always be correlated with imaging*

Red Flags for Immediate ENT Referral

- *Sudden SNHL*
- *Asymmetric SNHL*
- *Severe vertigo with hearing loss*
- *Poor speech discrimination inconsistent with PTA*

Traumatic Inner Ear Disorders

1. Transverse Temporal Bone Fracture

Usually associated with high velocity trauma.

Clinical features

- *Sudden profound SNHL*
- *Severe vertigo*
- *Facial nerve involvement may occur*

Audiology

- *Profound SNHL*
- *Absent speech discrimination*
- *Vestibular areflexia*

2. Acoustic Trauma

Damage due to excessive noise exposure.

Types

- ***Acute acoustic trauma***
- ***Chronic noise induced hearing loss***

Audiology

- ***High frequency SNHL***
- ***Classic 4000 Hz notch***
- ***OAEs reduced or absent in affected frequencies***

Audiology red flag

Sudden hearing loss after noise exposure requires urgent ENT referral.

III. Inflammatory Inner Ear Disorders

Labyrinthitis

Inflammation of the membranous labyrinth.

Types

- ***Serous labyrinthitis***
- ***Suppurative labyrinthitis***

Clinical features

- ***Acute vertigo***
- ***Nausea and vomiting***
- ***Hearing loss, usually SNHL***

Audiology

- ***SNHL, often unilateral***
- ***Poor speech discrimination***
- ***Vestibular testing shows reduced or absent responses***

Important note

Suppurative labyrinthitis is a medical emergency.

IV. Tumors of the Inner Ear

Acoustic Neuroma (Vestibular Schwannoma)

Benign tumor arising from the vestibular division of the eighth nerve.

Clinical features

- ***Unilateral progressive SNHL***
- ***Tinnitus***
- ***Poor speech discrimination disproportionate to PTA***
- ***Imbalance rather than true vertigo***

Audiology

- ***Asymmetric SNHL***
- ***Poor speech discrimination***
- ***Abnormal ABR, prolonged interpeak latencies***

Audiology red flag

Asymmetric SNHL always requires MRI referral.

V. Miscellaneous Inner Ear Disorders

Meniere Disease

Disorder of endolymphatic hydrops.

Clinical triad

- ***Episodic vertigo***
- ***Fluctuating SNHL***
- ***Tinnitus with aural fullness***

Audiology

- ***Fluctuating low frequency SNHL early***
- ***Progressive flat SNHL later***
- ***Reduced speech discrimination during attacks***

Vestibular findings

- ***Reduced caloric response on affected side***

Audiology Focus Summary

- ***Inner ear diseases primarily cause SNHL***
- ***Speech discrimination often guides retrocochlear suspicion***
- ***Asymmetry equals pathology until proven otherwise***
- ***Audiology findings must always be correlated with imaging***

Red Flags for Immediate ENT Referral

- ***Sudden SNHL***
- ***Asymmetric SNHL***
- ***Severe vertigo with hearing loss***
- ***Poor speech discrimination inconsistent with PTA***

Golden Rule

Inner ear pathology is diagnosed by pattern recognition, not by a single audiologic test.

DIAGNOSTIC ALGORITHM

Acoustic Neuroma, Vestibular

Patient with Sensorineural Hearing Loss

↓

Is hearing loss unilateral or asymmetric?

↓

YES ***NO***

↓ ↓

Suspect retrocochlear pathology ***Likely cochlear pathology***

↓ ↓

Speech discrimination test ***Manage as cochlear***

↓ ↓

Is SDS disproportionately poor compared to PTA? ***Hearing aids, monitoring***

↓

YES

↓

Check additional red flags

- ***Unilateral tinnitus***
- ***Imbalance without true vertigo***
- ***Facial numbness***

↓

Audiology referral pathway

↓

ABR testing

↓

ABR abnormal or inconclusive?

↓

YES ***NO***

↓ ↓

MRI referral ***MRI still***

to ENT ***recommended if***

clinical suspicion persists

↓

Diagnosis confirmed by MRI

Override Rule

Sudden SNHL or rapidly progressive asymmetry bypasses ABR and goes directly to urgent ENT and MRI referral.


TRANSCHOCLEAR
TRANSLABYRINTHINE
RETROLABYRINTHINE
RETROSIGMOID
A
SS
SSCC
CC
Co
PSCC
VA
JB
IS
VV
Tentorium
JV
10
9
E
S
PA
5
11S
11C
8
Ch
Fl
BS
Cerebellum
B
C

▲ *Figure 24–5. Surgical approaches to the cerebellopontine angle*

AUDIOLOGY CHECKLIST

Before ABR and MRI Referral

Suspected Acoustic Neuroma

1. History Checklist

- ☐ ***Unilateral or asymmetric hearing loss***
- ☐ ***Progressive course***
- ☐ ***Unilateral tinnitus***
- ☐ ***Difficulty understanding speech in noise***
- ☐ ***Imbalance rather than spinning vertigo***
- ☐ ***Facial numbness or tingling***

2. Audiologic Findings Checklist

- ☐ ***Asymmetric SNHL on PTA***
- ☐ ***Poor speech discrimination disproportionate to PTA***
- ☐ ***Absent or abnormal acoustic reflexes***
- ☐ ***Reflex decay positive, if tested***
- ☐ ***OAEs present despite SNHL***

3. ABR Indication Checklist

Proceed with ABR if:

- ☐ ***Asymmetry present***

☐ *Speech discrimination suspicious*

☐ *Patient medically stable*

ABR red flags:

☐ *Prolonged interpeak latencies*

☐ *Interaural latency difference*

Normal ABR does NOT exclude small tumors.

4. MRI Referral Checklist

Recommend MRI if any of the following are present:

☐ *Unilateral or asymmetric SNHL*

☐ *Poor speech discrimination*

☐ *Abnormal or inconclusive ABR*

☐ *Persistent unilateral tinnitus*

☐ *Vestibular symptoms without peripheral pattern*

5. Documentation Safe Statement

"Audiologic findings raise suspicion of retrocochlear pathology. Imaging is recommended for definitive diagnosis."

Golden Audiology Rule

Audiology identifies risk, MRI confirms diagnosis.

Acoustic Neuroma vs Meniere Disease

Feature	*Acoustic Neuroma*	*Meniere Disease*
Pathology	***Vestibular nerve tumor***	***Endolymphatic hydrops***
Onset	***Gradual***	***Episodic***
Hearing loss	***Progressive unilateral SNHL***	***Fluctuating SNHL***
Laterality	***Usually unilateral***	***Usually unilateral early***
Tinnitus	***Continuous, unilateral***	***Roaring, fluctuating***
Vertigo	***Mild imbalance***	***Severe episodic vertigo***
Speech discrimination	***Disproportionately poor***	***Reduced during attacks***
PTA pattern	***High frequency SNHL***	***Low frequency SNHL early***
OAEs	***May be present***	***Reduced or absent***
ABR	***Abnormal***	***Usually normal***
MRI	***Diagnostic***	***Usually normal***

Final Unified Clinical Message

Fluctuating hearing with vertigo suggests Meniere disease.

Progressive asymmetric hearing with poor speech discrimination suggests acoustic neuroma.

Chapter



25

Meniere Disease

Endolymphatic Hydrops

Meniere disease is a disorder of the inner ear characterized by endolymphatic hydrops, an abnormal accumulation of endolymph within the membranous labyrinth, leading to episodic vestibular and auditory symptoms.

Etiology

The exact cause is unknown. Proposed mechanisms include:

- ***Impaired endolymph absorption***
- ***Increased endolymph production***
- ***Abnormal endolymphatic sac function***
- ***Autoimmune factors***
- ***Viral insults***

Pathophysiology

- ***Accumulation of endolymph increases inner ear pressure***
- ***Distension of the membranous labyrinth***
- ***Rupture between endolymph and perilymph compartments***
- ***Ionic imbalance leading to hair cell dysfunction***


Endolymphatic sac
Vestibulo cochlear nerve
A
Normal membranous labyrinth
Endolymphatic sac
Vestibulocochlear nerve
B
Dilated membranous labyrinth

▲ *Figure 25–1. Meniere Disease*


Reissner's membrane

▲ *Figure 25–2. Bulging of Reissner's membrane*



▲ *Figure 25–3. Normal and engorged membranous labyrinth*

Classic Triad

- ***Episodic vertigo***
- ***Fluctuating sensorineural hearing loss***
- ***Tinnitus with aural fullness***

Vertigo

- ***Severe spinning sensation***
- ***Lasts minutes to hours***
- ***Associated with nausea and vomiting***
- ***Patient prefers to lie still***

Hearing Loss

- ***Fluctuating***
- ***Typically low frequency SNHL in early stages***
- ***Progresses to flat SNHL in advanced disease***

Tinnitus

- ***Roaring or buzzing***
- ***Fluctuates with attacks***

Aural Fullness

- ***Sensation of pressure or blockage***
- ***Often precedes vertigo attacks***

Otoscopic Findings

- ***Tympanic membrane appears normal***

Audiological Findings

Pure Tone Audiometry

- ***Early disease***

– Low frequency SNHL

– Rising audiogram

- ***Late disease***

– Flat or sloping SNHL

Speech Audiometry

- ***Speech discrimination usually good between attacks***
- ***May deteriorate during acute episodes***

Immittance

- ***Tympanometry usually normal***
- ***Acoustic reflexes usually present***

Otoacoustic Emissions

- ***Reduced or absent during active hydrops***
- ***May fluctuate over time***

Vestibular Testing

- ***Reduced caloric response on the affected side***
- ***Degree of hypofunction does not correlate with vertigo severity***

Diagnosis

Diagnosis is clinical, supported by audiologic pattern and symptom history. Imaging is used to exclude retrocochlear pathology, not to confirm Meniere disease.

Differential Diagnosis

- ***Vestibular migraine***
- ***Acoustic neuroma***
- ***Labyrinthitis***
- ***Autoimmune inner ear disease***

Management

Acute Attack

- ***Vestibular suppressants***
- ***Antiemetics***
- ***Bed rest***

Long Term Management

- ***Low salt diet***
- ***Diuretics***
- ***Avoid caffeine and nicotine***
- ***Stress control***

Advanced Cases

- ***Intratympanic gentamicin***
- ***Intratympanic steroids***
- ***Surgical options in refractory cases***



▲ *Figure 24–4. Intra tympanic gentamicin injection in low dose*

Prognosis

- ***Vertigo attacks may decrease over time***
- ***Hearing loss often becomes permanent***
- ***Bilateral involvement may occur in late stages***

Audiology Focus Summary

- ***Meniere disease causes fluctuating SNHL***
- ***Low frequency loss is characteristic early***
- ***Speech discrimination is relatively preserved***
- ***Audiology tracks progression, not just diagnosis***

Audiology Red Flags

- *Poor speech discrimination out of proportion to PTA*
- *Progressive asymmetric SNHL*
- *Persistent unilateral tinnitus without vertigo*

These suggest retrocochlear pathology, not Meniere disease.

Golden Rule

Fluctuating low frequency SNHL with episodic vertigo points to cochlear hydrops, not neural disease.

Patient Counseling Sheet

Meniere Disease (Endolymphatic Hydrops)

What is Meniere Disease?

Meniere disease is a condition of the inner ear caused by excess inner ear fluid.

It affects balance and hearing and usually comes in attacks.

What Symptoms Can Happen?

You may experience:

Episodes of spinning dizziness, vertigo
Hearing loss that may change from day to day
Ringing or buzzing noise in the ear, tinnitus
A feeling of pressure or fullness in the ear

Symptoms often come and go. Between attacks, you may feel normal.

Important Things to Know

Meniere disease is not a brain tumor

It is not contagious

Attacks are unpredictable

Hearing loss may become permanent over time

Lifestyle Advice

Very Important for Symptom Control

Diet

Reduce salt intake

Avoid salty foods, pickles, chips, processed foods

Eat small meals regularly

Drinks

Reduce caffeine, coffee, tea, energy drinks

Avoid alcohol

Drink water regularly throughout the day

Smoking

Stop smoking

Nicotine worsens inner ear circulation

Stress

Stress can trigger attacks

Try relaxation techniques

Ensure good sleep

During a Vertigo Attack

- Sit or lie down immediately
- Avoid sudden head movements
- Keep your eyes open and fixed on one point
- Do not drive or walk alone
- Take prescribed medications as instructed

Hearing and Balance Expectations

- Hearing may fluctuate early in the disease
- Over time, hearing loss may become stable but permanent
- Balance symptoms often improve with time due to brain compensation

Treatment Expectations

- Medicines help control symptoms, not cure the disease
- Diet and lifestyle changes are as important as medications
- Some procedures may be recommended if attacks are severe

When to Seek Medical Advice Urgently

Contact your doctor if you have:

- Sudden severe hearing loss
- Persistent dizziness not improving
- New weakness of the face
- Severe headache or visual problems

Prognosis

- Many patients have fewer attacks over time
- Vertigo often becomes less severe after years
- Hearing loss may slowly progress

Long term follow up is important

Role of Audiology Follow Up

Monitor hearing changes

Adjust hearing aids if needed

Detect changes that need ENT review

Key Message for Patients

With correct lifestyle changes and regular follow up,

many patients with Meniere disease can live a normal and active life.

Chapter

26

The Facial Nerve and Bell's Palsy

The Facial Nerve (Seventh Cranial Nerve)

Functions

The facial nerve is a **mixed nerve** with the following components:

- **Motor**
  - Muscles of facial expression
  - Stapedius muscle
- **Sensory**
  - Taste sensation from the anterior two thirds of the tongue
- **Parasympathetic**
  - Lacrimal gland
  - Submandibular and sublingual salivary glands

Course of the Facial Nerve

Intracranial Course

- Arises from the pons
- Enters the internal auditory canal with the vestibulocochlear nerve

Intratemporal Course

Within the temporal bone, the facial nerve passes through the facial canal and has three parts:

Labyrinthine segment

- ***From internal auditory canal to geniculate ganglion***
- ***Narrowest segment***

Tympanic segment

- ***Runs horizontally above the oval window***

Mastoid segment

- ***Runs vertically to exit the skull***

Branches within the temporal bone include:

- ***Greater petrosal nerve***
- ***Nerve to stapedius***
- ***Chorda tympani***

Extracranial Course

- ***Exits the skull through the stylomastoid foramen***
- ***Passes through the parotid gland***
- ***Divides into terminal branches:***

– Temporal

– Zygomatic

– Buccal

– Marginal mandibular

– Cervical


Lacrimal gland
Trigeminal ganglion
Geniculate ganglion
VI
Superior salivatory nucleus
Motor nucleus CN VII
Nasal and palatine glands
Nucleus solitarius
Fasciculus solitarius
Superficial petrosal nerve
Pterygopalatine ganglion
Nerve to stapedius
Chorda tympani
Tongue
Lingual nerve
Submandibular ganglion
Terminal branches
Sublingual gland
Submandibular gland

▲ *Figure 26–1. Facial nerve and its branches*


Frontal
Zygomatic
Buccal
Marginal
Cervical

▲ *Figure 26–2. Facial nerve anatomy.*


Trigeminal ganglion
Vagus nerve
Ophthalmic branch N.V
Facial nerve
5
7
9
10
Maxillary branch N.V
Glossopharyngeal nerve
Mandibular branch N.V
A
Maxillary process
Meckel cartilage
Pharyngeal cleft
Mandibular arch
Hyoid arch
Cardiac bulge
B
Lesser horn of hyoid bone
Meckel cartilage
Malleus
Incus
Stapes
Styloid process
Stylohyoid lig.
I
II
III
IV
VI
Body of hyoid bone
Thyroid cartilage
Cricoid cartilage
Tracheal rings
Greater horn of hyoid bone
C

▲ *Figure 26–3. A schematic illustration demonstrating the embryology of the facial nerve*


Posterior limb of internal capsule
Motor cortex (head region)
Corticobulbar tract
Internal acoustic meatus
Temporal branch
VII motor nucleus
Facial canal
Zygomatic branch
Buccal branch
Posterior auricular nerve
Nerve to stapedius
Cervical branch
Marginal mandibular branch

▲ *Figure 26–4. A schematic illustration of the complete pathway of the motor division of the facial nerve.*


Pons
VIII
Medulla
Nervus intermedius portion of VII
VI
VII

▲ *Figure 26–5.The anatomy of the facial nerve (CN VII), cochlear nerve, and vestibular nerve (CN VIII), as they exit the brainstem at the level of the pontomedullary junction.*


Dorsal longitudinal fasciculus
Hypothalamus
Trigeminal ganglion (elevated)
Mandibular nerve
Maxillary nerve
Internal carotid artery
Foramen lacerum
Lacrimal gland
Pterygo-palatime ganglion
Lingual branch of V3
Sublingual gland
Submandibular ganglion
Submandibular gland
Chorda tympani
Greater petrosal nerve
Greater petrosal foramen
Nervus intermedius
Superior salivatory (lacrimal) nucleus

▲ *Figure 26–6. The anatomy of the visceral motor portion of the facial nerve, making up the nervus intermedius, or Wrisberg nerve.*


Gustatory area
in sensory
cortex
Ventral
posterior
nucleus of
thalamus
Nucleus solitarius
(rostral gustatory portion)
Tractus solitarius
Nervus intermedius
Geniculate ganglion
Lesion B
Chorda tympani
(through petrotympanic
fissure)
Facial
nerve
Lingual branch of V3
Lesion A
Anterior 2/3 of
tongue
Submandibular
ganglion

▲ ***Figure 26–7. The anatomy of the special sensory component of the facial nerve, comprising the chorda tympani nerve.***


Facial nerve
Oval window niche
Pyramidal
eminence
Tympanic
sinus
Round window niche
Posterior
hypotympanum

▲ ***Figure 26–8. Anatomy of the posterior mesotympanum. Note the location of the facial nerve and ossicular chain.***


Temporal branch
Zygomatic branch
Pinna
Parotid duct
Buccal branch
Branches of facial nerve (CN. VI)
Mastoid process
Parotid gland
Mandibular branch
Masseter muscle
Sternocleidomastoid muscle
Cervical branch

▲ *Figure 26–9. Extratemporal course of facial nerve*


Main trunk of facial nerve
Temporal branch
Zygomatic branch
Buccal branches
Mandibular branch
Cervical branch

▲ *Figure 26–10. A portion of an illustration from Sir Charles Bell, demonstrating the exit of the facial nerve from the stylomastoid foramen.*

Facial Nerve Paralysis

Facial nerve paralysis is classified into:

1. Upper Motor Neuron Lesion

- *Central lesion*
- *Forehead muscles spared*
- *Associated neurological signs present*

2. Lower Motor Neuron Lesion

- *Peripheral lesion*
- *Entire half of the face affected*
- *Forehead involvement present*


Cerebral cortex
Cerebral cortex
Upper motor neuron lesion
Facial nerve nucleus
Lower motor neuron lesion
Facial nerve nucleus
Forehead sparing

▲ *Figure 26–11. Upper and lower motor neuron lesions of facial nerve*


Meatal
Supranuclear
Labyrinthine
Intracranial
LSCC
Processus cochleariformis
Tympanic
Facial nerve nucleus
Oval window
Mastoid
Extratemporal

▲ *Figure 26–12. Course of facial nerve (LSCC: Lateral semicircular canal) Note the important landmarks for facial nerve [oval window, processus cochleariformis and lateral semicircular canal (LSCC)]*


Site of injury
A
B
C
D
Nerve fiber:
Axon
Myelin
Distal to
site of injury
E
F
G
H
Connective tissue:
Endoneurium
Perineurium
Epineurium
Site of Injury:
A - Normal fascicle
B - 1st-degree injury – neuropraxia
C - 2nd-degree injury – axonotmesis
D - 3rd-degree injury
Distal to site of Injury:
E - Normal
F - 1st-degree injury – normal fascicle distally
G - 2nd-degree injury – Wallerian degeneration has produced axonal loss distally
H - 3rd-degree injury

▲ *Figure 26–13. A model of graded neural injury that details clinical–pathological classifications.*


Stimulator
R

▲ *Figure 26–14. Placement of recording and stimulating electrodes for facial EEMG recordings.*

Bell's Palsy

Bell's palsy is an acute, idiopathic, lower motor neuron facial nerve paralysis.

It is the most common cause of facial paralysis.

Etiology

The exact cause is unknown. Proposed mechanisms include:

- *Viral infection, especially herpes simplex virus*
- *Inflammation and edema of the facial nerve*
- *Compression within the facial canal*

Pathophysiology

- *Inflammation causes swelling of the facial nerve*
- *Compression occurs within the narrow facial canal*
- *Ischemia leads to nerve dysfunction*

Clinical Features of Bell's Palsy

Symptoms

- *Sudden onset facial weakness*
- *Drooping of the angle of the mouth*
- *Inability to close the eye*
- *Drooling of saliva*
- *Altered taste sensation*
- *Hyperacusis due to stapedius paralysis*

Signs

- *Loss of forehead wrinkling*
- *Incomplete eye closure*
- *Deviation of mouth to the normal side*
- *Bell's phenomenon, upward rolling of the eyeball on attempted eye closure*

Audiological Relevance

- *Hyperacusis due to stapedius muscle paralysis*
- *Normal hearing thresholds*
- *Reduced acoustic reflex on the affected side*

Clinical Audiology Note

Absent stapedial reflex with normal audiogram suggests facial nerve involvement.

Investigations

- *Diagnosis is mainly clinical*
- *Audiological tests for baseline documentation*
- *Imaging only if atypical features are present*

Treatment of Bell's Palsy

Medical Treatment

- *Oral corticosteroids started early*
- *Antiviral drugs may be added*
- *Eye care is essential:*

– Artificial tears

– Eye patch at night

Physiotherapy

- *Facial muscle exercises*
- *Electrical stimulation in selected cases*

Prognosis

- *Most patients recover completely*
- *Improvement usually begins within weeks*
- *Poor prognosis associated with:*

– Complete paralysis

– Delayed treatment

– Severe pain


Anterior
Tympanic segment
Labyrinthine segment
Medial
Lateral
Canalicular segment
External auditory canal
A
Facial Nerve Decompression: Bell Palsy
25 uV
Electrical Stimulation
Pregeniculate
M5
M6
M7
Distal
M8
100 ms
B

▲ Figure 26–15. (A) Schematic of the approach to the intracanalicular and labyrinthine segments of the facial nerve via a middle cranial fossa. (B) Electrically evokedresponses obtained after decompression of the meatal Foramen in a patient with Bell paralysis and > 90% reduction on preoperative EEMG.



▲ Figure 26–16. (A) Prototypic case of Bell palsy. This 28-year-old woman experienced the onset of an acute, left-sided paralysis of the face over a 24-hour period, in a pattern consistent with peripheral nerve dysfunction. Treatment consisted of oral steroid dosing at pharmacological doses for 10 days, followed by a 2-week taper. (B) Full recovery of facial motor function 2 months after onset.

Differential Diagnosis of Facial Paralysis

- *Ramsay Hunt syndrome*
- *Otitis media related facial palsy*
- *Temporal bone fracture*
- *Acoustic neuroma*
- *Stroke*

Audiology and ENT Red Flags

- *Facial paralysis with ear pain and vesicles*
- *Facial palsy with hearing loss or vertigo*

- ***Recurrent facial palsy***
- ***Progressive facial weakness***

These require urgent ENT evaluation.

Key Points

- ***Facial nerve is a mixed nerve with a long intratemporal course***
- ***Bell's palsy is idiopathic LMN facial paralysis***
- ***Hyperacusis is a key audiologic clue***
- ***Early treatment improves outcome***

Golden Rule

Any facial paralysis with ear symptoms is not Bell's palsy until proven otherwise.

Chapter

27

Hearing Loss

Hearing loss is a reduction in the ability to perceive sound and may affect one or both ears. It can be temporary or permanent and varies in type and degree.

Classification of Hearing Loss

Hearing loss is classified according to:

Site of lesion

Degree of loss

Time of onset

Configuration

I. Classification According to Site of Lesion

1. Conductive Hearing Loss (CHL)

Hearing loss due to impaired transmission of sound through the external or middle ear.

Common causes

- *Wax impaction*
- *Otitis externa*

- *Otitis media*
- *Tympanic membrane perforation*
- *Ossicular fixation or discontinuity*

Audiology

- *Air conduction worse than bone conduction*
- *Air bone gap present*
- *Speech discrimination usually good*

2. Sensorineural Hearing Loss (SNHL)

Hearing loss due to pathology of the cochlea or auditory nerve.

Common causes

- *Noise induced hearing loss*
- *Presbycusis*
- *Meniere disease*
- *Ototoxic drugs*
- *Acoustic neuroma*

Audiology

- *Both air and bone conduction reduced*
- *No air bone gap*
- *Speech discrimination reduced*

3. Mixed Hearing Loss

Combination of conductive and sensorineural components.

Audiology

- *Both air and bone conduction reduced*
- *Air bone gap present*


Middle ear
Inner ear
Outer ear
Conductive
Sensorineural
Mixed

▲ *Figure 27–1.* ***Hearing pathway with type of hearing losses***


Bone conduction (BC)
Air conduction (AC)

▲ *Figure 27–2.* ***Air and Bone conduction pathway Blue arrow–Bone conduction, Red arrow–Air conduction***



Fig. 3.7A: Audiogram



Legends	Right	Left
Air conduction	○	X
Air conduction with masking	△	□
Bone conduction	<	>
Bone conduction with masking	[	]
No response	↙	↘

▲ *Figure 27–3. Legends used in audiogram*



▲ *Figure 27–4. (A) Normal hearing; (B) Conductive hearing loss; (C) Sensorineural hearing loss; (D) Mixed Hearing loss*

II. Classification According to Degree of Hearing Loss

Degree	Hearing Level (dB HL)
Normal	0–25
Mild	26–40
Moderate	41–55
Moderately severe	56–70
Severe	71–90
Profound	> 90



▲ Figure 27–5. Speech discrimination audiogram

III. Classification According to Time of Onset

1. Congenital Hearing Loss

- *Present at birth*
- *Genetic or non genetic*

2. Acquired Hearing Loss

- *Occurs after birth*
- *May be sudden or progressive*

IV. Classification According to Laterality

- *Unilateral*
- *Bilateral*
- *Symmetrical*
- *Asymmetrical*

Audiology Red Flag

Asymmetrical SNHL requires ENT and imaging referral.

V. Configuration of Hearing Loss

- *Flat*
- *Sloping*
- *Rising*
- *Notched*
- *Cookie bite*

Each configuration suggests a different etiology.

Audiological Evaluation

Basic Tests

- ***Pure tone audiometry***
- ***Speech audiometry***
- ***Tympanometry***

Advanced Tests

- ***Otoacoustic emissions***
- ***Auditory brainstem response***

Impact of Hearing Loss

- ***Communication difficulties***
- ***Delayed speech and language development in children***
- ***Academic and social problems***
- ***Reduced quality of life***

Management of Hearing Loss

Management depends on:

- ***Type***
- ***Degree***
- ***Cause***
- ***Age of patient***

Options include:

- ***Medical treatment***
- ***Surgical treatment***
- ***Hearing aids***
- ***Cochlear implantation***
- ***Rehabilitation and counseling***

Audiology Focus Summary

- *Always identify the type of hearing loss*
- *Speech discrimination adds diagnostic value*
- *Asymmetry and sudden loss are red flags*
- *Audiology guides management, not just diagnosis*

Audiology Red Flags

- *Sudden hearing loss*
- *Asymmetric SNHL*
- *Poor speech discrimination out of proportion to PTA*
- *Hearing loss with vertigo or neurological symptoms*

Golden Rule

The first step in hearing loss management is correct classification, not amplification.

Audiology Diagnostic Flowchart

Hearing Loss

Patient complains of hearing difficulty

↓

Step 1: History

- *Onset (sudden or gradual)*
- *Laterality (unilateral or bilateral)*
- *Associated symptoms (pain, discharge, vertigo, tinnitus)*

↓

Step 2: Otoscopic Examination

↓

Is the external ear and TM normal?

↓ ***YES*** ↓ ***NO***

↓ ↓

Step 3: Tympanometry ***Conductive cause likely***

↓ ↓

Middle ear status normal? ***Manage ear pathology***

↓

YES

↓

Step 4: Pure Tone Audiometry

↓

Is there an air bone gap?

↓ ***YES*** ↓ ***NO***

↓ ↓

Conductive or Mixed HL ***Sensorineural HL***

↓ ↓

Step 5: Speech Audiometry ***Step 5: Speech Audiometry***

↓ ↓

Is speech discrimination ***Is speech discrimination***
appropriate for thresholds? ***disproportionately poor?***

↓ ↓

YES ***YES***

↓ ↓

Cochlear pathology likely ***Suspect retrocochlear***

↓ ↓

Routine management ***ABR and MRI referral***

Override Rules

- ***Sudden SNHL goes directly to urgent ENT referral***
- ***Asymmetric SNHL always requires imaging***
- ***Hearing loss with vertigo or neurological signs is an emergency***

Key Audiology Message

Do not jump to amplification before localization.

Diagnosis always comes before devices.

Patient Counseling Sheet

Hearing Loss

What Is Hearing Loss?

Hearing loss means difficulty hearing sounds clearly.

It can affect one ear or both ears and may be temporary or permanent.

Types of Hearing Loss

- ***Conductive hearing loss***

Problem in the outer or middle ear. Often treatable.

- ***Sensorineural hearing loss***

Problem in the inner ear or hearing nerve. Usually permanent.

- ***Mixed hearing loss***

Combination of both types.

Common Symptoms

- *Difficulty hearing speech*
- *Needing higher volume on TV or phone*
- *Ringing in the ears, tinnitus*
- *Difficulty hearing in noise*
- *Asking others to repeat*

Important Facts for Patients

- *Hearing loss is common and manageable*
- *Early diagnosis improves outcomes*
- *Hearing aids do not make hearing worse*
- *Ignoring hearing loss affects communication and quality of life*

What Tests Are Needed?

Audiology tests may include:

- *Hearing test, audiogram*
- *Speech understanding test*
- *Middle ear test*
- *Special tests if needed*

These tests help identify the type and cause of hearing loss.

Treatment Options

Treatment depends on the type and degree of hearing loss:

- *Medical treatment*
- *Surgical treatment*
- *Hearing aids*

- ***Cochlear implant in selected cases***
- ***Hearing rehabilitation and counseling***

Daily Living Advice

- ***Reduce background noise when talking***
- ***Face the speaker***
- ***Ask people to speak clearly, not loudly***
- ***Use hearing devices as advised***
- ***Attend regular hearing follow up***

When to Seek Medical Help Urgently

- ***Sudden hearing loss***
- ***Hearing loss with dizziness***
- ***Hearing loss with severe pain***
- ***Hearing loss in one ear only***

Prognosis

- ***Some hearing loss is reversible***
- ***Some types are permanent but manageable***
- ***Regular follow up helps preserve hearing and communication***

Key Message for Patients

Hearing loss is not just about ears,

it affects communication, safety, and quality of life.

Early testing and follow up make a real difference.

Audiology Diagnostic Report

Hearing Loss, Flowchart-Linked Template

Reason for Referral

Audiologic evaluation for reported hearing difficulty.

Step 1. History Summary

(as per diagnostic flowchart)

Onset

☐ Sudden ☐ Gradual

Laterality

☐ Right ☐ Left ☐ Bilateral ☐ Asymmetrical

Associated symptoms

☐ Tinnitus

☐ Vertigo or imbalance

☐ Ear pain

☐ Ear discharge

☐ Neurological symptoms

Relevant history notes:

Step 2. Otoscopic Examination

External auditory canal

☐ Normal ☐ Abnormal

Tympanic membrane

☐ Intact ☐ Perforated ☐ Congested ☐ Retracted

Otoscopic impression:

Step 3. Immittance Findings

Tympanometry

☐ Type A ☐ Type As ☐ Type Ad ☐ Type B ☐ Type C

Acoustic reflexes

☐ Present ☐ Absent ☐ Not tested

Middle ear status:

☐ Normal

☐ Abnormal, consistent with conductive component

Step 4. Pure Tone Audiometry

Right ear

☐ Normal ☐ Conductive ☐ Sensorineural ☐ Mixed

Left ear

☐ Normal ☐ Conductive ☐ Sensorineural ☐ Mixed

Degree of hearing loss

☐ Mild ☐ Moderate ☐ Moderately severe ☐ Severe ☐ Profound

Air bone gap

☐ Present ☐ Absent

Step 5. Speech Audiometry

Speech discrimination score

Right ear ______ %

Left ear ______ %

Speech discrimination

- ☐ *Consistent with PTA*
- ☐ *Disproportionately poor*

Step 6. Diagnostic Flowchart Conclusion

Based on history, otoscopy, immittance, pure tone, and speech audiometry:

- ☐ *Conductive hearing loss*
- ☐ *Sensorineural hearing loss, cochlear pattern*
- ☐ *Sensorineural hearing loss, retrocochlear pattern suspected*
- ☐ *Mixed hearing loss*

Red Flag Assessment

- ☐ *Sudden SNHL*
- ☐ *Asymmetrical SNHL*
- ☐ *Poor speech discrimination not matching PTA*
- ☐ *Hearing loss with vertigo*
- ☐ *Neurological symptoms*

If any box is checked, escalation is required.

Clinical Impression

Audiologic findings indicate ___________________ hearing loss, consistent with ___________________ pathology.

Findings are interpreted according to the institutional diagnostic flowchart for hearing loss.

Recommendations

☐ Routine audiologic follow up

☐ Medical or surgical ENT evaluation

☐ Hearing aid evaluation

☐ ABR testing

☐ MRI referral

☐ Urgent ENT referral due to red flag findings

Testing Limitations

Audiologic testing was limited by:

☐ Active ear disease

☐ Patient tolerance

☐ Safety considerations

Key Documentation Rule

Each conclusion in the report must be traceable to a step in the diagnostic flowchart.

No conclusion without data. No data without documentation.

Auditory Neuropathy Spectrum Disorder (ANSD)

***Auditory Neuropathy Spectrum Disorder (ANSD)* is a hearing disorder characterized by *impaired neural transmission of auditory signals*, despite *preserved cochlear outer hair cell function*.**

Children with ANSD may have:

- ***Normal or near-normal otoacoustic emissions***
- ***Absent or severely abnormal auditory brainstem responses***
- ***Disproportionately poor speech perception***

ANSD represents a spectrum, not a single disease entity.

Pathophysiology

ANSD results from dysfunction at one or more levels of the auditory pathway:

- ***Inner hair cells***
- ***Synapse between inner hair cells and auditory nerve***
- ***Auditory nerve fibers***

This leads to temporal dys-synchrony, which severely affects speech understanding, especially in noise.

Etiology

Common Causes

- ***Prematurity***
- ***Hyperbilirubinemia***
- ***Hypoxia***

- ***NICU stay***
- ***Genetic mutations***
- ***Auditory nerve deficiency***

ANSD may be:

- ***Congenital***
- ***Acquired***
- ***Transient or permanent***

Audiologic Characteristics

Typical audiologic findings include:

- ***Present otoacoustic emissions in early stages***
- ***Absent or grossly abnormal ABR***
- ***Variable pure tone thresholds***
- ***Disproportionately poor speech perception***
- ***Absent acoustic reflexes***

Key Point

Pure tone thresholds alone do not reflect functional hearing ability in ANSD.

Diagnosis of ANSD

Diagnosis is based on a test battery, not a single test.

Required Findings

- *Absent or abnormal ABR*
- *Preserved OAEs or cochlear microphonic*
- *Poor speech perception relative to audiogram*

Diagnosis must exclude:

- *Middle ear pathology*
- *Technical recording errors*

Pediatric Considerations

- *ANSD is frequently identified during newborn hearing screening*
- *AABR is critical for detection*
- *OAE-only screening will miss ANSD*

Medico-Legal Risk

Relying solely on OAE screening in high-risk infants.

Natural History

- *Some infants show improvement over time*
- *Others demonstrate stable or progressive deficits*
- *OAEs may disappear with age*
- *Speech and language outcomes are highly variable*

Regular reassessment is mandatory.

Management Principles

There is no single management strategy suitable for all ANSD cases.

Management depends on:

- *Functional auditory performance*

- ***Speech and language development***
- ***Benefit from amplification***
- ***Family and educational factors***

Role of Hearing Aids

- ***May provide benefit in selected cases***
- ***Benefit is unpredictable***
- ***Must be assessed functionally, not assumed***

Trial amplification requires close monitoring.

Cochlear Implantation in ANSD

- ***CI can improve neural synchrony***
- ***Particularly beneficial when lesion is pre-synaptic***
- ***Outcomes vary depending on neural integrity***

CI candidacy is based on:

- ***Poor functional hearing***
- ***Limited benefit from amplification***
- ***Delayed speech and language development***

Role of Audiology

Audiology is responsible for:

- ***Accurate diagnosis using test battery***
- ***Monitoring auditory function over time***
- ***Functional assessment of hearing ability***
- ***Counseling families***
- ***Coordinating referrals***

Audiology does not predict outcomes with certainty.

Counseling of Families

Families must understand:

- ANSD is variable and unpredictable
- Audiograms may not reflect communication ability
- Progress requires close follow-up
- Outcomes cannot be guaranteed

Clear documentation of counseling is essential.

Educational Considerations

- Classroom listening may be severely affected
- Assistive listening devices may be required
- Visual communication strategies may be beneficial

Medico-Legal Considerations

High-risk areas include:

- Failure to detect ANSD in newborn screening
- Over-reliance on OAEs
- Delayed referral for CI evaluation
- Inadequate documentation of functional performance

Key Points

- ANSD is a disorder of neural synchrony
- OAEs may be normal

Unilateral Hearing Loss

UHL

Unilateral hearing loss is hearing loss affecting ***one ear only****, with the other ear having normal or significantly better hearing.*

It may be ***conductive, sensorineural, or mixed****.*

Clinical Importance

Unilateral hearing loss is ***never benign until evaluated****.*

It requires careful assessment to exclude ***retrocochlear and central causes****.*

Classification of Unilateral Hearing Loss

According to Type

Unilateral Conductive Hearing Loss

Unilateral Sensorineural Hearing Loss

Unilateral Mixed Hearing Loss

According to Onset

- *Sudden*
- *Gradual*

Sudden unilateral SNHL is treated as ***SSNHL****.*

Common Causes

I. Unilateral Conductive Hearing Loss

- ***Impacted cerumen***
- ***Otitis externa***
- ***Otitis media***
- ***Tympanic membrane perforation***
- ***Otosclerosis***
- ***Ossicular discontinuity***

II. Unilateral Sensorineural Hearing Loss

- ***Sudden sensorineural hearing loss***
- ***Acoustic neuroma (vestibular schwannoma)***
- ***Meniere disease***
- ***Noise induced hearing loss***
- ***Viral or vascular causes***

Acoustic neuroma must always be excluded.

III. Unilateral Mixed Hearing Loss

- ***Chronic otitis media***
- ***Cholesteatoma***
- ***Otosclerosis with cochlear involvement***

Audiological Evaluation

History

- ***Onset and duration***
- ***Sudden or progressive***
- ***Associated tinnitus***

- *Vertigo or imbalance*
- *Neurological symptoms*

Pure Tone Audiometry

- *Confirm type and degree of loss*
- *Identify air bone gap*
- *Document asymmetry*

Speech Audiometry

- *Speech discrimination score*
- *Disproportionately poor score suggests retrocochlear pathology*

Tympanometry

- *To exclude middle ear pathology*
- *Usually normal in SNHL*

Definition of Asymmetry

Unilateral hearing loss or asymmetry is suspected when:

- *≥ 15 dB difference at two consecutive frequencies*
- *≥ 20 dB difference at one frequency*
- *Speech discrimination difference > 15–20 percent*

Red Flags in Unilateral Hearing Loss

Immediate ENT referral if:

- *Sudden onset unilateral SNHL*
- *Unilateral tinnitus*
- *Unilateral hearing loss with vertigo*
- *Poor speech discrimination*
- *Progressive asymmetry*

Audiology Role

Audiology must:

- ***Confirm and document asymmetry***
- ***Identify SNHL vs CHL***
- ***Trigger ENT referral***
- ***Avoid reassurance without exclusion of serious causes***

Audiology must not:

- ***Attribute unilateral SNHL to aging***
- ***Delay referral for imaging***

ENT Role

ENT is responsible for:

- ***Medical diagnosis***
- ***Imaging, usually MRI***
- ***Medical or surgical management***

Management Overview

Conductive Causes

- ***Medical or surgical treatment***
- ***Hearing usually improves after correction***

Sensorineural Causes

- ***SSNHL requires urgent steroids***
- ***Retrocochlear causes require imaging***
- ***Rehabilitation if permanent loss remains***

Rehabilitation Options

- *Hearing aid in affected ear*
- *CROS or BiCROS systems*
- *Bone conduction devices*
- *Cochlear implant in selected cases*

Medico-Legal Considerations

- *Failure to investigate unilateral SNHL is a common source of litigation*
- *Audiology reports must document:*

– Degree of asymmetry
– Referral recommendation

Safe wording:

"Unilateral sensorineural hearing loss was identified. ENT evaluation and imaging are recommended to exclude retrocochlear pathology."

Meningitis

Meningitis is *inflammation of the meninges* surrounding the brain and spinal cord, most commonly caused by *infection*.

Meningitis is a *life threatening medical emergency*.

Classification of Meningitis

According to Etiology

Bacterial meningitis

Viral meningitis

Tuberculous meningitis

Fungal meningitis

According to Clinical Course

- Acute
- Subacute
- Chronic

Clinical Presentation

Classic Symptoms

- Fever
- Headache
- Neck stiffness
- Altered consciousness

Additional Features

- *Vomiting*
- *Photophobia*
- *Seizures*
- *Rash, in some bacterial types*

Infants may present with nonspecific symptoms.

Otologic and Audiologic Importance

Meningitis is one of the most important causes of acquired sensorineural hearing loss, especially in children.

Hearing Loss in Meningitis

Type

- ***Usually sensorineural hearing loss***
- ***Often bilateral***
- ***May be sudden or progressive***

Mechanism

- ***Spread of infection to the inner ear***
- ***Inflammatory damage to cochlea***
- ***Labyrinthitis***
- ***Cochlear ossification after infection***

Audiology Role in Meningitis

Audiology must:

- ***Identify hearing loss early***
- ***Document baseline hearing after stabilization***
- ***Monitor progression***
- ***Guide rehabilitation***

Audiology involvement is mandatory after recovery from meningitis.

Timing of Hearing Assessment

Recommended

- As soon as patient is medically stable
- Within 4 weeks after recovery
- Earlier in children if possible

Audiologic Evaluation

- Pure tone audiometry
- Speech audiometry
- Tympanometry
- Otoacoustic emissions in children
- ABR if behavioral testing not possible

Red Flags in Post Meningitis Hearing

Immediate ENT referral if:

- Sudden deterioration in hearing
- Asymmetric hearing loss
- Vertigo
- Facial nerve involvement

Imaging Considerations

- CT or MRI may show inner ear involvement
- Imaging is critical before cochlear implantation to assess ossification

Rehabilitation After Meningitis

Mild to Moderate SNHL

- Hearing aids
- Speech therapy

Severe to Profound SNHL

- Cochlear implant evaluation
- Early implantation is crucial before cochlear ossification

Delay may make implantation difficult or impossible.

Pediatric Considerations

- High risk of permanent hearing loss
- Early screening is essential
- Family counseling mandatory

Medico Legal Considerations

- Failure to assess hearing after meningitis is a recognized medico-legal risk
- Audiology must document:

– Date of meningitis diagnosis
– Date of hearing assessment
– Follow-up plan

Safe wording:

"Hearing assessment was performed following recovery from meningitis to evaluate possible inner ear involvement."

Occupational Hearing Loss

Noise-Induced Hearing Loss in the Workplace

Occupational Hearing Loss (OHL) refers primarily to noise-induced sensorineural hearing loss resulting from chronic exposure to hazardous noise levels in the workplace.

It is one of the most common and most preventable occupational diseases worldwide

Epidemiology and Importance

- ***Millions of workers are affected by OHL***
- ***Prevalence increases with duration and intensity of exposure***
- ***OHL has major human, economic, and medico-legal consequences***
- ***Despite this, it is often under-recognized because it does not visibly affect work performance***

Pathophysiology

Occupational noise exposure leads to:

- ***Damage to cochlear outer hair cells***
- ***Progressive loss of inner hair cell function***
- ***Degeneration of auditory nerve fibers***

Initial injury typically affects frequencies between 3000–6000 Hz, classically around 4000 Hz, but this pattern is not specific to noise alone

Audiologic Characteristics

Typical audiologic findings include:

- ***Bilateral, usually symmetrical SNHL***
- ***High-frequency notch, often at 4 kHz***
- ***Gradual progression with continued exposure***
- ***Relative stability after removal from noise exposure***

Important

A 4-kHz notch alone does not confirm OHL.



▲ Figure 27–6. Noise-induced hearing loss. The audiogram shows a typical bilateral high-frequency sensorineural hearing loss, most severe at 4000 Hz, with a normal speech discrimination score.



▲ Figure 27–7. Ear plugs and muffs

Diagnostic Criteria for OHL

A diagnosis of occupational hearing loss must be based on positive evidence, including:

- *Documented exposure to hazardous noise*
- *Audiogram consistent with noise-induced patterns*
- *Appropriate temporal relationship between exposure and loss*
- *Absence of alternative causes of hearing loss*
- *Stability of hearing after noise exposure ends*

OHL is a diagnosis of exclusion, not assumption.

Differential Diagnosis

Conditions that may mimic OHL include:

- *Presbycusis*
- *Ototoxicity*
- *Genetic hearing loss*
- *Otosclerosis*
- *Retrocochlear pathology*

Failure to consider these alternatives is a medico-legal error

Audiology Evaluation Protocol

Audiologic assessment should include:

- *Pure tone audiometry (air and bone)*
- *Speech audiometry*
- *Serial audiograms when available*
- *Review of exposure history*

Audiologists must document inconsistencies and recommend medical evaluation when findings are atypical.

Hearing Conservation Programs

Effective prevention includes:

- *Noise surveys to identify hazardous exposure*
- *Regular audiometric monitoring*
- *Properly fitted hearing protection*
- *Training and retraining of personnel*
- *Medical evaluation of abnormal audiograms*

Hearing conservation programs are inexpensive and highly effective

Medico-Legal Considerations

Key Principles

- *Employment in a noisy environment does not automatically equal OHL*
- *Diagnosis must be evidence-based*
- *Large compensation claims increase false attribution risk*
- *Clinicians must remain objective and independent*

Common Medico-Legal Pitfalls

- *Labeling hearing loss as occupational without full evaluation*
- *Ignoring asymmetry or sudden changes*
- *Inadequate documentation of exposure history*
- *Failure to recommend medical referral*

Audiology Ethical Responsibilities

Audiologists must:

- *Maintain diagnostic neutrality*

- *Avoid advocacy bias*
- *Separate clinical findings from compensation issues*
- *Clearly document limitations and uncertainty*

Management

Primary Prevention

- *Noise reduction when possible*
- *Hearing protection use*

Secondary Prevention

- *Early detection through surveillance*
- *Removal from hazardous exposure when indicated*

Rehabilitation

- *Hearing aids when appropriate*
- *Counseling on realistic expectations*

Key Points

- *OHL is common but preventable*
- *Diagnosis requires positive evidence*
- *Audiogram patterns are suggestive, not definitive*
- *Medico-legal implications are significant*

Golden Rule

Noise exposure explains hearing loss
only after other causes are excluded.

Sudden Sensorineural Hearing Loss SSNHL

Sudden sensorineural hearing loss is defined as a rapid onset sensorineural hearing loss of at least 30 dB affecting three consecutive frequencies, occurring within 72 hours or less.

SSNHL is a medical emergency.

Epidemiology

- *Can occur at any age*
- *More common in adults*
- *Usually unilateral*
- *Bilateral SSNHL is rare and more serious*

Etiology

In most cases, the cause is idiopathic.

Proposed Causes

- *Viral infection*
- *Vascular compromise*
- *Autoimmune inner ear disease*
- *Membrane rupture*
- *Retrocochlear pathology*

Clinical Presentation

Key Features

- *Sudden hearing loss, often noticed on waking*
- *Usually unilateral*

- *Ear fullness*
- *Tinnitus common*
- *Vertigo may be present*

Presence of vertigo indicates worse prognosis.

Audiological Findings

Pure Tone Audiometry

- *Sensorineural hearing loss*
- *Any configuration possible*
- *Low frequency loss has better prognosis*

Speech Audiometry

- *Reduced speech discrimination*
- *May be disproportionate to PTA*

Tympanometry

- *Usually Type A*
- *Helps exclude middle ear pathology*

Diagnosis

Diagnosis is clinical and audiometric.

Criteria:

- *Sudden onset*
- *SNHL pattern*
- *No conductive component*

Audiology confirms the loss.

ENT initiates treatment.

Audiology Role in SSNHL

Audiology must:

- ***Perform urgent baseline audiogram***
- ***Confirm sensorineural nature***
- ***Document severity and configuration***
- ***Identify asymmetry***
- ***Trigger immediate ENT referral***

Audiology must not delay referral for additional tests.

Management Principles

ENT Management

- ***Systemic steroids***
- ***Intratympanic steroids when indicated***
- ***Imaging to exclude retrocochlear pathology***

Timing is Critical

Best outcomes occur when treatment starts:

- ***Within 72 hours***
- ***Ideally within the first 7 days***

Prognostic Factors

Better Prognosis

- ***Early treatment***
- ***Mild to moderate loss***
- ***Low frequency loss***
- ***Absence of vertigo***

Poor Prognosis

- ***Delayed treatment***
- ***Severe or profound loss***
- ***Presence of vertigo***
- ***Advanced age***

Follow Up Audiology

Initial Follow Up

- ***Repeat audiogram after treatment course***

Ongoing Follow Up

- ***Monitor recovery or stabilization***
- ***Compare to baseline***

Rehabilitation After SSNHL

If incomplete recovery:

- ***Hearing aid fitting***
- ***Tinnitus management***
- ***Counseling***

Red Flags in SSNHL

Immediate ENT referral if:

- ***Any sudden SNHL***
- ***SSNHL with vertigo***
- ***Bilateral SSNHL***
- ***SSNHL with neurological symptoms***

Medico Legal Considerations

- ***Delay in diagnosis or referral increases liability***
- ***Audiology must document:***

– Time of onset

– Date of testing

– Urgency of referral

Safe wording:

"Findings are consistent with sudden sensorineural hearing loss.

Urgent ENT evaluation was advised on the same day."

Key Points

- *SSNHL is an emergency*
- *Audiology confirms, ENT treats*
- *Early steroids improve outcomes*
- *Delay worsens prognosis*

Golden Rule

Sudden hearing loss is SSNHL until proven otherwise.

Test fast. Refer immediately.

SSNHL Audiology Emergency Flowchart

Patient reports sudden hearing loss

↓

Step 1 Confirm urgency

- Onset within 72 hours
- Usually unilateral

↓

Step 2 Immediate audiology

- Pure tone audiometry
- Bone conduction
- Speech audiometry if possible
- Tympanometry to exclude CHL

↓

Is sensorineural hearing loss confirmed?

↓ NO ↓ YES

↓ ↓

Evaluate other causes SSNHL confirmed

↓

Step 3 Red flag screening

- Vertigo
- Neurological symptoms
- Bilateral loss

↓

Step 4 Immediate action

- *Same day ENT referral*
- *Document time of onset and testing*

↓

Step 5 Follow up

- *Repeat audiogram after treatment*
- *Compare to baseline*

Critical Rule

Audiology must not delay ENT referral for additional testing.

Key Safety Message

SSNHL is diagnosed in minutes, not days.

Referral must be same day.

ENT Referral Template

Sudden Sensorineural Hearing Loss (SSNHL)

Date: ____________________

Urgency: SAME DAY

To: ENT Consultant

From: Audiology Department

Patient Information

Name: ____________________

Age: ____________________

MRN: ____________________

Reason for Referral

Urgent ENT evaluation for sudden sensorineural hearing loss.

History Summary

- Onset of hearing loss: ____________________
- Duration since onset: ____________________
- Laterality: ☐ Right ☐ Left ☐ Bilateral
- Associated symptoms

☐ Tinnitus

☐ Vertigo

☐ Aural fullness

☐ Neurological symptoms

Audiologic Findings

- *Pure tone audiometry shows SNHL*
- *Degree: ☐ Mild ☐ Moderate ☐ Severe ☐ Profound*
- *Configuration: ____________________*
- *Speech discrimination: ____________________*
- *Tympanometry: ☐ Type A ☐ Other _________*

Clinical Impression

Audiologic findings are consistent with SSNHL.

Urgent ENT management is advised.

Recommendation

- *Immediate ENT assessment*
- *Steroid therapy as indicated*
- *Imaging to exclude retrocochlear pathology*

Medico Legal Statement

Audiologic assessment confirms sudden sensorineural hearing loss based on patient history and testing performed on the date above.

Urgent ENT referral was advised without delay.

Referral Rule

Every SSNHL referral must carry a time stamp.

Chapter

28

Tinnitus

Tinnitus is the perception of sound in the absence of an external acoustic stimulus.

It is a symptom, not a disease.

Types of Tinnitus

I. Subjective Tinnitus

- ***Heard only by the patient***
- ***Most common type***
- ***Usually related to auditory system pathology***

II. Objective Tinnitus

- ***Rare***
- ***Can be heard by the examiner***
- ***Usually vascular or muscular in origin***

Classification of Tinnitus

According to Sound Quality

- ***Ringing***
- ***Buzzing***

- ***Hissing***
- ***Roaring***
- ***Clicking***

According to Laterality

- ***Unilateral***
- ***Bilateral***

Red Flag

Unilateral tinnitus requires further evaluation.

According to Rhythm

- ***Non pulsatile***
- ***Pulsatile***

Pulsatile tinnitus is vascular until proven otherwise.

Causes of Tinnitus

Cochlear Causes

- ***Noise induced hearing loss***
- ***Presbycusis***
- ***Meniere disease***
- ***Ototoxic drugs***

Retrocochlear Causes

- ***Acoustic neuroma***
- ***Brainstem lesions***

Middle Ear Causes

- ***Otosclerosis***
- ***Middle ear effusion***
- ***Glomus tumors***

Vascular Causes

- ***Glomus tumor***
- ***High jugular bulb***
- ***AV malformations***

Muscular Causes

- ***Palatal myoclonus***
- ***Middle ear muscle spasm***

Pathophysiology

- ***Abnormal neural activity in the auditory pathway***
- ***Reduced auditory input leads to central gain increase***
- ***Brain generates phantom sound perception***

Key Concept

Tinnitus is mainly a central auditory processing phenomenon, even when triggered by peripheral pathology.

Clinical Evaluation

History

- ***Onset and duration***
- ***Unilateral or bilateral***
- ***Pulsatile or non pulsatile***
- ***Associated hearing loss***

- *Vertigo or imbalance*
- *Exposure to noise or drugs*

Otoscopic Examination

- *Usually normal*
- *Look for vascular middle ear masses*

Audiological Evaluation

Pure Tone Audiometry

- *Often shows SNHL*
- *Degree of hearing loss does not correlate with tinnitus severity*

Speech Audiometry

- *Helps identify cochlear vs retrocochlear pattern*

Immittance

- *To exclude middle ear pathology*

Special Tests

- *ABR if retrocochlear pathology suspected*
- *Imaging when red flags are present*

Red Flags in Tinnitus

Immediate ENT referral if:

- *Unilateral tinnitus*
- *Pulsatile tinnitus*
- *Tinnitus with asymmetric SNHL*
- *Tinnitus with vertigo*
- *Tinnitus with neurological symptoms*

Management of Tinnitus

Treat Underlying Cause

- Wax removal
- Treat middle ear disease
- Manage hearing loss

Audiologic Management

- Hearing aids
- Sound therapy
- Tinnitus masking
- Tinnitus retraining therapy

Counseling

- Explanation and reassurance
- Stress management
- Sleep hygiene

Prognosis

- Tinnitus may persist
- Severity often decreases with counseling and sound therapy
- Habituation is the goal, not silence

Audiology Focus Summary

- Tinnitus is a symptom, not a diagnosis
- Severity does not equal pathology severity
- Red flags guide imaging and referral
- Audiology plays a central role in management

Golden Rule

Unilateral or pulsatile tinnitus is pathological until proven otherwise.

Audiology Diagnostic Flowchart

Tinnitus

Patient complains of tinnitus

↓

Step 1 History

- ***Unilateral or bilateral***
- ***Pulsatile or non pulsatile***
- ***Sudden or gradual onset***
- ***Associated hearing loss, vertigo, neurological symptoms***

↓

Step 2 Otoscopic examination

↓

Is there visible middle ear pathology or vascular mass

↓ ***YES*** ↓ ***NO***

↓ ↓

ENT referral ***Step 3 Audiology tests***

- ***PTA***
- ***Speech audiometry***
- ***Tympanometry***

↓

Step 4 Hearing status

↓

Is hearing loss present

↓ ***YES*** ↓ ***NO***

↓ ↓

Manage hearing loss ***Consider central tinnitus***

and tinnitus together ***counseling and sound therapy***

↓

Step 5 Red flag assessment

- ***Unilateral tinnitus***
- ***Pulsatile tinnitus***
- ***Asymmetric SNHL***
- ***Poor speech discrimination***

↓

Any red flag present

↓ ***YES*** ↓ ***NO***

↓ ↓

ENT referral with ***Audiology based management***

imaging recommendation ***and follow up***

Override Rules

- ***Pulsatile tinnitus bypasses routine testing and goes to ENT***
- ***Unilateral tinnitus with asymmetry requires imaging***
- ***Tinnitus with vertigo or neurological signs is urgent***


Tinnitus
Annoying
Not troublesome transient
Organic
Functional/psychogenic
Peripheral
Central
• Temporal lobe epilepsy (auditory hallucination)
• Cerebrovascular disease
• Intracranial vascular tumors
• Migraine
Systemic
• Anemia
• Strychnine poisoning
• Hyper-/hypotension
In the ear
Outside the ear
• ET dysfunction
• Palatal myoclonus
• Impacted wisdom tooth
External ear
• Wax
• Foreign body
Middle ear
• Otosclerosis
• Fluid in the middle ear
• Glomus tumor
• AV malformation
Inner ear
• Meniere's disease
• Hair cell damage
• Ototoxicity
• Acoustic neuroma
• Noise trauma

▲ *Flowchart 28–1. Classification of tinnitus*

Patient Counseling Sheet

Tinnitus

What Is Tinnitus

Tinnitus is the sensation of hearing sound when no external sound is present.

It may sound like ringing, buzzing, hissing, or roaring.

Important Facts for Patients

- ***Tinnitus is a symptom, not a disease***
- ***It is very common***
- ***It does not mean you are going deaf***

- *It is rarely dangerous*
- *Stress and anxiety can make it worse*

Why Does Tinnitus Happen

Common reasons include:

- *Hearing loss*
- *Noise exposure*
- *Inner ear disorders*
- *Stress and fatigue*

Sometimes no single cause is found.

What Tests Are Needed

Audiology tests help to:

- *Check hearing levels*
- *Identify ear related causes*
- *Decide if ENT referral is needed*

Imaging is only required when warning signs are present.

Lifestyle Advice

Hearing Protection

- *Avoid loud noise*
- *Use ear protection when needed*

Sound Environment

- *Avoid complete silence*
- *Use background sound at night*
- *Low level music or white noise may help*

Stress and Sleep

- *Manage stress*
- *Practice relaxation techniques*
- *Maintain good sleep routine*

Treatment Expectations

- *There is no single cure*
- *Treatment aims to reduce awareness and distress*
- *Hearing aids often reduce tinnitus*
- *Sound therapy and counseling are effective*

When to Seek Medical Advice Urgently

Contact your doctor if you have:

- *Pulsatile tinnitus*
- *Tinnitus in one ear only*
- *Tinnitus with hearing loss or dizziness*
- *Sudden change in hearing*

Prognosis

- *Many patients improve over time*
- *Habituation is common*
- *Quality of life improves with proper management*

Key Message for Patients

Tinnitus is manageable.

Understanding it reduces fear,

and reduced fear reduces tinnitus impact.

Chapter

29

Earache



Otalgia

Otalgia is pain felt in or around the ear.

It is a symptom, not a diagnosis.

Classification of Otalgia

Otalgia is classified into two main types:

Primary (Otogenic) Otalgia

Secondary (Referred) Otalgia

This classification is clinically very important.

I. Primary (Otogenic) Otalgia

Pain originates from ear pathology itself.

Common Causes

External Ear Causes

- Otitis externa
- Furunculosis
- Impacted wax
- Foreign body
- Trauma to external auditory canal

Middle Ear Causes

- *Acute otitis media*
- *Otitis media with effusion, dull ache*
- *Barotrauma*
- *Acute mastoiditis*

Inner Ear Causes

- *Inner ear diseases rarely cause pain*
- *Severe labyrinthitis may cause discomfort but true otalgia is uncommon*

Clinical Features of Primary Otalgia

- *Ear pain often severe*
- *Pain increases with ear manipulation, especially in otitis externa*
- *Otoscopic examination usually reveals pathology*

II. Secondary (Referred) Otalgia

Pain is felt in the ear but originates outside the ear.

This occurs because the ear shares nerve supply with other structures.

Nerve Supply Involved

- *Trigeminal nerve (V)*
- *Facial nerve (VII)*
- *Glossopharyngeal nerve (IX)*
- *Vagus nerve (X)*
- *Upper cervical nerves (C2, C3)*

Common Causes of Referred Otalgia

Dental Causes

- *Dental caries*
- *Impacted molar*
- *Dental abscess*

Throat and Pharyngeal Causes

- *Acute tonsillitis*
- *Pharyngitis*
- *Peritonsillar abscess*
- *Tumors of the tonsil or base of tongue*

Laryngeal Causes

- *Laryngeal tumors*
- *Chronic laryngitis*

Temporomandibular Joint (TMJ)

- *TMJ arthritis*
- *Malocclusion*
- *Bruxism*

Cervical Spine

- *Cervical spondylosis*
- *Neck muscle spasm*

Clinical Features of Referred Otalgia

- *Otoscopic examination is normal*
- *Pain is often vague or dull*
- *Pain may be aggravated by swallowing, chewing, or neck movement*


Larynx
Vagus nerve supplying to external ear, larynx and laryngopharynx
Oropharynx
C2, 3
Glossopharyngeal nerve supplying to middle ear and oropharynx
CN V
Teeth and gums
Cervical vertebra
Temporo-mandibular joint
Base
Tongue
Anterior 2/3

▲ *Figure 29–1 Referred otalgia. Common causes of referred otalgia and nerve supply of ear*

Red Flag

Referred otalgia in adults may be the only symptom of head and neck malignancy.

Immediate ENT referral if:

- ***Severe otalgia with normal ear examination***
- ***Persistent unilateral ear pain in adults***
- ***Otalgia associated with weight loss or dysphagia***
- ***Otalgia with cranial nerve symptoms***


Cervical spondylosis, caries spine
C2, C3
via 5th nerve
via 9th nerve
via 10th nerve
• Temporomandibular joint disorder
• Impacted tooth, dental caries
• Acute tonsillitis, peritonsillar abscess
• Eagle's syndrome (styalgia)
• Oropharyngeal malignancy

▲ *Figure 29–2.Diagrammatic represenation of referred pain to the ear*

Key Points

- ***Otalgia is a symptom, not a disease***
- ***Normal otoscopy suggests referred pain***
- ***Audiology plays a role in exclusion and referral***

Golden Rule

Ear pain with a normal ear must be investigated beyond the ear.

Chapter

30

Otorrhea

Ear Discharge

Otorrhea is discharge coming from the external auditory canal.

It is a sign, not a diagnosis.

Classification of Otorrhea

Otorrhea can be classified according to:

- ***Site of origin***
- ***Nature of discharge***
- ***Duration***

I. Classification According to Site of Origin

1. External Ear Causes

• Otitis externa

– Purulent discharge

– Painful ear

– Tender tragus

- ***Furunculosis***

– *Localized painful swelling*

– *Thick purulent discharge after rupture*

- ***Foreign body***

– *Offensive discharge, especially in children*

- ***Trauma to EAC***

– *Blood-stained discharge*

2. Middle Ear Causes

- ***Acute otitis media with perforation***

– *Mucopurulent discharge*

– *Relief of pain after discharge appears*

- ***Chronic suppurative otitis media (CSOM)***

– *Persistent or recurrent discharge*

– *Central perforation, safe type*

– *Attic or marginal perforation, unsafe type*

- ***Otitis media with effusion after grommet insertion***

3. Inner Ear and Other Causes

- ***CSF otorrhea***

– *Clear, watery discharge*

– *Following trauma or surgery*

– *Serious emergency*

II. Classification According to Nature of Discharge

1. Serous

- *Clear, watery*
- *Seen in early AOM or CSF leak*

2. Mucoid

- *Thick, sticky*
- *Common in chronic otitis media*

3. Purulent

- *Yellow or green*
- *Suggests bacterial infection*

4. Blood-stained

- *Trauma*
- *Tumors*
- *Acute severe infection*

5. Foul-smelling

Highly suggestive of cholesteatoma

III. Classification According to Duration

Acute Otorrhea

- *Sudden onset*
- *Usually painful*
- *Common in AOM and otitis externa*

Chronic Otorrhea

- *Lasts more than 6 weeks*
- *Usually painless*
- *Common in CSOM*

Otoscopic Examination

- *Condition of external canal*
- *Tympanic membrane status*
- *Presence of perforation*
- *Granulation tissue or polyp*
- *Cholesteatoma debris*

Important rule

Never clean or probe the ear blindly before full examination.

Audiological Relevance

- *Otorrhea often causes conductive hearing loss*
- *Audiologic testing may be limited during active discharge*
- *Tympanometry usually unreliable*
- *Hearing should be reassessed after ear becomes dry*

Red Flags in Otorrhea

Immediate ENT referral if:

- *Foul-smelling discharge*
- *Attic or marginal perforation*
- *Otorrhea with vertigo*
- *Otorrhea with facial nerve weakness*
- *Clear watery discharge suspected CSF*

Management Principles

General Measures

- ***Keep ear dry***
- ***Avoid ear syringing***
- ***Aural toilet by trained personnel***

Medical Treatment

- ***Topical antibiotics***
- ***Systemic antibiotics when indicated***
- ***Avoid ototoxic drops if TM is perforated***

Surgical Treatment

- ***Indicated in unsafe CSOM***
- ***Mastoid surgery for cholesteatoma***

Audiology Focus Summary

- ***Otorrhea indicates active ear disease***
- ***Foul smell suggests unsafe pathology***
- ***Audiology supports diagnosis but does not replace ENT evaluation***
- ***Safety comes before testing***

Golden Rule

Any ear discharge with foul smell or attic perforation is unsafe until proven otherwise.

A wet ear is a medical condition first and an audiologic case second.

Chapter

31

Vertigo



Vertigo is an illusion of movement, usually described as a sensation that the patient or surroundings are spinning.

It is a symptom, not a disease.

Vertigo vs Dizziness (Important Distinction)

Feature	Vertigo	Dizziness
Sensation	Spinning or rotational	Light headedness, imbalance
Origin	Vestibular system	Non vestibular causes common
Associated nausea	Common	Less common
Nystagmus	Usually present	Usually absent

Classification of Vertigo

Vertigo is classified into:

- Peripheral Vertigo
- Central Vertigo

This distinction is critical clinically and medico legally.

I. Peripheral Vertigo

Originates from the inner ear or vestibular nerve.

Common Causes

- ***Benign paroxysmal positional vertigo (BPPV)***
- ***Meniere disease***
- ***Vestibular neuritis***
- ***Labyrinthitis***
- ***Acoustic neuroma***
- ***Ototoxicity***



▲ *Figure31–1.Characteristic nystagmus in posterior canal BPPV.*


Vergence
Saccade
Optokinetic tracking
Smooth pursuit
Vestibulo-ocular reflex (VOR)
VOR cancellation
Head rotates and object fixed
Head and object rotate together

▲ *Figure 31–2.Classes of eye movements.*



▲ *Figure 31–3. (A) 45-degree side head rotation; (B) 30-degree head down position*


Gravity
Superior canal
45º
Posterior canal
Gravity
A
Particles
Utriculus
Ampulla of posterior canal
Gravity
Utriculus
Superior canal
Gravity
B
Particles
Posterior canal

▲ *Figure31–4. Dix-Hallpike test.*



First degree: Nystagmus appears only when the patient looks in the direction of fast component (opposite the side of hypoactive labyrinth)



Second degree: Nystagmus is visible in primary neutral position but not in the direction of slow component



Third degree: Nystagmus can be seen even in the direction of its slow component

▲ ***Figure 31–5.*** ***Alexander's law and the degree (severity) of nystagmus***


Utriculus
Superior canal
Posterior canal ampulla
Posterior canal
Particles
A
Posterior canal ampulla
Utriculus
Superior canal
Particles
Posterior canal
B
Superior canal
Posterior canal
Particles
Utriculus
Posterior canal ampulla
D
Posterior canal
Posterior canal ampulla
Particles
Utriculus
Superior canal
C

▲ *Figure 31–6. Epley maneuver.*

Clinical Features of Peripheral Vertigo

- ***Severe spinning sensation***
- ***Sudden onset***
- ***Associated nausea and vomiting***
- ***Hearing symptoms may be present***
- ***Nystagmus is usually horizontal or rotatory***
- ***Symptoms are positional in many cases***

II. Central Vertigo

Originates from the brainstem or cerebellum.

Common Causes

- *Stroke*
- *Multiple sclerosis*
- *Brain tumors*
- *Migraine related vertigo*
- *Vertebrobasilar insufficiency*

Clinical Features of Central Vertigo

- *Usually milder spinning sensation*
- *Gradual onset*
- *Neurological symptoms present*
- *Nystagmus may be vertical or direction changing*
- *Hearing usually normal*
- *Poor response to vestibular suppressants*

Red Flag

Central vertigo is potentially life threatening.

Peripheral vs Central Vertigo

Feature	Peripheral	Central
Severity	Severe	Mild to moderate
Onset	Sudden	Gradual
Hearing loss	Common	Rare
Tinnitus	Common	Rare
Nystagmus	Unidirectional	Direction changing
Fatigability	Present	Absent

Feature	Peripheral	Central
Neurological signs	Absent	Present


Central vertigo
Intermediate vertigo
Peripheral vertigo

▲ Figure 31–7. Diagrammatic representation of sites of the


Stimulation of crista ampullaris of semicircular canal
Ipsilateral vestibular nerve
Ipsilateral vestibular nuclei
MLF
Ipsilateral oculomotor complex
Contralateral abducent complex
Ipsilateral medial rectus
Contralateral lateral rectus
Threshold stimulus
Reticular formation of brain stem
Contralateral conjugate deviation of the eyes
Counter impulses to jerk eyes to original midline position
Slow phase of nystagmus
Fast phase of nystagmus

▲ *Flowchart 31–1. Mechanism of origin of nystagmus*


Vertigo
Full history and examination
Medical investigation
Neuro-otologic investigation
Neurologic investigation
Diagnosis and treatment
Diagnosis and treatment
Specific diagnosis
Peripheral labyrinthine vertigo
Medical treatment
Otologic surgery
Medical treatment of acute attack
Medical treatment of recurrent vertigo
Therapeutic procedures
Surgery
Destructive procedures
• Physical exercise
• Regimens of maneuver
• Psychological support
• Vestibular sedatives

▲ *Flowchart 31–2. Approach to a vertiginous patient*


Direction of view
straight lateral
Dark cells
Utricle
Saccule
Cochlea
Posterior
semicircular
canal
Displaced
otoconia

▲ Figure 31–8. Pathogenesis of BPPV showing displaced otoconia


Cupula
Cupulolithiasis
Ampulla
Canalithiasis

▲ Figure 31–9. Theories of BPPV



▲ *Figure 31–10. Epley manoeuvre steps*

Audiological Relevance

- ***Vertigo often coexists with hearing loss***
- ***Audiology helps in:***

– Differentiating cochlear from retrocochlear pathology

– Identifying asymmetry

– Supporting ENT diagnosis

Audiologic Findings in Vertigo

- ***Meniere disease***

– Low frequency SNHL

- ***Acoustic neuroma***

– Asymmetrical SNHL

– Poor speech discrimination

- ***Central vertigo***

– Audiogram often normal

Evaluation of Vertigo

History

- ***Onset and duration***
- ***Triggering factors***
- ***Positional relation***
- ***Hearing loss or tinnitus***
- ***Neurological symptoms***

Examination

- ***Otoscopic examination***
- ***Nystagmus assessment***
- ***Balance testing***
- ***Cranial nerve examination***

Investigations

- ***Pure tone audiometry***
- ***Speech audiometry***

- *Vestibular tests*
- *Imaging when indicated*

Management Principles

Peripheral Vertigo

- *Vestibular suppressants (short term)*
- *Etiology specific treatment*
- *Vestibular rehabilitation*

Central Vertigo

- *Urgent neurological evaluation*
- *Imaging*
- *Multidisciplinary management*

Audiology Red Flags in Vertigo

Immediate ENT or neurology referral if:

- *Acute vertigo with neurological deficits*
- *Persistent vertigo with normal ear findings*
- *Vertigo with sudden hearing loss*
- *Vertigo with facial weakness*
- *New onset vertigo in elderly patients*

Key Points

- *Vertigo is a symptom*
- *Peripheral and central causes must be distinguished*
- *Audiology supports diagnosis but does not replace medical evaluation*

Golden Rule

Vertigo with neurological signs is central until proven otherwise.

Audiology Checklist
Before Vestibular Testing

1. Patient Safety Screening

- ☐ ***Acute severe vertigo***
- ☐ ***Persistent vomiting***
- ☐ ***New neurological symptoms***
- ☐ ***Severe headache***
- ☐ ***Recent head trauma***
- ☐ ***Facial weakness or numbness***

STOP RULE

If any item is positive, do not proceed with vestibular testing and refer immediately.

2. History Review

- ☐ ***Onset, sudden or gradual***
- ☐ ***Duration of vertigo episodes***
- ☐ ***Positional trigger present or absent***

☐ Associated hearing loss

☐ Tinnitus or aural fullness

☐ Previous similar attacks

3. Otologic Assessment

☐ Otoscopic examination normal

☐ No active ear infection

☐ No ear discharge

☐ Tympanic membrane intact

Vestibular testing is deferred in active ear disease.

4. Audiologic Prerequisites

☐ Baseline pure tone audiometry completed

☐ Speech audiometry reviewed

☐ Asymmetry documented if present

5. Contraindications to Vestibular Testing

Unstable medical condition

Severe cervical spine disease

6. Documentation Statement

"Audiology safety screening was completed prior to vestibular testing.

Testing proceeded only after exclusion of medical red flags."

Golden Rule

Vestibular testing is diagnostic, not emergent care.

Safety comes first.

Peripheral vs Central Clues

Feature	Peripheral	Central
Severity	Severe	Mild
Hearing loss	Common	Rare
Tinnitus	Common	Rare
Nystagmus	One direction	Changing or vertical
Neuro signs	No	Yes

Central features → Emergency

32

Ear Operations

Ear operations are surgical procedures performed on the ear to treat disease, restore hearing, prevent complications, or improve quality of life.

Classification of Ear Operations

Ear operations are classified according to the part of the ear involved:

External ear operations

Middle ear operations

Inner ear operations

I. External Ear Operations

Common Indications

- ***Congenital deformities***
- ***Trauma***
- ***Infection or tumors***
- ***External auditory canal obstruction***

Common Procedures

1. Otoplasty

- ***Correction of protruding ear or deformity***
- ***Mainly cosmetic***
- ***No effect on hearing***

2. Excision of Preauricular Sinus

- ***Recurrent infection***
- ***Cosmetic reasons***

3. Excision of External Ear Tumors

- ***Benign or malignant lesions***
- ***Requires margin control***

4. Canalplasty

- ***Widening of the external auditory canal***
- ***Indicated in canal stenosis or exostosis***

II. Middle Ear Operations

These operations mainly aim at eradicating disease and restoring conductive hearing.

1. Myringotomy

- ***Incision in the tympanic membrane***
- ***Indications:***

– Acute otitis media with severe pain

– Otitis media with effusion

- ***Temporary procedure***

2. Tympanostomy Tube Insertion

- *Ventilation tube placed in TM*
- *Indications:*

– Persistent otitis media with effusion

– Recurrent acute otitis media

- *Common in children*

3. Tympanoplasty

- *Repair of tympanic membrane perforation*
- *May include ossicular reconstruction*
- *Goals:*

– Dry ear

– Improved hearing

4. Ossiculoplasty

- *Reconstruction of ossicular chain*
- *Indications:*

– Ossicular erosion

– Discontinuity

- *May use prosthesis*

5. Mastoidectomy

Cortical Mastoidectomy

- *Removal of mastoid air cell disease*
- *Often combined with tympanoplasty*

Modified Radical Mastoidectomy

- ***Canal wall down***
- ***For cholesteatoma with hearing preservation***

Radical Mastoidectomy

- ***Extensive disease***
- ***Hearing usually sacrificed***

III. Inner Ear Operations

These procedures are less common and carry higher risk.

1. Stapedectomy or Stapedotomy

- ***Surgery for otosclerosis***
- ***Improves conductive hearing loss***
- ***Requires careful audiologic follow up***

2. Labyrinthectomy

- ***Destruction of vestibular labyrinth***
- ***Indicated in intractable vertigo***
- ***Causes complete hearing loss in operated ear***

3. Endolymphatic Sac Surgery

- ***For Meniere disease***
- ***Aims to control vertigo***
- ***Hearing preservation varies***



▲ *Figure 32–1.* ***(A) Endolymphatic sac surgery. (B) Vestibular nerve section***

4. Cochlear Implant Surgery

- ***For severe to profound SNHL***
- ***Bypasses damaged cochlea***
- ***Requires extensive audiologic rehabilitation***

Audiology Considerations Before Ear Surgery

- ***Baseline pure tone audiometry***
- ***Speech audiometry***
- ***Tympanometry when appropriate***
- ***Counseling regarding expected hearing outcome***

Audiology Considerations After Ear Surgery

- ***Post operative hearing assessment***
- ***Monitoring of hearing improvement or deterioration***

- *Identification of complications*
- *Hearing aid fitting if required*

Red Flags After Ear Operations

Immediate ENT referral if:

- ***Severe vertigo***
- ***Sudden hearing loss***
- ***Facial nerve weakness***
- ***Persistent ear discharge***
- ***Severe otalgia***

Key Points

- ***Ear operations are classified by ear part involved***
- ***Middle ear surgery aims at disease control and hearing restoration***
- ***Inner ear surgery carries higher risk***
- ***Audiology plays a key role before and after surgery***

Golden Rule

No ear surgery is complete without proper audiologic assessment.

Audiology Checklist

Before Ear Surgery

Patient and History

☐ Procedure planned: ____________________

☐ Side: ☐ Right ☐ Left ☐ Bilateral

☐ Main complaint: ☐ Hearing loss ☐ Otorrhea ☐ Otalgia ☐ Vertigo ☐ Tinnitus

☐ Duration and progression documented

☐ Previous ear surgery: ☐ No ☐ Yes, details __________

☐ Noise exposure or ototoxic drugs reviewed

☐ Fitness for testing, patient cooperation confirmed

2. Safety and Otoscopy

☐ Otoscopic exam documented

☐ TM status: ☐ Intact ☐ Perforated ☐ Retracted ☐ Ventilation tube

☐ Active discharge: ☐ No ☐ Yes

☐ Suspicion of cholesteatoma or vascular mass: ☐ No ☐ Yes

If unsafe ear features present, limit testing and refer to ENT.

3. Baseline Audiology Tests

☐ Pure tone audiometry, air and bone

☐ Masking applied when indicated

☐ Speech audiometry, SRT and SDS

☐ Tympanometry if TM intact and ear dry

☐ Acoustic reflexes when appropriate

4. Interpretation and Surgical Relevance

☐ *Type of loss confirmed:* ☐ *Conductive* ☐ *SNHL* ☐ *Mixed*

☐ *Air bone gap size documented*

☐ *Speech discrimination adequacy documented*

☐ *Asymmetry assessed*

☐ *Retrocochlear red flags excluded*

5. Pre Operative Counseling Documentation

☐ *Expected hearing outcome explained*

☐ *Possibility of no hearing improvement discussed*

☐ *Risk of SNHL explained for relevant surgeries*

☐ *Hearing aid plan discussed when applicable*

6. Required Report Statements

"Audiologic findings represent baseline functional hearing status prior to surgery."

"Post operative audiologic reassessment is recommended according to protocol."

Golden Rule

No ear surgery without a documented baseline audiogram and speech scores.

Post Operative Audiology Follow Up Protocol

Ear Surgery

1. Immediate Post Op Phase

Audiology testing is generally not required in the first 1–2 weeks unless urgent symptoms occur.

Urgent referral if:

☐ ***Sudden hearing loss***

☐ ***Severe vertigo***

☐ ***Facial weakness***

☐ ***Persistent bleeding or discharge***

2. Early Follow Up

4–6 Weeks Post Op

☐ ***Pure tone audiometry, air and bone***

☐ ***Speech audiometry***

☐ ***Compare to pre op baseline***

☐ ***Document ABG change if conductive component expected***

3. Standard Follow Up

3 Months Post Op

☐ Repeat PTA and speech

☐ Tympanometry if appropriate

☐ Determine hearing stabilization

☐ Hearing aid assessment if needed

4. Long Term Follow Up

6–12 Months Post Op

☐ Annual audiogram for chronic ear disease

☐ Ongoing monitoring after cholesteatoma surgery

☐ CI patients follow dedicated mapping schedule

Documentation Standards

"All post operative results are compared directly to the pre operative baseline audiogram."

"Any unexpected deterioration triggers ENT review."

Key Message

Post op audiology is not one test, it is trend monitoring.

Comparison Table

Ear Operations and Expected Hearing Outcome

Operation	*Main Goal*	*Typical Hearing Outcome*	*Audiology Notes*
Myringotomy	*Drainage*	*Temporary change*	*Baseline audiogram useful*
Ventilation tube	*Ventilation*	*Often improves CHL*	*Tympanometry changes expected*
Tympanoplasty	*Repair TM*	*Improves CHL*	*ABG reduction expected*
Ossiculoplasty	*Rebuild chain*	*Variable improvement*	*ABG may persist*
Mastoidectomy	*Eradicate disease*	*Variable*	*Dry ear priority over hearing*
Stapedotomy	*Otosclerosis*	*Excellent CHL improvement*	*Watch for SNHL risk*
Labyrinthectomy	*Control vertigo*	*Hearing sacrificed*	*Counsel clearly*
Endolymphatic sac surgery	*Control vertigo*	*Hearing variable*	*Monitor fluctuation*
Cochlear implant	*Restore access to sound*	*Speech improves*	*Needs mapping and rehab*

Final Clinical Message

Surgical success is not only a dry ear,

it is also documented hearing outcomes and counseling aligned with reality.

Chapter



33

Rehabilitation of the Impaired Hearing

Hearing rehabilitation is the process of reducing the impact of hearing loss on communication, daily activities, and quality of life through medical, audiologic, technological, and rehabilitative interventions.

Rehabilitation focuses on the person, not only the audiogram.

Goals of Hearing Rehabilitation

- Improve speech understanding
- Enhance communication ability
- Reduce listening effort
- Improve quality of life
- Promote social and emotional well-being

Multidisciplinary Approach

Effective hearing rehabilitation involves:

- Audiologist
- ENT specialist
- Speech and language therapist
- Patient and family

Types of Hearing Rehabilitation

Hearing rehabilitation depends on:

- ***Type of hearing loss***
- ***Degree of hearing loss***
- ***Age of patient***
- ***Communication needs***

I. Medical and Surgical Rehabilitation

Managed primarily by ENT.

Examples

- ***Treatment of otitis media***
- ***Tympanic membrane repair***
- ***Ossicular reconstruction***
- ***Stapes surgery***
- ***Cochlear implantation***

Audiology role:

- ***Baseline assessment***
- ***Outcome measurement***
- ***Post operative follow up***

II. Audiologic Rehabilitation

Core responsibility of Audiology.

1. Hearing Aids

Indications

- ***Mild to severe hearing loss***
- ***Conductive, sensorineural, or mixed loss***

Benefits

- ***Amplify sound***
- ***Improve speech audibility***
- ***Reduce listening effort***

Limitations

- ***Do not restore normal hearing***
- ***Require adaptation and counseling***


Ear hook
Volume control
Battery compartment
Earmold
Battery compartment
Volume control
Volume control
Battery compartment
Ear canal
Pull string
Tympanic membrane

▲ *Figure 33–1. Different types of hearing aids*

2. Cochlear Implants

Indications

- *Severe to profound SNHL*
- *Limited benefit from hearing aids*

Audiology role:

- *Pre implantation evaluation*
- *Candidacy assessment*
- *Post implantation mapping*
- *Auditory training*


Transmitter coil
Receiver internal coil
Electrodes
8th nerve
Speech processor

▲ *Figure 33–2. Multichannel cochlear implant*

3. Bone Conduction Devices

- *For conductive or mixed hearing loss*
- *For single sided deafness*

Examples:

- ***Bone anchored hearing aids***
- ***Bone conduction implants***

III. Auditory Training and Rehabilitation Therapy

Auditory training helps patients:

- ***Improve speech perception***
- ***Adapt to hearing devices***
- ***Enhance listening skills***

Includes:

- ***Speech reading***
- ***Listening exercises***
- ***Environmental sound awareness***

IV. Communication Strategies

Patients are trained to:

- ***Face the speaker***
- ***Reduce background noise***
- ***Use visual cues***
- ***Request clarification***

Family counseling is essential.

V. Pediatric Hearing Rehabilitation

Key Principles

- ***Early identification***
- ***Early amplification***
- ***Family centered intervention***

Includes:

- ***Hearing aids or implants***
- ***Speech and language therapy***
- ***Educational support***

Delay in rehabilitation may lead to permanent language deficit.

VI. Rehabilitation of Special Hearing Problems

Tinnitus Rehabilitation

- ***Counseling***
- ***Sound therapy***
- ***Hearing aids when indicated***

Auditory Processing Disorders

- ***Auditory training***
- ***Classroom accommodations***

VII. Psychosocial Aspects

Hearing loss may cause:

- ***Social withdrawal***
- ***Depression***
- ***Anxiety***

Rehabilitation must address emotional impact.

Role of Audiology in Hearing Rehabilitation

Audiology is responsible for:

- ***Accurate diagnosis***
- ***Device selection and fitting***

- *Verification and validation*
- *Counseling and follow up*

Audiology does not promise cure, but functional improvement.

Outcome Measures in Rehabilitation

- *Speech audiometry*
- *Questionnaires*
- *Patient reported benefit*
- *Quality of life scales*

Key Points

- *Hearing rehabilitation is individualized*
- *Technology alone is not enough*
- *Counseling is essential*
- *Early intervention improves outcomes*

Golden Rule

The success of hearing rehabilitation is measured by communication improvement, not by decibels alone.

Chapter



34

Hearing Aid Fitting and Follow Up Protocol

1. Pre Fitting Assessment

☐ Complete audiologic evaluation

☐ Type and degree of hearing loss confirmed

☐ Speech audiometry completed

☐ Medical clearance obtained if required

☐ Patient communication needs assessed

2. Counseling Before Fitting

☐ Realistic expectations explained

☐ Adaptation period discussed

☐ Limitations of hearing aids explained

☐ Maintenance and daily use reviewed

3. Hearing Aid Selection

☐ Style selected based on audiogram and anatomy

☐ Technology level matched to needs

☐ Monaural or binaural decision documented

4. Hearing Aid Fitting

☐ ***Initial programming completed***

☐ ***Comfort and loudness verified***

☐ ***Patient instruction provided***

☐ ***Safety and volume limits set***

5. Early Follow Up

2 to 4 weeks

☐ ***Patient comfort reviewed***

☐ ***Gain adjustment if needed***

☐ ***Counseling reinforced***

6. Outcome Evaluation

6 to 8 weeks

☐ ***Speech testing with aids***

☐ ***Subjective benefit assessed***

☐ ***Daily usage discussed***

7. Long Term Follow Up

☐ ***Review every 6 to 12 months***

☐ ***Audiogram update***

Golden Rule

A hearing aid fitting is a process, not a single visit.

Implantable Hearing Devices

Implantable hearing devices are surgically placed systems designed to improve hearing by delivering sound to the auditory system through mechanical vibration or electrical stimulation, bypassing part or all of the external and middle ear.

They are used when conventional hearing aids are insufficient, contraindicated, or poorly tolerated.

Goals of Implantable Hearing Devices

- ***Improve audibility and speech understanding***
- ***Bypass diseased or malformed ear structures***
- ***Provide stable and predictable sound transmission***
- ***Improve quality of life***

Classification of Implantable Hearing Devices

Implantable hearing devices are broadly classified into:

Middle Ear Implantable Hearing Devices
Bone Conduction Implantable Devices
Cochlear Implants

35

Middle Ear Implantable Hearing Devices

Devices that mechanically stimulate the ossicular chain or inner ear structures.

Indications

- ***Moderate to severe sensorineural hearing loss***
- ***Mixed hearing loss***
- ***Patients intolerant to conventional hearing aids***
- ***External ear canal problems***

Types

- ***Partially implantable middle ear devices***
- ***Fully implantable middle ear devices***

Advantages

- ***Improved sound quality***
- ***Reduced feedback***
- ***No occlusion of ear canal***

Limitations

- ***Surgical risks***
- ***Higher cost***

- ***Limited availability***
- ***Variable outcomes***


FMT
VORP
VORP 503
12345
VIBRANT
Audio processor receiver
Vibrating ossicular prosthesis (VORP) implant
Conductor link
Floating mass transducer
Detail of floating mass transducer
A
B
SP
LP
RW
Clip

Figure 35–1. Vibrant Soundbridge (MED EL) AMEI and couplers.



Figure 35–2. MED- EL audiometric indications for the Vibrant Soundbridge AMEI.

II. Bone Conduction Implantable Devices

Devices that transmit sound via bone vibration directly to the cochlea, bypassing the outer and middle ear.

Indications

- ***Conductive hearing loss***
- ***Mixed hearing loss***
- ***Congenital aural atresia***
- ***Chronic otitis media***
- ***Single sided deafness***

Types

- ***Percutaneous bone anchored devices***
- ***Transcutaneous bone conduction implants***

Advantages

- ***Reliable hearing improvement***
- ***Bypasses middle ear pathology***
- ***Suitable for congenital ear anomalies***

Limitations

- ***Surgical procedure required***
- ***Skin or implant related complications***
- ***Cosmetic concerns in some patients***

III. Cochlear Implants

A cochlear implant is an electronic medical device that directly stimulates the auditory nerve, bypassing damaged cochlear hair cells.

Indications

- ***Severe to profound sensorineural hearing loss***
- ***Limited benefit from hearing aids***
- ***Post lingual or pre lingual deafness***

Components

- ***External sound processor***
- ***Internal receiver stimulator***
- ***Electrode array***

Audiology Role

- ***Candidacy assessment***
- ***Pre implant counseling***
- ***Post implant mapping***
- ***Auditory rehabilitation***

Limitations

- ***Requires surgery***
- ***Intensive rehabilitation***
- ***Speech outcomes vary***

Cochlear Implant Audiology Pathway

1. Referral to CI Program

☐ Severe to profound SNHL

☐ Limited benefit from hearing aids

☐ Motivation and realistic expectations assessed

2. Pre Implant Audiology Assessment

☐ Unaided audiogram

☐ Aided audiogram

☐ Speech perception testing

☐ Hearing aid trial documented

3. Multidisciplinary CI Candidacy

- Audiology
- ENT
- Speech and language therapy
- Psychology when indicated

4. Pre Implant Counseling

☐ Nature of CI explained

☐ Surgical and auditory outcomes discussed

☐ Rehabilitation commitment explained

5. Post Surgical Activation

3 to 4 weeks post surgery

☐ Initial device activation

☐ Basic mapping performed

☐ Sound awareness counseling

6. Early Mapping Phase

First 3 months

☐ Multiple mapping sessions

☐ Progressive stimulation increase

☐ Auditory training initiated

7. Rehabilitation Phase

☐ Speech perception therapy

☐ Environmental sound training

☐ Family involvement encouraged

8. Long Term Follow Up

☐ Regular mapping

☐ Annual speech evaluation

☐ Device integrity check

Mandatory Documentation Line

"Cochlear implantation provides access to sound. Speech understanding improves gradually with rehabilitation."

Final Clinical Message

Hearing rehabilitation succeeds when technology, training, and counseling work together.



▲ Figure 35–3. The audiological criteria for cochlear implantation

Comparison of Implantable Hearing Devices

Device Type	*Hearing Loss*	*Primary Benefit*	*Key Limitation*
Middle ear implant	*SNHL or mixed*	*Natural sound quality*	*Cost and surgery*
Bone conduction implant	*Conductive or mixed*	*Bypasses middle ear*	*Skin or implant issues*
Cochlear implant	*Severe to profound SNHL*	*Access to sound*	*Rehab dependent*

Pre Implant Audiologic Assessment

Audiology must perform:

- *Complete audiologic evaluation*
- *Speech perception testing*
- *Aided and unaided assessment*
- *Counseling on realistic expectations*

Audiology does not decide surgical candidacy alone.

Post Implant Follow Up

- *Device activation*
- *Programming or mapping*
- *Objective and subjective outcome measures*
- *Long term monitoring*

Pediatric Considerations

- *Early implantation improves outcomes*
- *Family counseling is essential*
- *Multidisciplinary approach required*

Medico Legal and Ethical Considerations

- *Informed consent is mandatory*
- *Outcomes must not be guaranteed*
- *Device selection must be patient centered*
- *Conflicts of interest must be disclosed*

Key Points

- *Implantable devices bypass external amplification limits*
- *Proper patient selection is critical*

- *Audiology plays a central role before and after implantation*
- *Rehabilitation determines success*

Golden Rule

An implant is not a cure.

It is a rehabilitation tool that requires commitment and follow up.

Cochlear Implants VS Auditory Brainstem Implants

Cochlear implants (CI) and auditory brainstem implants (ABI) are advanced implantable hearing devices designed for patients with severe to profound hearing loss who derive limited or no benefit from conventional hearing aids.

While both devices aim to provide access to sound, they differ fundamentally in:

- *Site of stimulation*
- *Indications*
- *Expected outcomes*
- *Rehabilitation requirements*

Cochlear Implants

A cochlear implant is an electronic medical device that directly stimulates the auditory nerve fibers within the cochlea, bypassing damaged or non-functional hair cells.

Components of a Cochlear Implant

- ***External sound processor***
- ***Microphone and speech processor***
- ***Transmitting coil***
- ***Internal receiver stimulator***
- ***Electrode array inserted into the cochlea***

Mechanism of Action

Sound is:

- ***Captured by the microphone***
- ***Processed into electrical signals***
- ***Transmitted to the implanted receiver***
- ***Delivered to the auditory nerve via electrodes***

This provides auditory perception, not normal hearing.

Indications for Cochlear Implantation

- ***Severe to profound bilateral sensorineural hearing loss***
- ***Poor speech perception despite appropriately fitted hearing aids***
- ***Congenital or acquired deafness***
- ***Post meningitis hearing loss***
- ***Progressive hearing loss***

Contraindications

- ***Absence of the auditory nerve***
- ***Severe cochlear aplasia***
- ***Active infection***
- ***Medical contraindications to surgery***

Audiology Role in Cochlear Implants

Audiology is responsible for:

- ***Pre implant audiologic assessment***
- ***Hearing aid trial documentation***
- ***Speech perception testing***
- ***Counseling regarding expectations***
- ***Post implant activation and mapping***
- ***Long term auditory rehabilitation***

Audiology does not guarantee outcomes.

Outcomes of Cochlear Implants

- ***Improved sound awareness***
- ***Improved speech understanding***
- ***Outcomes vary widely***
- ***Depend on:***

– Age at implantation

– Duration of deafness

– Auditory nerve integrity

– Rehabilitation compliance

Auditory Brainstem Implants

An auditory brainstem implant is a device that directly stimulates the cochlear nucleus in the brainstem, bypassing both the cochlea and auditory nerve.

It is used when cochlear implantation is not possible.

Indications for Auditory Brainstem Implant

- ***Bilateral absence of the auditory nerve***
- ***Cochlear nerve aplasia or severe hypoplasia***
- ***Neurofibromatosis type II***
- ***Severe cochlear malformations***
- ***Failed cochlear implantation***

Components of an ABI

- ***External sound processor***
- ***Internal receiver***
- ***Electrode array placed on the cochlear nucleus***

Mechanism of Action

- ***Sound is converted into electrical signals***
- ***Signals directly stimulate the brainstem auditory pathways***
- ***Auditory sensations are produced without cochlear input***

Outcomes of ABI

- ***Sound awareness is common***
- ***Speech understanding is limited***
- ***Outcomes are less predictable than CI***
- ***Rehabilitation is prolonged and intensive***

Pediatric Considerations

- ***Early cochlear implantation improves language outcomes***
- ***ABI in children requires careful ethical consideration***
- ***Family counseling is essential***
- ***Multidisciplinary decision making is mandatory***



▲ *Figure 35–4. technique for auditory brainstem implant (ABI*

Cochlear Implant vs Auditory Brainstem Implant

Feature	*Cochlear Implant*	*Auditory Brainstem Implant*
Site of stimulation	*Cochlea*	*Brainstem (cochlear nucleus)*
Auditory nerve required	*Yes*	*No*
Typical outcome	*Speech understanding*	*Sound awareness*

Feature	Cochlear Implant	Auditory Brainstem Implant
Indications	Severe SNHL	Absent auditory nerve
Rehab complexity	High	Very high
Availability	Widely available	Limited, specialized centers

Rehabilitation and Follow Up

Cochlear Implant

- Initial activation after wound healing
- Regular mapping sessions
- Auditory training
- Speech and language therapy

Audiology and ENT Shared Responsibility

- Audiology: assessment, mapping, rehabilitation, counseling
- ENT: diagnosis, imaging, surgical implantation, medical follow up

Clear documentation and communication are essential to reduce medico-legal risk.

Key Points

- Cochlear implants stimulate the auditory nerve
- Auditory brainstem implants bypass the auditory nerve
- CI outcomes are generally superior to ABI
- Rehabilitation determines functional benefit

Before Cochlear Implant or Auditory Brainstem Implant

1. Patient and History

☐ Type of hearing loss confirmed

☐ Duration of deafness documented

☐ Onset: congenital or acquired

☐ Etiology suspected or known

☐ Post meningitis or progressive loss noted

2. Audiologic Assessment

☐ Unaided pure tone audiometry

☐ Aided audiogram

☐ Speech perception testing

☐ Hearing aid trial documented

☐ Limited benefit clearly recorded

SECTION 5

Pediatric Otology

Chapter

36

Pediatric Otology

Pediatric otology focuses on ear diseases and hearing disorders in infants and children, with special emphasis on early detection, timely intervention, and prevention of long-term auditory and language consequences.

Children are not small adults.

Otologic disorders in childhood require age-specific evaluation, interpretation, and management.

Importance of Pediatric Otology

- ***Hearing is critical for speech and language development***
- ***Delayed diagnosis leads to permanent communication deficits***
- ***Many pediatric ear conditions are silent or asymptomatic***
- ***Audiology plays a central role in early identification***

Anatomical and Physiological Considerations

Key pediatric differences include:

- ***Narrower external auditory canal***
- ***More horizontal and shorter Eustachian tube***

- *Immature immune system*
- *Ongoing maturation of auditory pathways*

These factors increase susceptibility to:

- *Otitis media*
- *Conductive hearing loss*
- *Fluctuating hearing levels*

Audiologic Assessment in Children

Principles

- *Age appropriate test selection*
- *Objective testing when behavioral responses are unreliable*
- *Repeated assessment when needed*

Common Tests

- *Otoacoustic emissions*
- *Auditory Brainstem Response*
- *Behavioral audiometry*
- *Tympanometry*

Audiology must document test reliability and limitations.

Red Flags in Pediatric Otology

Immediate referral is required if:

- *Failed newborn hearing screening*
- *Speech or language delay*
- *Recurrent otitis media with hearing concerns*
- *Post meningitis recovery*
- *Suspected unilateral hearing loss*

Role of Audiology

Audiology is responsible for:

- *Early identification of hearing loss*
- *Objective hearing assessment*
- *Monitoring fluctuating hearing*
- *Rehabilitation planning*
- *Family counseling*

Audiology does not make medical diagnoses.

Role of ENT

ENT is responsible for:

- *Medical diagnosis*
- *Surgical decision making*
- *Imaging*
- *Medical management*

Multidisciplinary Approach

Effective pediatric otology care involves:

- *Audiology*
- *ENT*
- *Pediatrics*
- *Speech and language therapy*
- *Family*

Medico-Legal Considerations

- *Delay in diagnosis may result in lifelong disability*
- *Documentation of follow up is essential*

- *Parental counseling must be recorded*
- *Missed post meningitis hearing loss is a major liability risk*

Key Points

- *Pediatric ear disorders differ from adults*
- *Early hearing access is critical*
- *Audiology is central to detection and monitoring*
- *Team-based care improves outcomes*

Golden Rule

In children, hearing loss is a developmental emergency until proven otherwise.

Genetics of Hearing Loss

Genetic factors are responsible for a significant proportion of hearing loss, particularly in children.

More than 50% of permanent childhood sensorineural hearing loss has a genetic basis, either isolated or as part of a syndrome.

Genetic hearing loss may be:

- *Congenital or delayed onset*
- *Progressive or stable*
- *Syndromic or non-syndromic*

Understanding genetic mechanisms is essential for accurate diagnosis, prognosis, counseling, and management planning.

Classification of Genetic Hearing Loss

1. Non-Syndromic Hearing Loss

- *Hearing loss is the only clinical manifestation*
- *Accounts for approximately 70% of genetic hearing loss*
- *Usually sensorineural*
- *Most commonly inherited as autosomal recessive*

2. Syndromic Hearing Loss

- *Hearing loss occurs with other systemic abnormalities*
- *Accounts for approximately 30% of genetic cases*
- *May involve vision, balance, heart, kidneys, thyroid, or skeletal system*

Modes of Inheritance

1. Autosomal Recessive

- *Most common inheritance pattern*
- *Parents usually have normal hearing*
- *Hearing loss often congenital and severe to profound*

2. Autosomal Dominant

- *Progressive hearing loss is common*
- *Onset may be delayed*
- *Often milder initially*

3. X-Linked

- *Rare*
- *Males more severely affected*

4. Mitochondrial

- *Maternal inheritance*
- *Often associated with aminoglycoside ototoxicity susceptibility*

Common Genetic Causes of Non-Syndromic Hearing Loss

Connexin 26 (GJB2) Mutation

- ***Most common genetic cause worldwide***
- ***Autosomal recessive***
- ***Congenital, bilateral SNHL***
- ***Variable severity***
- ***Normal imaging***

Clinical Insight

Normal CT or MRI does not exclude genetic hearing loss.

Common Syndromic Hearing Loss Disorders

Usher Syndrome

- ***Hearing loss + retinitis pigmentosa***
- ***Vestibular dysfunction common***
- ***Progressive visual loss***

Waardenburg Syndrome

- ***SNHL***
- ***Pigmentary abnormalities***
- ***Dystopia canthorum***

Pendred Syndrome

- ***SNHL***
- ***Thyroid involvement***
- ***Enlarged vestibular aqueduct***

Alport Syndrome

- *Progressive SNHL*
- *Renal disease*
- *Ocular abnormalities*

Audiologic Characteristics of Genetic Hearing Loss

- *Bilateral and symmetrical in most cases*
- *May be present at birth or appear later*
- *May be progressive*
- *Speech perception varies with neural involvement*

Red Flag

Asymmetry or sudden onset suggests non-genetic pathology and requires medical evaluation.

Audiology Role in Genetic Hearing Loss

Audiology responsibilities include:

- *Early identification*
- *Accurate characterization of hearing loss*
- *Monitoring progression*
- *Counseling families*
- *Coordinating referrals*

Audiologists do not make genetic diagnoses.

Indications for Genetic Referral

Referral for genetic evaluation should be considered when:

- ***Congenital bilateral SNHL***
- ***Positive family history***
- ***Syndromic features suspected***
- ***Enlarged vestibular aqueduct***
- ***Progressive childhood hearing loss***

Pediatric Considerations

- ***Early diagnosis improves language outcomes***
- ***Genetic counseling helps family planning***
- ***Results guide prognosis and CI candidacy***

Management Implications

Hearing Rehabilitation

- ***Hearing aids***
- ***Cochlear implants when indicated***
- ***Early intervention services***

Common Pitfall

Labeling hearing loss as "idiopathic" without considering genetic causes.

Key Points

- ***Genetics is a major cause of pediatric hearing loss***
- ***Non-syndromic cases are more common***

- *Audiology plays a key role in detection and monitoring*
- *Genetic diagnosis guides prognosis, not just etiology*

Golden Rule

When hearing loss starts early and runs in families,
genetics must be considered, even if imaging is normal.

Otologic Syndromes

Otologic syndromes are genetic or developmental disorders in which hearing loss is a major clinical feature, occurring alongside abnormalities in other organ systems.

Recognition of otologic syndromes is critical because:

- ***Hearing loss may be progressive***
- ***Associated systemic conditions may be life-threatening***
- ***Early diagnosis affects prognosis, rehabilitation, and counseling***

Classification of Otologic Syndromes

Otologic syndromes are broadly classified into:

Syndromes with predominant sensorineural hearing loss
Syndromes with mixed or conductive hearing loss
Syndromes with associated vestibular dysfunction

Common Otologic Syndromes

1. Usher Syndrome

Key Features

- *Congenital or early-onset SNHL*
- *Retinitis pigmentosa*
- *Possible vestibular dysfunction*

Audiologic Characteristics

- *Bilateral SNHL*
- *Variable severity*
- *Progressive visual impairment affects rehabilitation*

Audiology Note

Speech perception may be limited by combined auditory and visual loss.

2. Waardenburg Syndrome

Key Features

- *Congenital SNHL*
- *Pigmentary abnormalities*
- *Dystopia canthorum*

Audiologic Characteristics

- *SNHL ranging from mild to profound*
- *May be unilateral or bilateral*

3. Pendred Syndrome

Key Features

- *SNHL*
- *Thyroid dysfunction*
- *Enlarged vestibular aqueduct*

Audiologic Characteristics

- *Fluctuating or progressive SNHL*
- *Sudden drops in hearing may occur*

Red Flag

Minor head trauma may cause sudden hearing deterioration.

4. Alport Syndrome

Key Features

- *Progressive SNHL*
- *Renal disease*
- *Ocular abnormalities*

Audiologic Characteristics

- *High-frequency SNHL*
- *Progressive course*

5. Branchio-Oto-Renal Syndrome

Key Features

- *External ear anomalies*
- *Conductive or mixed hearing loss*
- *Renal malformations*

Audiologic Characteristics

- *Variable hearing loss type*
- *Structural abnormalities common*

6. CHARGE Syndrome

Key Features

- *Ear anomalies*

- ***Cranial nerve involvement***
- ***Balance dysfunction***
- ***Multisystem involvement***

Audiologic Characteristics

- ***Conductive, sensorineural, or mixed loss***
- ***Absent or malformed semicircular canals***

Audiologic Characteristics of Otologic Syndromes

Common patterns include:

- ***Bilateral hearing loss***
- ***Early onset***
- ***Progressive or fluctuating course***
- ***Associated vestibular dysfunction***

Red Flag

Asymmetry or sudden onset requires exclusion of non-syndromic pathology.

Audiology Role in Otologic Syndromes

Audiologists are responsible for:

- ***Early identification of hearing loss***
- ***Accurate classification of hearing type***
- ***Monitoring progression***
- ***Rehabilitation planning***
- ***Family counseling***

Audiology does not establish the genetic diagnosis.

Indications for Multidisciplinary Referral

Referral is recommended when:

- ***Hearing loss is congenital***
- ***Syndromic features are present***
- ***Family history is positive***
- ***Hearing loss is progressive***

Team members include:

- ***ENT***
- ***Genetics***
- ***Pediatrics***
- ***Ophthalmology***
- ***Nephrology when indicated***

Management Principles

Hearing Rehabilitation

- ***Hearing aids***
- ***Cochlear implantation when appropriate***
- ***Early intervention programs***

Long-Term Monitoring

- ***Regular audiologic follow up***
- ***Surveillance for systemic complications***

Medico-Legal Considerations

Audiology documentation must:

- ***Avoid assigning specific syndromic diagnosis***
- ***Recommend appropriate referrals***

- *Document counseling clearly*
- *Avoid false reassurance*

Common Pitfall

Treating syndromic hearing loss as isolated SNHL without further referral.

Key Points

- *Otologic syndromes are common in pediatric hearing loss*
- *Hearing loss is often bilateral and progressive*
- *Audiology plays a key role in detection and monitoring*
- *Multidisciplinary care is essential*

Golden Rule

When hearing loss comes with other system findings,
think syndrome until proven otherwise.

Congenital Hearing Loss

***Congenital hearing loss** is hearing loss that is **present at birth or develops shortly after**, and it represents one of the **most critical conditions in pediatric otology** because of its direct impact on **speech, language, and cognitive development**.*

Congenital hearing loss may be:

- *Genetic or non-genetic*
- *Syndromic or non-syndromic*

- *Conductive, sensorineural, or mixed*
- *Stable, progressive, or fluctuating*

Early identification and intervention are essential to prevent permanent developmental consequences

Epidemiology

- *Occurs in approximately 1–3 per 1,000 live births*
- *Higher prevalence in:*

– NICU graduates

– Premature infants

– Children with congenital anomalies

- *One of the most common congenital disorders worldwide*

Etiology of Congenital Hearing Loss

1. Genetic Causes

- *Account for more than 50% of congenital permanent hearing loss*
- *May be syndromic or non-syndromic*
- *Often autosomal recessive*

2. Non-Genetic Causes

Prenatal

- *Intrauterine infections, TORCH*
- *Ototoxic drug exposure*
- *Maternal metabolic disease*

Perinatal

- *Prematurity*
- *Hypoxia*

- ***Hyperbilirubinemia***
- ***NICU stay***

Postnatal Early Life

- ***Meningitis***
- ***Ototoxic medications***
- ***Severe neonatal illness***

Types of Congenital Hearing Loss

Sensorineural Hearing Loss

- ***Most common type***
- ***Usually bilateral***
- ***Often permanent***
- ***May be progressive***

Conductive Hearing Loss

- ***External or middle ear malformations***
- ***Otitis media with effusion***
- ***May be transient or correctable***

Mixed Hearing Loss

- ***Combination of structural and cochlear pathology***

Audiologic Characteristics

Typical features include:

- ***Absent or abnormal newborn hearing screening***
- ***Bilateral symmetrical loss in many cases***
- ***Normal tympanometry in SNHL***
- ***Speech delay as the primary presenting complaint***

Red Flag

Unilateral congenital hearing loss must never be considered benign.

Audiologic Assessment

Audiologic evaluation must be age-appropriate and objective.

Core Tests

- ***Otoacoustic emissions***
- ***Auditory Brainstem Response***
- ***Tympanometry***
- ***Behavioral audiometry when age allows***

Audiology must document:

- ***Test reliability***
- ***Limitations***
- ***Need for repeat testing***

Newborn Hearing Screening

- ***Universal newborn hearing screening is mandatory***
- ***Failed screening requires prompt diagnostic evaluation***
- ***"Refer" does not equal diagnosis***

Delay in follow-up is a major medico-legal risk.

Management Principles

Early Intervention

- ***Hearing aids when indicated***
- ***Cochlear implantation for severe to profound loss***
- ***Early speech and language therapy***

Multidisciplinary Care

- ***Audiology***
- ***ENT***
- ***Pediatrics***
- ***Speech and language therapy***
- ***Genetics when indicated***

Prognosis

- ***Outcomes depend on:***

– Age at diagnosis

– Severity of loss

– Consistency of amplification

– Family involvement

Early identification before 6 months of age significantly improves outcomes.

Golden Rule

In congenital hearing loss,
time lost equals language lost.

Management of Children with Inner Ear Malformations

Inner ear malformations are congenital abnormalities of the cochlea, vestibular system, or both, resulting from arrested or abnormal embryologic development.

They represent a significant cause of congenital sensorineural hearing loss and pose unique challenges in:

- *Audiologic diagnosis*
- *Hearing rehabilitation*
- *Cochlear implant candidacy*
- *Surgical and medico-legal decision making*

Embryologic Basis

- *Inner ear development occurs early in gestation*
- *Arrest at different stages produces distinct malformation patterns*
- *Severity of hearing loss correlates with degree of structural abnormality*

Classification of Inner Ear Malformations

1. Complete Labyrinthine Aplasia (Michel Deformity)

- *Absence of cochlea and vestibular system*
- *Profound bilateral SNHL*
- *Cochlear implantation not possible*

2. Cochlear Aplasia

- *Absence of cochlea*
- *Vestibular system may be partially present*
- *Profound SNHL*

3. Common Cavity Deformity

- *Cochlea and vestibule form a single cavity*
- *Severe to profound SNHL*
- *CI outcomes variable*

4. Incomplete Partition (IP)

IP Type I

- *Cystic cochlea without modiolus*
- *Severe to profound SNHL*

IP Type II (Mondini Malformation)

- *One and a half turns of cochlea*
- *Commonly associated with enlarged vestibular aqueduct*
- *Hearing loss may be progressive or fluctuating*

5. Enlarged Vestibular Aqueduct (EVA)

- *One of the most common inner ear malformations*
- *Hearing loss may be sudden or progressive*
- *Often associated with Pendred syndrome*

Audiologic Characteristics

Children with inner ear malformations typically present with:

- *Congenital or early onset SNHL*
- *Bilateral involvement in most cases*

- ***Progressive or fluctuating thresholds***
- ***Possible vestibular dysfunction***

Red Flag

Sudden hearing deterioration after minor head trauma is characteristic of EVA.

Audiologic Assessment

Audiology evaluation must include:

- ***Objective testing, ABR and OAE***
- ***Behavioral audiometry when possible***
- ***Speech perception testing***
- ***Vestibular screening when indicated***

Audiology must document:

- ***Stability or progression***
- ***Asymmetry***
- ***Functional hearing impact***

Role of Imaging

- ***High resolution CT of temporal bone***
- ***MRI for auditory nerve assessment***

Imaging is essential to:

- ***Confirm type of malformation***
- ***Assess cochlear nerve presence***
- ***Guide CI or ABI candidacy***

Audiology does not interpret imaging but must integrate findings into management planning.

Management Principles

Management depends on:

- ***Type of malformation***
- ***Degree of hearing loss***
- ***Auditory nerve integrity***
- ***Child's developmental status***

Hearing Rehabilitation Options

Conventional Hearing Aids

- ***Useful in mild to moderate losses***
- ***Limited benefit in severe malformations***

Cochlear Implantation

- ***Possible in many inner ear malformations***
- ***Outcomes depend on anatomy and nerve integrity***
- ***Higher surgical risk in some anomalies***

Auditory Brainstem Implant

- ***Indicated when cochlear nerve is absent***
- ***Considered in severe malformations or failed CI***
- ***Requires specialized centers***

Vestibular Considerations

- ***Vestibular dysfunction is common***
- ***May affect motor development***
- ***Referral for vestibular rehabilitation may be required***

Pediatric and Family Counseling

Families must be counseled regarding:

- ***Variability of hearing outcomes***
- ***Possible progression of hearing loss***
- ***Surgical risks and limitations***
- ***Long-term rehabilitation commitment***

Audiology must avoid guaranteeing outcomes.

Multidisciplinary Management

Optimal care requires:

- ***Audiology***
- ***ENT / Otology***
- ***Radiology***
- ***Genetics***
- ***Speech and language therapy***

Medico-Legal Considerations

Audiology documentation must:

- ***Clearly describe degree and progression of hearing loss***
- ***Recommend appropriate referrals***
- ***Document counseling and expectations***
- ***Avoid definitive prognostic statements***

Common Pitfall

Delaying implant referral in progressive malformations.

Key Points

- *Inner ear malformations are a major cause of congenital SNHL*
- *Imaging is essential for diagnosis and planning*
- *Hearing loss may be progressive or fluctuating*
- *CI is possible in many cases, ABI in selected cases*

Golden Rule

In inner ear malformations,
anatomy guides technology,
but rehabilitation determines outcome.

Balance Disorders in Children

Common Causes of Balance Disorders in Children

Peripheral Vestibular Causes

- *Vestibular neuritis*
- *Labyrinthitis*
- *Otitis media related vestibular symptoms*
- *Congenital inner ear malformations*
- *Ototoxicity*
- *Cochlear implant related vestibular dysfunction*

Central Causes

- *Migraine associated vertigo*
- *Seizure disorders*
- *Central nervous system lesions*
- *Developmental neurological disorders*

Systemic and Other Causes

- *Visual impairment*
- *Musculoskeletal disorders*
- *Metabolic disorders*
- *Psychological factors*

Clinical Presentation

Children with balance disorders may present with:

- *Delayed sitting or walking*
- *Poor balance during play*
- *Motion intolerance*
- *Difficulty with sports*
- *Nausea or vomiting*
- *Headaches*
- *Poor concentration*

Important

Absence of vertigo complaint does not exclude vestibular dysfunction.

History Taking in Pediatric Balance Disorders

Key history elements include:

- *Age at symptom onset*

- ***Developmental milestones***
- ***History of ear disease***
- ***Head trauma***
- ***Ototoxic drug exposure***
- ***Migraine history***
- ***Family history***

History must involve parents or caregivers.

Audiologic and Vestibular Assessment

Audiologic Evaluation

- ***Pure tone audiometry when age appropriate***
- ***Speech audiometry***
- ***Tympanometry***
- ***Otoacoustic emissions***

Audiology must evaluate hearing concurrently.

Vestibular Assessment

Test selection depends on age and cooperation:

- ***Vestibular evoked myogenic potentials***
- ***Video head impulse testing***
- ***Rotational chair testing***
- ***Postural and balance observation***

Formal caloric testing may be limited in young children.

Special Pediatric Conditions

Vestibular Migraine

- ***Common cause of episodic dizziness***
- ***Often associated with headache history***
- ***Normal audiologic findings***

Inner Ear Malformations

- ***Frequently associated with vestibular dysfunction***
- ***May present as delayed motor development***

Post Meningitis

- ***High risk of bilateral vestibular loss***
- ***May cause severe balance impairment***

Role of Audiology

Audiology responsibilities include:

- ***Identification of vestibular related hearing patterns***
- ***Selection of age appropriate tests***
- ***Documentation of functional impact***
- ***Referral for vestibular evaluation***
- ***Monitoring over time***

Audiology does not diagnose central neurologic disease.

Management Principles

Management depends on etiology and severity:

- ***Treatment of underlying otologic disease***
- ***Vestibular rehabilitation therapy***
- ***Developmental support***
- ***Educational accommodations***

Early intervention improves compensation and outcomes.

Vestibular Rehabilitation in Children

- *Adapted exercises for age*
- *Focus on balance, gaze stabilization, and coordination*
- *Requires family participation*
- *Often play based*

Multidisciplinary Approach

Effective management requires:

- *Audiology*
- *ENT*
- *Neurology*
- *Physical therapy*
- *Pediatrics*
- *Education services*

Key Points

- *Balance disorders in children are common and underdiagnosed*
- *Symptoms differ from adults*
- *Vestibular dysfunction affects development*
- *Early identification improves outcomes*

Golden Rule

In children, balance problems often speak
through development, not dizziness.

Pediatric Cochlear Implant Referral Decision

Audiologic Criteria

• Severe to profound bilateral SNHL• Limited benefit from hearing aids

Additional Evaluation

☐ Imaging

☐ Speech and language assessment

☐ Developmental evaluation

Decision

- ***CI candidate***
- ***Continue amplification and monitoring***

Post-Meningitis Hearing Assessment

Immediate Action

☐ Urgent audiologic evaluation

Results

- ***Normal→ Close follow-up***
- ***HL detected→ Imaging and CI fast-track consideration***

Evidence Anchor

Delayed assessment risks cochlear ossification.

Global Pediatric Otology Golden Rule

In children, evidence guides timing,

but delay costs development.

Final section

What Otology Expects from Audiology

What Otology Expects from Audiology

This Book was written with a clear intention.

Not to turn audiologists into otologists, and not to blur professional boundaries.

It was written to clarify responsibility, expectation, and clinical reality at the point where audiology decisions directly shape otologic outcomes.

Otology does not operate in isolation.

Every surgical decision, every medical escalation, and every reassurance begins with audiologic data and, more importantly, audiologic interpretation.

This chapter is not a summary of what you have read.

It is a statement of what comes next.

Otology and Audiology

Code of Ethic

1. Purpose

This Code of Ethics defines the professional, ethical, and legal standards governing practice in otology and audiology, with the primary goal of protecting patients, clinicians, and institutions.

Ethical practice is an essential component of safe and defensible clinical care.

2. Core Ethical Principles

All otologic and audiologic practice shall be guided by the following principles:

- ***Beneficence** – Act in the best interest of the patient*
- ***Non-maleficence** – Do no harm*
- ***Autonomy** – Respect patient choice and informed decision making*
- ***Justice** – Provide fair and equitable care*
- ***Integrity** – Maintain honesty and professional accountability*

3. Professional Competence

Clinicians must:

- *Practice within their defined scope*
- *Maintain professional knowledge and skills*
- *Use evidence-based methods*
- *Seek consultation or referral when appropriate*

No clinician shall provide services beyond their training, credentials, or institutional privileges.

4. Scope of Practice and Role Clarity

Audiology

Audiologists are responsible for:

- *Assessment of hearing and balance function*
- *Interpretation of audiologic test results*

- Rehabilitation and counseling
- Referral for medical evaluation when indicated

Audiologists **do not provide medical or surgical diagnoses**.

Otology (ENT)

Otologists are responsible for:

- Medical diagnosis
- Medical and surgical management
- Interpretation of imaging
- Informed consent for medical or surgical interventions

5. Patient-Centered Care

Clinicians must:

- Respect patient dignity and privacy
- Communicate clearly and honestly
- Provide information in understandable language
- Avoid coercion or undue influence

Patient welfare takes precedence over institutional, financial, or personal interests.

6. Informed Consent and Transparency

- Consent is a process, not a signature
- Risks, benefits, and limitations must be explained
- Alternatives, including no intervention, must be discussed
- Consent must be documented

Incomplete disclosure is ethically unacceptable.

7. Confidentiality and Data Protection

Clinicians must:

- *Protect patient confidentiality*
- *Secure audiologic and medical records*
- *Share information only on a need-to-know basis*
- *Follow institutional and legal data protection policies*

Audiograms, reports, and recordings are legal medical records.

8. Conflict of Interest

Clinicians must:

- *Avoid financial or personal conflicts influencing care*
- *Disclose any real or perceived conflict*
- *Separate diagnosis from device sales*
- *Document patient choice and alternatives*

Ethical practice requires transparency.

9. Professional Integrity and Honesty

Clinicians must:

- *Provide accurate information*
- *Avoid guarantees or unrealistic promises*
- *Acknowledge uncertainty and limitations*
- *Correct errors promptly and transparently*

Misrepresentation is unethical and legally risky.

10. Interprofessional Collaboration

- *Respect the roles of other professionals*
- *Communicate clearly and promptly*

- *Document referrals and shared decisions*
- *Support coordinated patient care*

Breakdown in communication is an ethical failure.

11. Research, Teaching, and Publication

- *Obtain appropriate consent*
- *Protect patient identity*
- *Follow ethical review processes*
- *Avoid data manipulation or plagiarism*

Clinical education must never compromise patient rights.

12. Advertising and Professional Representation

Clinicians must:

- *Avoid misleading claims*
- *Represent qualifications accurately*
- *Comply with institutional and regulatory advertising policies*

Ethical marketing reflects professional integrity.

13. Reporting Ethical Concerns

Clinicians have an ethical obligation to:

- *Report unsafe or unethical practices*
- *Protect patient safety*
- *Follow institutional reporting pathways*

Silence in the presence of harm is unethical.

14. Enforcement and Accountability

Violation of this Code may result in:

- *Institutional disciplinary action*

- Professional regulatory consequences
- Legal liability

Ethical accountability is a professional duty.

Ethical Key Statements

- Ethics is preventive medicine in medico-legal practice
- Scope boundaries protect patients and clinicians
- Transparency builds trust and reduces litigation

Final Ethical Principle

Clinical excellence without ethics is unsafe.

Ethics without documentation is defenseless.

Professional integrity protects everyone.

Audiology vs ENT

Liability Comparison Table

Aspect	Audiology Liability	ENT Liability
Scope of responsibility	Functional assessment and rehabilitation	Medical diagnosis and surgical management
Legal duty	Accurate testing, interpretation, referral	Diagnosis, treatment, surgery, and follow up
Common allegations	Missed SSNHL, failure to refer unilateral SNHL, delayed post meningitis screening	Delayed diagnosis, surgical complications, inadequate follow up

Aspect	*Audiology Liability*	*ENT Liability*
	Does not provide medical diagnosis	*Provides definitive medical diagnosis*
High risk conditions	*SSNHL, unilateral SNHL, pediatric hearing loss, CI candidacy*	*Cholesteatoma, acoustic neuroma, otologic tumors, complications*
Time sensitive liability	*Delay in referral or escalation*	*Delay in treatment or surgery*
Documentation focus	*Test selection, interpretation, counseling, referral*	*Diagnosis, treatment plan, consent, operative notes*
Consent responsibility	*Test limitations and rehabilitation counseling*	*Medical and surgical informed consent*
Error of omission risk	*Very high, failure to refer, failure to document*	*Moderate, failure to treat or follow up*
Error of commission risk	*Moderate, acting beyond scope*	*High, surgical or invasive errors*
Device related liability	*Hearing aid and CI counseling, conflict of interest*	*Surgical device selection and implantation*
Medico legal exposure trend	*Increasing with expanded scope*	*Traditionally high but better defined*
Legal defense strength	*Documentation and timely referral*	*Documentation, consent, and standard of care adherence*

1. What Otology Assumes Audiology Already Knows

Otologists often assume that audiology:

- ***Understands the difference between measurement and meaning***
- ***Recognizes when results are reliable and when they are not***
- ***Can distinguish cochlear pathology from neural dysfunction***
- ***Identifies red flags early, not after complications occur***
- ***Knows when silence is dangerous and reassurance is harmful***

These assumptions are not always correct.

This volume was written to close that gap.

2. What Otology Needs from Audiology in Real Practice

Otology does not need more numbers.

It needs clarity.

It needs audiology to answer questions such as:

- ***Is this hearing loss functional or structural***
- ***Is poor speech due to audibility or neural synchrony***
- ***Is this asymmetry clinically meaningful***
- ***Is this child safe to wait or does time matter***
- ***Is this patient a surgical candidate or a rehabilitation candidate***

When audiology hesitates to interpret, otology is forced to guess.

Guessing is where risk begins.

3. Where Miscommunication Commonly Occurs

Miscommunication between audiology and otology rarely comes from lack of testing.

It comes from lack of context.

Common failure points include:

- Reporting thresholds without functional interpretation
- Ignoring speech performance discrepancies
- Underestimating mild or unilateral losses
- Delaying referral while waiting for “clearer” results
- Using technically correct but clinically misleading language

A correct test does not guarantee a correct decision.

4. Where Medico-Legal Risk Truly Starts

In most legal cases involving hearing care, the problem is not the equipment.

It is not calibration.

It is not even test selection.

It is interpretation.

Statements such as:

- “Findings are within normal limits”
- “Consistent with age”
- “No significant hearing loss”

can become legally dangerous when context is ignored.

The test may be correct.

The conclusion may not be defensible.

5. Audiology as a Clinical Guide, Not a Technical Service

Audiology is often viewed as a supportive service.

In reality, it is a guiding discipline.

Audiology determines:

- ***Who is reassured***
- ***Who is referred***
- ***Who waits***
- ***Who acts***

Otology acts on what audiology frames.

This is not power.

It is responsibility.

6. The Central Message of This Volume

Otology for Audiology was written to establish one principle:

Audiology does not support otology.

Audiology guides otology.

Guidance requires:

- ***Evidence awareness***
- ***Clear communication***
- ***Ethical restraint***
- ***Precise documentation***

Anything less exposes patients and professionals to avoidable harm.

A Letter to the Audiologist

This book was not written to impress you.
It was written to protect you.
It was written for the moment when an otologist asks,
“Are you sure?”
It was written for the moment when a family asks,
“Is it safe to wait?”
It was written for the report that will be read years later,
not just today.
If this book helped you pause before reassuring,
question before concluding,
and document before deciding,
then it has done its job.

Otology and audiology are not parallel disciplines.

They are sequential.



Audiology thinks first.

Otology acts next.

audiology must understand otology to guide decisions safely and responsibly.

Otology and Audiology

Speak One Language

“Two Specialties. One Patient. One Language.”

Sincerely

Dr. Yahia Al Ahmari

Otology, Neurotology & Cochlear Implant Leader

Dr. Hany Abdou

Audio-Vestibular Medicine

Otology for Audiology

Otology and Audiology

Speak One Language

“Two Specialties. One Patient. One Language.”

* * *

Dr. Yahia Al Ahmari

Otology, Neurotology & Cochlear Implant Leader

Dr. Hany Abdou

Audio-Vestibular Medicine